

**ISOFLAVONE EFFECTS ON
INSULIN LIKE GROWTH FACTORS AND
BREAST CANCER**

by

Maeli Campbell-McNulty

A Thesis Submitted for the Degree of

Doctor of Philosophy

of the University of London

Department of Surgery

University College London

September 2007

UMI Number: U591434

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI U591434

Published by ProQuest LLC 2013. Copyright in the Dissertation held by the Author.
Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against
unauthorized copying under Title 17, United States Code.



ProQuest LLC
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106-1346

SIGNED DECLARATION

I, Maeli Campbell-McNulty, confirm that the work presented in this thesis is my own. Where information has been derived from other sources, I confirm that this has been indicated in the thesis.

Signature:

Date: 14TH MARCH 2008

ABSTRACT

An increased concentration of insulin-like growth factor-1 (IGF-1) is recognized to be an independent risk factor for pre-menopausal breast cancer. Tamoxifen is thought to initially reduce IGF-1 concentrations and increase levels of the IGF binding proteins 1 and 3. Isoflavones are oestrogen-like plant compounds, which may, because of their structural similarities to tamoxifen, also alter IGF status.

This thesis first compares IGF-1, IGF binding protein 1 (BP-1) and IGF binding protein 3 (BP-3) levels in breast cancer patients (n=14) compared with control subjects (n=23) and then assesses the effect of tamoxifen on IGF status after 9, 18 and 27 months of treatment. Concentrations of IGF-1, BP-1 and BP-3 at baseline did not differ between cases and controls. However on tamoxifen treatment, BP-1 and BP-3 were both significantly increased after 18 and 27 months while the IGF-1/BP-3 ratio was significantly decreased after 9 and 18 months.

A feasibility study was then conducted to compare the effects of acute (single 80 mg load) versus chronic (80 mg/day for 7 days) administration of isoflavone-containing soy and linseed cereal bars on IGF-1, IGFBP-1 and IGFBP-3 in healthy female volunteers (n=10). Assays were established for the analyses of serum IGF-1, BP1 and BP3 concentrations and GCMS analysis of isoflavones in urine. Concentrations of IGF-1 were unchanged following the single 80 mg load. However, IGF-1 and BP-3 concentrations were significantly elevated after a week of supplementation with the soy and linseed bar.

A larger study was then carried out to assess the effect of one-month isoflavone supplementation (80 mg/d) in tablet form, on IGF status in healthy pre- (n=16) and post-menopausal (n=7) women. This was a randomized, placebo-controlled

crossover study with a minimum two-month washout period. Hormonal, antioxidant and lipid altering effects of the supplement were also examined, as was background diet. For pre-menopausal subjects, while there was an effect of time on IGF-1 ($p=0.005$), BP-1 ($p=0.004$), and BP-3 ($p<0.001$) confirming that IGF profile is influenced by menstrual cycle, this did not differ between placebo and isoflavone supplement. When the change in IGF-1 over the whole supplementation period was compared between the supplement and placebo phases, there was a non-significant reduction in change in IGF-1 ($p=0.06$) on isoflavone supplement compared to placebo. However, this may have been due to the non-significantly higher baseline concentrations of IGF-1 during the supplement phase. In post-menopausal subjects, there was no effect of isoflavone supplementation in comparison with placebo on IGF-1, BP-1 or BP-3. Finally, experiments were done *in vitro* to study the effect of isoflavones on DNA synthesis and proliferation in ER α positive MCF-7 and ER α negative MDA-MB 231 human breast cancer cells. In MCF-7 cells, low concentrations of isoflavones (0.1 μ M-10 μ M) stimulated DNA synthesis while high concentrations of genistein and equol were able to inhibit DNA synthesis with IC₅₀ values of 93 μ M and 55 μ M respectively compared with 1.17 μ M tamoxifen. In MDA-MB231 cells the Isoflavones did not stimulate DNA synthesis at any concentrations but significantly inhibited DNA synthesis with IC₅₀ values of 114 μ M, 14.1 μ M and 14.55 μ M for daidzein, genistein and equol respectively compared with 1.35 μ M tamoxifen.

This work suggests that isoflavone supplementation may alter IGF profile in pre-menopausal women, and could suggest a role for these dietary compounds in breast cancer prevention. This should be further investigated in long term intervention studies with isolated isoflavone supplements.

For my parents

PUBLICATIONS ARISING FROM WORK CONTRIBUTING TO THIS THESIS

Campbell, M.J., Woodside, J.V., Secker-Walker, J., Titcomb, A. & Leathem, A.J. (2001). IGF status is altered by tamoxifen in patients with breast cancer. *Mol Pathol*, **54**, 307-10.

Campbell, M.J., Woodside, J.V., Honour, J.W., Morton, M.S. & Leathem, A.J. (2004). Effect of red clover-derived isoflavone supplementation on insulin-like growth factor, lipid and antioxidant status in healthy female volunteers: a pilot study. *Eur J Clin Nutr*, **58**, 173-9.

Woodside, J.V. & Campbell, M.J. (2001). Isoflavones and breast cancer. *J Br Menopause Soc*, **7**, S17-S21.

Woodside, J.V., Campbell, M.J., Denholm, E.E., Newton, L., Honour, J.W., Morton, M.S., Young, I.S. & Leathem, A.J. (2006). Short-term phytoestrogen supplementation alters insulin-like growth factor profile but not lipid or antioxidant status. *J Nutr Biochem*, **17**, 211-5.

ACKNOWLEDGEMENTS

I would like to thank my daughter Grace who gave me the opportunity to finish writing, my mother and father for helping look after her and my husband Keith for his support and understanding throughout.

Many thanks to Dr Jayne Woodside for her invaluable scientific guidance, support and friendship. Ann Titcomb for her help and support and Dr Anthony Leathem for his supervision and for giving me the opportunity to pursue this research. Thanks to Dr Hazel Welch for her help with tissue culture and the Ludwig Institute for donating the breast cancer cell lines. Thanks to Brian Rafferty for his supervision with assay development at NIBS. My acknowledgement also goes to my colleagues, Jayne Woodside and Erica Denholm in the Breast Cancer Research Group, Department of Surgery, UCL. The phytoestrogen, antioxidant, lipid analyses and Comet assays were conducted by them.

I would also like to thank my friends Lisa Leeton and Devon Wilson for their practical help at the University of Queensland and Kim Jackson for invigilating the viva. I would like to thank Novogen (North Ryde, Sydney, Australia) for kindly supplying the supplements and Action Against Breast Cancer for their financial support.

Most of all I would like to thank my heavenly Father who makes all things possible.

CONTENTS

<u>SIGNED DECLARATION</u>	<u>2</u>
<u>ABSTRACT</u>	<u>3</u>
<u>PUBLICATIONS ARISING FROM WORK CONTRIBUTING TO THIS THESIS 6</u>	
<u>ACKNOWLEDGEMENTS</u>	<u>7</u>
<u>CONTENTS</u>	<u>8</u>
<u>FIGURES</u>	<u>14</u>
<u>TABLES</u>	<u>15</u>
<u>CHAPTER ONE – INTRODUCTION</u>	<u>18</u>
1.1 ROLE OF DIETARY FACTORS	18
1.2 ISOFLAVONES.....	20
1.2.1 <i>IN VITRO</i> EVIDENCE.....	22
1.2.2 ANIMAL EVIDENCE	24
1.2.3 EPIDEMIOLOGICAL EVIDENCE	25
1.2.3.1 Soy/Isoflavone intake	25
1.2.3.2 Hormone levels and breast cancer risk	30
1.2.4 INTERVENTION STUDIES EXAMINING EFFECT OF ISOFLAVONE SUPPLEMENTATION ON HORMONE LEVELS	31

1.3	INSULIN LIKE GROWTH FACTOR -1 AND BINDING PROTEINS	34
1.3.1	IGF-1 AND BREAST CANCER RISK	35
1.3.2	IGF-1 AND OESTRADIOL	38
1.3.3	IGF-1 AND TAMOXIFEN	38
1.3.4	IGF-1 AND DIET	39
1.3.5	IGF-1 AND ISOFLAVONE/SOY INTAKE	40
1.4	CONCLUSION.....	40
 <u>CHAPTER TWO – METHODS</u>		<u>51</u>
2.1	COLLECTION AND PREPARATION OF BIOLOGICAL SAMPLES	51
2.2	ENDPOINT MEASUREMENTS	51
2.2.1	DIET	51
2.2.2	PHYTOESTROGENS	51
2.2.3	ASCORBIC ACID	52
2.2.4	RETINOL, α -TOCOPHEROL AND CAROTENOIDS	53
2.2.5	SERUM LIPIDS AND LIPID PEROXIDATION.....	54
2.2.6	COMET ASSAY	55
2.2.7	CREATININE.....	55
2.2.8	HORMONES	55
2.2.9	IGF-1, BP-1 AND BP-3	57
2.3	DEVELOPMENT OF AN ENZYME LINKED IMMUNOSORBANT ASSAY (ELISA) FOR THE QUANTITATIVE MEASUREMENT OF IGF-BP3 IN HUMAN SERUM.....	59
2.3.1	INTRODUCTION	59
2.3.2	METHOD	59
2.3.3	RESULTS.....	60
2.3.4	DISCUSSION	77

2.4 CELL CULTURE	79
2.5 BRDU ASSAY	79
2.5.1 BRDU INCORPORATION	80
2.5.2 OPTIMAL CONDITIONS.....	81

CHAPTER THREE – THE EFFECT OF TAMOXIFEN ON IGF STATUS IN BREAST CANCER PATIENTS – A CASE CONTROL STUDY.....83

3.1 INTRODUCTION	83
3.2 MATERIALS AND METHODS	85
3.2.1 LABORATORY METHODS.....	85
3.2.2 STATISTICAL METHODS	86
3.3 RESULTS	86
3.4 DISCUSSION	92
3.5 CONCLUSION.....	96

CHAPTER FOUR – SHORT-TERM PHYTOESTROGEN SUPPLEMENTATION IN HEALTHY FEMALE VOLUNTEERS – PILOT STUDY.....97

4.1 INTRODUCTION	97
4.2 MATERIALS AND METHODS	98
4.2.1 STUDY DESIGN.....	98
4.2.2 PHASE 1	100
4.2.3 PHASE 2	101
4.2.4 COLLECTION AND PREPARATION OF BIOLOGICAL SAMPLES	101
4.2.5 ENDPOINT MEASUREMENTS	101
4.2.6 STATISTICAL METHODS.....	102
4.3 RESULTS	102
4.4 DISCUSSION	107

**CHAPTER FIVE – THE EFFECT OF ISOFLAVONE SUPPLEMENTATION ON
INSULIN-LIKE GROWTH FACTOR, LIPID AND ANTIOXIDANT STATUS IN
HEALTHY FEMALE VOLUNTEERS.....115**

5.1	INTRODUCTION	115
5.2	MATERIALS AND METHODS	116
5.2.1	STUDY DESIGN.....	116
5.2.2	ELIGIBILITY CRITERIA	117
5.2.3	SAMPLE COLLECTION	117
5.2.4	ENDPOINT MEASUREMENTS	118
5.2.5	SUPPLEMENT.....	119
5.2.6	STATISTICAL ANALYSES	119
5.3	RESULTS	120
5.3.1	ISOFLAVONES	121
5.3.2	HORMONES	128
5.3.3	INSULIN-LIKE GROWTH FACTORS.....	134
5.3.4	LIPIDS AND ANTIOXIDANTS	139
5.3.5	NUTRIENTS.....	144
5.3.6	ASSOCIATIONS BETWEEN CONTINUOUS VARIABLES	148
5.3.6.1	Associations with isoflavone status	148
5.3.6.2	Associations with hormonal status	149
5.3.6.3	Associations with IGF-1 and BPs	150
5.3.6.4	Associations with lipid and antioxidant status.....	151
5.4	DISCUSSION	152
5.4.1	ISOFLAVONES	152
5.4.2	EQUOL METABOLISM AND DIET.....	153
5.4.3	ISOFLAVONES AND HORMONES	156

5.4.4	IGF-1 AND BINDING PROTEIN CONCENTRATIONS	161
5.4.5	IGF AND MENSTRUAL CYCLE	162
5.4.6	ISOFLAVONES AND IGF	164
5.4.7	ISOFLAVONES AND LIPIDS	172
5.4.8	ISOFLAVONES AND ANTIOXIDANTS	173
5.4.9	DIET	174
5.5	CONCLUSION	175

CHAPTER SIX – THE EFFECT OF ISOFLAVONES ON DNA SYNTHESIS IN MCF-7 AND MDA-MB-231 HUMAN BREAST CANCER CELLS 176

6.1	INTRODUCTION	176
6.2	MATERIALS AND METHODS	182
6.2.1	CELL CULTURE	182
6.2.2	BRDU ASSAY	182
6.3	RESULTS	183
6.3.1	EFFECTS OF E ₂ AND IGF-1 ON DNA SYNTHESIS	183
6.3.2	EFFECTS OF ISOFLAVONES AND TAMOXIFEN ON DNA SYNTHESIS	183
6.3.3	INTERACTIONS BETWEEN ISOFLAVONES AND TAMOXIFEN WITH OESTRADIOL	188
6.3.4	INTERACTIONS BETWEEN ISOFLAVONES WITH TAMOXIFEN	188
6.3.5	INTERACTIONS BETWEEN ISOFLAVONES AND TAMOXIFEN WITH IGF-1	188
6.4	DISCUSSION	191
6.4.1	ISOFLAVONES	191
6.4.2	INTERACTIONS BETWEEN ISOFLAVONES AND OESTRADIOL	194
6.4.3	INTERACTIONS BETWEEN OESTRADIOL AND ISOFLAVONES WITH TAMOXIFEN	196
6.4.4	INTERACTIONS BETWEEN ISOFLAVONES AND TAMOXIFEN WITH IGF-1	198
6.5	CONCLUSION	201

CHAPTER SEVEN – CONCLUSION	203
APPENDIX.....	206
REFERENCES	246

FIGURES

Figure 1.2.1: Chemical structure of isoflavones and oestradiol.....	21
Figure 3.3.1: Effect of tamoxifen supplementation on insulin-like growth factor binding protein 1 (IGFBP-1).....	88
Figure 3.3.2: Effect of tamoxifen supplementation on insulin-like growth factor binding protein 3 (IGFBP-3).....	89
Figure 3.3.3: Effect of tamoxifen supplementation on insulin-like growth factor 1 (IGF-1).....	90
Figure 3.3.4: Effect of tamoxifen supplementation on insulin-like growth factor 1 (IGF-1) to IGF binding protein 3 (IGFBP-3) ratio.....	91
Figure 4.3.1: Association between change in IGF-1 and IGFBP-3 over 1 week of supplementation ($r=0.721$, $P<0.05$; Spearman correlation coefficient). ...	106
Figure 5.2.1 Study design – for pre-menopausal women blood samples were taken throughout the menstrual cycle as shown. For post-menopausal women, fasting blood and urine were only collected at the start and end of the 28 day intervention period.....	117
Figure 5.3.1.1 Effect of supplementation on urinary excretion of isoflavone metabolites in pre-menopausal subjects.....	125
Figure 5.3.1.2. Effect of supplement on urinary excretion of isoflavone metabolites in post-menopausal subjects.	125
Figure 5.3.2.1: Effect of supplementation on LH levels in pre-menopausal women. Mean (SE).....	129
Figure 5.3.2.2: Effect of supplementation on FSH levels in pre-menopausal women. Mean (SE).....	129
Figure 5.3.2.3: Effect of supplementation on Oestradiol levels in pre-menopausal women. Mean (SE).....	130
Figure 5.3.2.4: Effect of supplementation on Progesterone levels in pre-menopausal women. Mean (SE).....	130
Figure 5.3.2.5: Effect of supplementation on SHBG levels in pre-menopausal women. Mean (SE). * $P=0.003$ paired t test supplement vs placebo (Day 21-23).	131
Figure 5.3.3.1: IGF-1 levels in pre-menopausal women on placebo or supplement. Mean (SE).....	137
Figure 5.3.3.2: IGFBP-3 levels in pre-menopausal women on placebo or supplement. Mean(SE).....	137
Figure 5.3.3.3: IGFBP-1 levels in pre-menopausal women on placebo or supplement. Mean (SE).....	138
Figure 5.3.4.1:HDL-cholesterol (mmol/L) in pre and post-menopausal women after supplementation. Mean(SE).	139
Figure 6.3.1.2: Dose response of MDA-MB-231 cells incubated with IGF-1 and oestradiol for 24 hours before labelling with BrDU. Mean (SE).....	186
Figure 6.3.2.1: Dose response of MCF-7 cells incubated with isoflavones and Tamoxifen for 24 hours before labelling with BrDU. Mean (SE).....	187
Figure 6.3.2.2: Dose response of MDA-MB-231 cells incubated with isoflavones and Tamoxifen for 24 hours before labelling with BrDU. Mean (SE).....	187

TABLES

Table 1.2.3.1: Studies on the association between phyto-oestrogens in plasma and urine or phyto-oestrogen-rich diet and breast cancer risk.....	42
Table 1.2.4.1: Interventions with soy foods/supplements in premenopausal women-effects on hormones.....	43
Table 1.2.4.2: Interventions with soy foods/supplements in postmenopausal women- effects on hormones.....	44
Table 1.3.4.1: Epidemiological studies- Dietary correlations with IGF-1, BP3 and BP1 concentrations.....	45
Table 1.3.5.1: Interventions with soy foods or supplements – effects on insulin like growth factors and binding proteins.....	49
Table 3.3.1: Baseline concentrations of insulin-like growth factor 1 (IGF-1), IGF binding protein 1 (IGFBP-1), and IGFBP-3 in patients and controls	87
Table 3.3.2: Patient concentrations of insulin-like growth factor 1 (IGF-1), IGF binding protein 1 (IGFBP-1), and IGFBP-3 at baseline and during tamoxifen treatment.....	87
Table 4.3.1: Effect of one phytoestrogen meal on urinary phytoestrogen excretion- Phase 1	104
Table 4.3.2: Effect of 1 week of phytoestrogen supplementation on urinary phytoestrogen excretion – Phase 2	104
Table 5.3.1 Baseline characteristics of study subjects	120
Table 5.3.1.1. Effect of supplementation on urinary excretion of isoflavone metabolites in pre-menopausal subjects.....	123
Table 5.3.1.2. Effect of supplementation on urinary excretion of isoflavone metabolites in post-menopausal subjects.	124
Table 5.3.1.3. Comparison of Isoflavone excretion with other studies	126
Table 5.3.1.4: Comparison of serum lipids, tocopherols and IGF-1, BP-3 and BP-1 concentrations in equol excreters and non excreters.....	127
Table 5.3.2.1: Effect of isoflavone supplementation on circulating hormone levels in pre-menopausal women. Mean \pm (SD).	132
Table 5.3.2.2: Effect of isoflavone supplementation on circulating hormone levels in post-menopausal women. Mean \pm (SD).....	132
Table 5.3.2.3: Comparison of IGFs and hormone levels measured by menopausal status.....	133
Table 5.3.3.1: Effect of supplementation on insulin-like growth factor and binding protein concentrations in pre-menopausal women	135
Table 5.3.3.2: Effect of supplementation on insulin-like growth factor and binding protein concentrations in post-menopausal women	136
Table 5.3.4.1: Effect of isoflavone supplementation on antioxidant and lipid status in pre-menopausal subjects	140
Table 5.3.4.2: Effect of isoflavone supplementation on antioxidant and lipid status in post-menopausal subjects.....	142
Table 5.3.5.1: Dietary intake, on placebo and supplement in pre and post-menopausal women. Mean (SD)	146
Table 6.3.1.1: Dose response of MCF-7 cells incubated with Isoflavones, Tamoxifen, IGF-1 and oestradiol for 24 hours.	185
Table 6.3.1.2: Dose response of MDA-231 cells incubated with Isoflavones, Tamoxifen, IGF-1 and oestradiol for 24 hours.	185
Table 6.3.3.1 Response of MCF-7 cells incubated for 24 hours with different combinations of Isoflavones, oestradiol, Tamoxifen and IGF-1.	190

Table 6.3.3.2: Response of MDA231 cells incubated for 48 hours with different combinations of Isoflavones, oestradiol, Tamoxifen and IGF-1.	190
Table 5.A.1: Correlations of equol excretion with serum lipids, serum antioxidants, dietary intake (pre and post-menopausal women) and anthropometric measurements.	206
Table 5.A.2: Correlations of Enterolactone excretion with serum lipids, serum antioxidants and dietary intake (pre and postmenopausal women).	208
Table 5.A.3: Correlations of Enterodiol excretion with serum lipids, serum antioxidants and dietary intake (pre and postmenopausal women)	209
Table 5.A.4: Correlations of Genistein excretion with serum lipids, serum antioxidants and dietary intake (pre and post-menopausal women)	211
Table 5.A.5: Correlations of Daidzein excretion with serum lipids, serum antioxidants, dietary intake and anthropometric measurements (pre and postmenopausal women).....	213
Table 5.A.6: Correlations of serum concentrations of Isoflavones (ug/day) in Pre-menopausal women with Serum IGF-1, BP-3 and BP-1 throughout the menstrual cycle.....	216
Table 5.A.7: Correlations of serum concentrations of Enterolactone (ug/day) in pre-menopausal women with Serum IGF-1, BP-3 and BP-1 throughout the menstrual cycle.....	217
Table 5.A.8: Correlations of serum concentrations of Isoflavones (ug/day) in post-menopausal women with Serum IGF-1, BP-3 and BP-1.....	218
Table 5.A.9: Correlations of hormones with excretion of Isoflavones (ug/day) in post-menopausal women.....	219
Table 5.A.10: Correlations of hormones with excretion of Isoflavones (ug/day) in pre-menopausal women.	221
Table 5.A.11: Correlations of Estradiol and Progesterone with IGF-1, BP-3 and BP-1 in pre-menopausal women (n=16).	224
Table 5.A.12: Correlations of SHBG with IGF-1 and BP-1 in pre-menopausal women (n=16).....	225
Table 5.A.13: Correlations of hormones with IGF-1, BP-1 and BP-3 in post - menopausal women (n=7).	226
Table 5.A.14: Correlations of serum concentrations of IGF-1 & BP-1 and Serum Lipids (no correlations seen with BP-3) in pre-menopausal women.	227
Table 5.A.15: Correlations of serum concentrations of IGF1 with Serum antioxidants in pre menopausal women.....	228
Table 5.A.16: Correlations of serum concentrations of BP1 and Serum antioxidants in premenopausal women.....	229
Table 5.A.17: Correlations of serum concentrations of BP-3 and Serum antioxidants in pre-menopausal women.	230
Table 5.A.18: Correlations of serum concentrations of IGF-1, & BP-1 with anthropometric data in premenopausal women.	232
Table 5.A.19:Correlations of BP-1 with serum lipids in post-menopausal women.	233
Table 5.A.20: Correlations of IGF-1, BP-3 and BP-1 with serum antioxidants in postmenopausal women.	233
Table 5.A.21: Correlations of Serum Lipid concentrations and FFQ/FD dietary intake in pre and postmenopausal women (n=23).	234
Table 5.A.22: Correlations of Serum concentrations of antioxidants with FFQ/FD intake in pre and postmenopausal women (n=23). Serum vitamin C vs dietary vitamin C, Serum α and γ tocopherol vs dietary Vitamin E and serum α and β carotene vs dietary Carotene.....	235

Table 5.A.23: Correlations of IGF-1 concentrations with FFQ/FD intake in pre-menopausal women.....	237
Table 5.A.24: Correlations of BP-1 concentrations with FFQ/FD intake in pre-menopausal women.....	238
Table 5.A.25: Correlations of BP-3 concentrations with FFQ/FD intake in pre-menopausal women.....	239
Table 5.A.26: Correlations of IGF-1 with FFQ/FD in post-menopausal women	242
Table 5.A.27: Correlations of BP-3 and BP-1 with FFQ/FD in post-menopausal women	244

CHAPTER ONE – INTRODUCTION

Breast cancer is one of the most important public health problems facing women in the UK and other Western populations today. With over 40,000 new cases diagnosed in the UK in 2000 alone, the disease accounts for 17% of female cancer mortality with almost 13,000 deaths in UK each year (Toms, 2004). Incidence rates have increased by 70% since 1971 with one in ten women developing breast cancer at some point in their lives (ONS, 31 October 2003). Breast cancer is a complex, multifactorial disease, but established risk factors include nulliparity, late first pregnancy, early menarche, late menopause, oral contraceptive use, hormone replacement therapy (HRT), obesity, increased alcohol consumption, exposure to ionising radiation and inheritance of mutations in the BRCA1 and BRCA2 genes (McPherson et al., 2000).

1.1 Role of dietary factors

There is strong epidemiological evidence that environmental factors such as diet have a role in breast cancer aetiology (Miller et al., 1994). This evidence initially came from population and migrant studies where Chinese, Japanese or Filipino's who had lived in the United States for a decade or longer had a risk 80% higher than more recent migrants (Ziegler et al., 1993). The timing of immigration to the US was also important in determining breast cancer risk and when migration occurred later in life, rates for breast cancer were substantially lower than when migration occurred early, although they were still much higher than in the homeland populations (Hunter & Willett, 1996; Shimizu et al., 1991).


Subsequently, cohort and case-control studies have been carried out, mainly focussing on the hypothesis that a diet rich in fat can increase the risk of breast cancer (Hunter & Willett, 1996). Assessing this relationship is complicated and study design, dietary methodology and choice of study population is important (Miller et al., 1994). The results of prospective studies have largely been equivocal (Hunter & Willett, 1996), while fat-reducing intervention studies have shown some effects on total and free oestradiol in post-menopausal but not pre-menopausal women (Bingham et al., 1998). Other studies suggest that olive oil may be protective for its high levels of monounsaturated fatty acids and/or phenolic compounds (Hashim et al., 2005). However recent findings from large prospective studies in the UK and Europe (EPIC) suggest that risk of breast cancer is associated with intake of saturated fat (HR 1.22, 95% CI 1.06-1.4), with 35g /day vs 10g/day approximately doubling risk (Bingham et al., 2003) in peri and post-menopausal women studied. While total or specific fruit and vegetable intake was not found to be associated with risk of breast cancer in pre or postmenopausal women in the initial analysis of data in women across 10 countries in Europe (van Gils et al., 2005), the UK women's cohort study (n= 35,792) has found that for pre-menopausal women, total fibre is significantly protective against breast cancer, especially fibre from cereals and possibly fruit (Cade et al., 2007), while high consumption of meat, especially processed meat, significantly increased risk of breast cancer (Taylor et al., 2007). It is possible that specific components of dietary fibre such as lignans present in seeds and wholegrains are responsible for reducing risk of breast cancer (McCann et al., 2004). There is also interest in the protective effect of other dietary factors, such as vitamins, other micronutrients and phytochemicals (Bingham et al., 1998).

1.2 Isoflavones


Plants contain a large number of potential antioxidant/anticancer type compounds, known as phytoestrogens (Bingham et al., 1998), many of which can be classed as phenols, as they contain an active benzene ring with a hydroxyl group attached. An example of such compounds is the flavonoid family which are present in fruit and vegetables as the biologically inactive phenolic glycosides. The aglycone (sugar free) part of the compound is characterised by a C6-C3-C6 backbone. Sub groups such as Isoflavones and Flavanols have been classified to encompass differences in the position of ring B and varying degrees of oxidation and saturation of the pyrone ring (**Figure 1.2.1**).

Isoflavones are found predominantly in legumes (peas, beans, lentils, etc). Soybeans and soy products are perhaps the most common food source of isoflavones and contain large amounts of the isoflavones genistein and daidzein (Coward et al., 1993). But other legumes, especially clovers, contain higher total levels and, in addition to genistein and daidzein, also contain formononetin and biochanin (Francis et al., 1967). Interest has focused on isoflavones and breast cancer due to the relatively low breast cancer mortality rates in Asian countries where soy foods are commonly consumed. In Japan and China, for example, the breast cancer mortality rate is about one quarter of that in the United Kingdom (Society, 2006). In addition, a lower proportion of breast biopsies from Asian women contain hyperplasia, atypical hyperplasia, and apocrine metaplasia than breast biopsies from American women (Sasano et al., 1975; Schuerch et al., 1982). The evidence linking isoflavone or soy intake with breast cancer risk comes from a number of sources and these will be considered in turn.

Genistein and Biochanin A



Daidzein and Fomononetin



Oestradiol-17 β




Figure 1.2.1: Chemical structure of isoflavones and oestradiol.

Basic structure comprised of two benzene rings (A and B) linked through a heterocyclic pyran or pyrone (with a double bond) ring (c) in the middle

1.2.1 *In vitro* evidence

Isoflavones have been considered weak oestrogenic compounds *in vitro*, binding to both oestrogen receptors α and β with a relative binding affinity of 4 and 87 respectively for genistein (compared to 100 for oestradiol) (Kuiper et al., 1997; Kuiper et al., 1998). Numerous studies in breast cancer cells show that Isoflavones stimulate growth at low concentrations but inhibit both hormone dependent and independent cells at high concentrations with genistein IC₅₀ between 5 and 40 μ M (Messina, 1999) (Magee & Rowland, 2004). Studies suggest that these biphasic effects are a result of estrogen dependent and independent mechanisms (Chen et al., 2003) (Chen et al., 2004).

Apart from anti-oestrogenic activity the mechanism whereby isoflavones reduce cell growth, particularly in oestrogen receptor-negative cells, is still to be fully elucidated. Other possible mechanisms include: antioxidant properties observed *in vitro*, and stimulating several antioxidative enzymes like glutathione reductase and catalase (Ruiz-Larrea et al., 1997; Sierens et al., 2001; Sierens et al., 2002; Wei et al., 1993), inhibitory actions on several enzymes including aromatase and those involved in signal transduction, like tyrosine protein kinases (Akiyama et al., 1987), MAP kinase (Thorburn & Thorburn, 1994) and ribosomal S6 kinase (Linassier et al., 1990). Genistein has been shown to inhibit the activity of DNA topoisomerase I and II (Constantinou et al., 1990), inhibit angiogenesis (Fotsis et al., 1995) and metastasis (Li et al., 1999), increase apoptosis (Shao et al., 1998a) (Sarkar & Li, 2003), increase the *in vitro* concentrations of transforming growth factor β (Kim et al., 1998) and increase protein and mRNA expression of the IGF-1 receptor and IRS-1 (Chen & Wong, 2004). Genistein can also regulate signalling pathways involved in cell proliferation (NFKB and AKT) (Davis et al., 1999; Li & Sarkar, 2002), and has

recently been shown to increase expression of all genes involved in cell cycle control apoptosis and DNA repair including BRAC1, p53, p21, Rad51, BARD, Bax GADD45 and RF1 (Caetano et al., 2006). Genistein can also regulate many genes for which roles are still being discovered (Barnes, 2004; Naciff et al., 2002). It is also possible that phytoestrogen effects are mediated via their action on growth factors and growth factor binding proteins (Adlercreutz, 2002a; Chen & Wong, 2004) with vegetarian diets rich in phytoestrogens associated with low plasma concentration of insulin, insulin Like Growth Factor-1 (IGF-1) and higher levels of IGF-1 binding proteins and SHBG (Giovannucci, 1999). Although most of these studies have been carried out with genistein, daidzein also exhibits anti-cancer properties (Scholar & Toews, 1994), although it has received less attention (Jing et al., 1993), and biochanin produced greater growth inhibition than genistein in MCF-7 breast cancer cells (Peterson et al., 1996) and has potent anti-mutagenic activity (Chae et al., 1991). The metabolite equol has also received greater attention recently with superior oestrogenic and antioxidant activity (Setchell et al., 2002).

In vitro studies suggest that physiological levels of oestrogen may also impact on the effect of isoflavones *in vivo* (Messina et al., 2006) and some findings suggest that the clinical relevance of these genomic and non-genomic effects found *in vitro* may not actually be significant (Santell et al., 2000). Thus caution is warranted in the interpretation of some of these studies because some of the effects may have been observed at higher concentrations than would be found *in vivo*.

1.2.2 Animal evidence

Approximately 70% of early studies in rat and mice models (Barnes, 1995) have found that feeds of soy protein or whole soy beans had protective effects, lowering the number of carcinogen-induced tumours when compared with standard diets. Isoflavone-depleted soy was found to be inactive in these models (Barnes, 1995). A recent review (Messina et al., 2006) also found that the majority of rodent studies showed inhibition of tumour development with soy or isoflavone supplementation prior to carcinogen- induced tumours (Magee & Rowland, 2004) (Constantinou et al., 2001; Yan et al., 2002) or implanted cancer cells (Gallo et al., 2006; Zhou et al., 2004). However there were exceptions to this (Allred et al., 2004a; Cohen et al., 2000; Day et al., 2001; Thomsen et al., 2005) and a review (Cooke, 2006) suggests that isoflavones have limited benefit once cancer is established in animal models.

Studies using ovariectomized mice injected with MCF-7 cells and implanted with estrogen pellets found that diets rich in genistein (Hsieh et al., 1998), isoflavone rich soy protein (Allred et al., 2001), or isoflavone extracts (Allred et al., 2004b) stimulated tumour growth compared with standard diets once implanted estrogen pellets were removed. Although this model is more inclined to show growth with weak estrogenic compounds, dietary genistein was still able to stimulate tumour growth in the presence of low levels of plasma estradiol mimicking conditions in postmenopausal women (Ju et al., 2006a).

The outcome of these studies are not only affected by the model used, but the type of soy product (Allred et al., 2004b; Constantinou et al., 2001) and specific isoflavone, with equol showing no stimulation of tumour growth when compared to daidzein (Ju et al., 2006b). Others have found that combinations of isoflavones with green/black tea (Zhou et al., 2004) or flaxseed (Thompson &

Ward, 2005) give even greater benefit than either alone. Furthermore, the timing of exposure to the carcinogen compared with ovariectomy and isoflavone intake can also affect outcomes, with genistein increasing tumours under certain conditions (Allred et al., 2004a) (Ueda et al., 2003).

Studies exposing pre-pubertal rats to genistein (Cotroneo et al., 2002; Lamartiniere et al., 2002a; Lamartiniere et al., 1995; Lamartiniere et al., 2002b) also reduced the number of tumours later induced by administering DMBA carcinogen to adults. It is thought that in this model genistein promotes cellular differentiation thereby causing breast epithelial cells to be less vulnerable to carcinogen initiation. Other studies have also shown that subcutaneous injection of genistein in newborn rats can protect against DMBA induced tumours while increasing tumour resistant biochemical markers (Cabanès et al., 2004; Hilakivi-Clarke et al., 1999; Lamartiniere et al., 1995).

Certainly there are limitations with rodent studies and the large quantities of equol produced by rat gut flora vs human should be considered when using rodent models given that equol's oestrogenic activities are comparable to genistein but with preferential binding to $Er\beta$ (Cooke, 2006; Gu et al., 2006; Setchell et al., 2002; Setchell et al., 2005).

1.2.3 Epidemiological evidence

1.2.3.1 Soy/Isoflavone intake

Although there is supportive evidence from *in vitro* and animal studies that isoflavone intake may influence the risk of breast cancer, this clearly requires confirmation in human studies. Epidemiologic studies indicate that in populations with a high isoflavone intake the risk of breast cancer is reduced (Adlercreutz, 1990; Ingram et al., 1997; Wu et al., 1998), and that within ethnic

groups whose diets are traditionally high in isoflavone intake, change in diet due to adoption of western lifestyle has led to an increase in breast cancer incidence (Buell, 1973; Lee et al., 1988).

Epidemiological studies of human dietary soy intake and breast cancer risk have recently been reviewed (Adlercreutz, 2002a; Adlercreutz, 2002b; Peeters et al., 2003) and include case control and prospective studies (**Table 1.2.3.1**).

To date, case control studies have assessed intake by dietary questionnaire (Hirose et al., 1995; Lee et al., 1991; Witte et al., 1997; Wu et al., 1996; Yuan et al., 1995) (Dai et al., 2001; Hirohata et al., 1985; Shu et al., 2001) and/or plasma/urine concentrations (Adlercreutz et al., 1986; Adlercreutz et al., 1982; Ingram et al., 1997; Pietinen et al., 2001). Those that have assessed intake through dietary questionnaires include the following:

Of studies conducted in China, (Yuan et al., 1995) found no association between intake of soy and risk of breast cancer in either pre or post menopausal women but did see a strong protective effect from fibre rich foods, while (Dai et al., 2001) did find a dose-response relation between soy food intake and breast cancer risk in women with high BMI ($p=0.02$). (Shu et al., 2001) found that soy intake during adolescence (age 13-15) was associated with lower risk of breast cancer in later life.

In Singapore, high intake of soy was associated with a reduced risk in pre-menopausal women but not in post-menopausal women (Lee et al., 1991) (Lee et al., 1992). In pre-menopausal women, high daily consumers (≥ 55.0 g soy product) showed a 60% (95% CI 0.2, 0.9) reduced risk compared with low daily consumers (<20.3 g). Similarly, in Japan, a high intake of soybean curd (tofu) was associated with a statistically non-significant reduced risk of breast cancer in pre-menopausal but not in post-menopausal women. However, in both

groups of women, there was no association between intake of miso soup, which was categorised in the food frequency questionnaire as daily or occasional-to-never, and breast cancer risk (Hirose et al., 1995). A recent smaller case control study in Japan, by the same author, found a statistically significant inverse association between tofu or isoflavone intake and risk of breast cancer in Japanese pre-menopausal women, but again this was not found with post-menopausal women (Hirose et al., 2005).

A study In the USA and Canada (Witte et al., 1997) found that tofu intake was associated with lower risk of pre-menopausal bilateral breast cancer. Among Asian Americans, intake of tofu was associated with a lower risk of breast cancer both overall and separately in pre- and post-menopausal women. Similar results were obtained after adjustment for relevant menstrual and reproductive factors and selected dietary factors (Wu et al., 1996). Asian American women in the highest intake group (>120 times/y) showed a 30% reduced risk (95% CI 0.4, 1.0) compared with women in the lowest intake group (<12 times/y).

Analysis by migrant status showed a significant protective effect in migrants but not in Western-born Asian Americans. Another case control study in a similar population of Asian Americans found that the risk of breast cancer was significantly inversely associated with soy intake during adolescence and adult life [OR=0.53, 95% CI=0.36-0.78], with women who reported soy intake at least once per week during adolescence showing a statistically significantly reduced risk of breast cancer (Wu et al., 2002). A recent case control study in Canada retrospectively assessed phytoestrogen intake during adolescence of breast cancer patients vs controls and also found that higher phytoestrogen intake (both isoflavones and lignans) during adolescence was associated with reduced risk of breast cancer, with a trend observed from the lowest to the highest

quartile [OR =0.91, 95% CI= 0.79-1.04, OR=0.71, 95% CI=0.62-0.82, p trend<0.001] (Thanos et al., 2006).

Recently a large case control study in Italy analysed intake of the six principal classes of flavonoids and found that only increased intake of flavones gave a significant reduced risk of breast cancer [OR=0.81, highest vs lowest quintiles, p-trend=0.02] (Bosetti et al., 2005). This appears to be in line with an earlier case control study done in Greece (Peterson et al., 2003).

Case Control studies that have assessed intake by plasma or urine concentrations include the following:

Two studies in Australia have looked at levels of isoflavones or their metabolites in biological fluids. (Ingram et al., 1997) carried out a case-control study of 144 breast cancer patients and an equal number of age-matched healthy controls in Australia. When examining isoflavone and metabolite excretion in quartiles, as equol excretion increased, the relative risk of breast cancer was significantly reduced. The risk for the highest quartile of excretion of equol, after adjustment for confounding variables, was one quarter that of the lowest quartile. Similar effects were seen with respect to the lignan, enterolactone. Daidzein showed a similar pattern, but the trend was not significant. Unfortunately, the laboratory assay for genistein was not reliable enough to report results. These results were surprising as the total phytoestrogen intake of the study subjects, as estimated from the urinary phytoestrogen excretion, was very low (Messina et al., 1997). A similar, preliminary study, also carried out in Australia but this time in post-menopausal women and controls, found that those with breast cancer had lower 24-h urinary daidzein excretion than controls (p=0.03) and a trend towards lower genistein (p=0.08) (Murkies et al., 2000). A study in China (Zheng et al., 1999) showed that breast cancer cases had substantially lower levels of

isoflavones in urine than age matched controls, while studies in Finland (Adlercreutz et al., 1986; Adlercreutz et al., 1982; Pietinen et al., 2001) have also found lower urinary and plasma enterolactone concentrations in women with breast cancer vs controls.

Two recent nested case control studies examining phytoestrogen concentrations from women participating in the large European Prospective Investigations into cancer (EPIC) have had conflicting results. In the UK population, where 333 women who developed breast cancer, higher equol and daidzein in serum and urine significantly increased breast cancer risk (Grace et al., 2004). In this population the majority of women were post-menopausal, although analysis was combined. Analysis of the Dutch cohort of 383 women who developed breast cancer found that high levels of all isoflavones were associated with lower breast cancer risk and this was significant for plasma levels of genistein. These results were the same in pre and post-menopausal women (Verheus et al., 2007).

A number of larger prospective studies (den Tonkelaar et al., 2001; Hirayama, 1992; Horn-Ross et al., 2002; Keinan-Boker et al., 2004; Key et al., 1999; Nomura et al., 1978; Yamamoto et al., 2003) have also been conducted using dietary assessments of intake as a marker for isoflavone status. Only one found significant results (Yamamoto et al., 2003), with consumption of miso soup and isoflavones (not soy foods) inversely associated with risk of breast cancer in Japanese women aged 40-59 years. Compared with those in the lowest quartile of isoflavone intake, the adjusted RRs for breast cancer for women in the second, third, and highest quartiles were 0.76 (95% CI = 0.47 to 1.2), 0.90 (95% CI = 0.56 to 1.5), and 0.46 (95% CI = 0.25 to 0.84), respectively (P for trend =

.043). This association was stronger in post-menopausal women (P for trend = .006) (Yamamoto et al., 2003).

Prospective studies that have examined plasma and urine concentrations of isoflavones include a study in pre-menopausal German women, which found a non-significant positive association between isoflavones and breast cancer risk (Piller et al., 2006) and a study in Dutch postmenopausal women which found a weak but non-significant negative association between genistein urine concentrations and breast cancer risk (den Tonkelaar et al., 2001).

More recently meta-analyses have been conducted to assess soy intake and breast cancer risk. (Trock et al., 2006) analysed 12 case control and 6 cohort studies taking original measures of soy (soyfood intake or urinary isoflavone excretion) and converting them to an estimate of grams of soy protein consumed daily. They found high soy intake was modestly associated with reduced breast cancer risk [OR =0.86, 95% CI =0.75-0.99] and this association was stronger in pre than post-menopausal women [OR=0.7, 95% CI= 0.58-0.85] vs [OR=0.77, 95% CI=0.6-0.98]. (Qin et al., 2006) looked at 14 case-control and 7 cohort studies, including studies published in Japanese and Chinese. They also found an inverse association with soy food/isoflavone intake and breast cancer [RR=0.75, 95% CI=0.59-0.95], with isoflavone intake resulting in a 20% decrease in risk [RR=0.81, 95% CI=0.67-0.99].

1.2.3.2 Hormone levels and breast cancer risk

Reviews of epidemiological and clinical data confirm that lifetime exposure to increasing concentrations of oestrogens and androgens significantly increases breast cancer risk (Clemons & Goss, 2001; Key, 1999; Peeters et al., 2003; Thomas et al., 1997). A review of data from prospective studies has found that levels of endogenous sex hormones are strongly associated with breast cancer

risk in post-menopausal women (Key et al., 2002), whilst a recent case control study nested within the European Prospective Investigation into Cancer and Nutrition cohort, found that elevated blood concentrations of androgens were associated with an increased risk of breast cancer from 370 pre-menopausal women who subsequently developed breast cancer vs 726 matched cancer free control subjects (Kaaks et al., 2005).

Breast cancer patients have been shown to have shorter menstrual cycle lengths than control subjects (Olsson et al., 1983), while Asian women have longer cycle lengths than those in Western populations (Treloar et al., 1967). Women with shorter cycles will have a greater lifetime exposure to oestrogen, and also spend more of their reproductive lifetime in the luteal phase, where the mitotic rate for breast tissue is almost fourfold greater than during the follicular phase. Isoflavone supplementation, by increasing the length of the follicular phase, may therefore reduce the risk of breast cancer. Thus soy isoflavones may reduce risk of breast cancer by changing levels of endogenous hormones and altering the menstrual cycle (Adlercreutz, 2002a) (Bingham et al., 1998).

1.2.4 Intervention studies examining effect of isoflavone supplementation on hormone levels

A number of studies have looked at the effect of isoflavone supplementation on hormonal endpoints, in both pre and post-menopausal women, with varying results, and these are summarised in Tables 1.2.4.1 and 1.2.4.2.

In the controlled environment of a metabolic unit, studies have shown that consumption of isoflavone-rich soy extends the length of the follicular phase and therefore overall menstrual cycle and decreases serum follicle-stimulating hormone and luteinizing hormone concentrations (Cassidy et al., 1995; Cassidy

et al., 1994). However, 36 oz of soy milk had no effect on LH or FSH levels in 10 pre-menopausal women with no change in mean menstrual cycle length (Lu et al., 2000a; Lu et al., 2001). Only two studies have shown a significant reduction in oestradiol concentrations (Lu et al., 1996a; Wu et al., 2000). All of the studies reviewed (Table 1.2.4.1) showed a reduction in total progesterone concentrations measured in the luteal phase but only (Lu et al., 1996a) found this to be a significant reduction. SHBG levels measured in these intervention studies also varied with only two studies showing increases, and they were in the follicular (Nagata et al., 1998) and luteal phase respectively (Wu et al., 2000). In a more recent randomised intervention study with 220 pre-menopausal women (Maskarinec et al., 2004), there appeared to be no significant difference in the oestrone, oestradiol, progesterone and SHBG levels of the intervention group consuming ~50mg isoflavones daily vs control group in samples taken in the luteal phase (day 19) after 3, 6, 12 and 24 months. The authors concluded that the effects of soy on the breast, if they exist, are mediated by mechanisms other than the lowering of circulating estrogen levels in pre-menopausal women.

In post-menopausal women Intervention studies have also produced mixed results (Table 1.2.4.2). In post-menopausal women most soy intervention studies reviewed have produced no difference in levels of oestradiol, LH, or FSH (Baird et al., 1995; Berrino et al., 2001; Brooks et al., 2004; Brzezinski, 1997; Duncan et al., 1999b; Foth & Nawroth, 2003; Murkies et al., 1995; Nicholls et al., 2002; Persky et al., 2002; Petrakis et al., 1996; Pino et al., 2000). In the large EPIC Oxford cross-sectional study (Verkasalo et al., 2001), there were no differences seen in oestradiol or SHBG levels of 456 postmenopausal women consuming daily soy milk. Another centre in the EPIC study (Norfolk)

found that urinary and serum levels of daidzein and genistein were negatively correlated with plasma oestradiol (Low et al., 2005) but that when women with certain genotypes were excluded these correlations were no longer significant. The author concluded that polymorphisms in genes encoding for enzymes involved in oestradiol metabolism (i.e. ESR1 PvuII CC genotype) may affect the concentration of plasma oestradiol, suggesting diet-gene interactions. Recent cross-sectional studies of post-menopausal women (Shoff et al., 1998), (Thomas et al., 1999), (Verkasalo et al., 2001), (Low et al., 2005) assessing phytoestrogen intakes through dietary questionnaires found correlations with SHBG and oestradiol levels were not significant.

Results from a number of human studies suggest that soy consumption may exert oestrogenic effects on breast tissue and highlight the fact that caution is necessary at this stage. In a 6-month study (Petrakis et al., 1996), breast fluid secretion in both pre- and post-menopausal women in response to soy protein isolate consumption (containing 37.4 mg genistein/day) increased 2-6 fold, as did the number of atypical cells in breast fluid. However, this was a pilot study, which did not include a control group.

In another study, this time in pre-menopausal women, (McMichael-Phillips et al., 1998), the rate of DNA synthesis by breast cells taken from biopsies of normal breast tissue from women with benign or malignant disease was enhanced by 14 days of soy feeding (60 g/day equivalent to 45 mg isoflavones) prior to surgery. (Hargreaves et al., 1999) also found nipple aspirate levels of apolipoprotein D were significantly lowered and pS2 levels raised in response to soy supplementation ($P < \text{or} = 0.002$), after two weeks of soy supplementation, which they suggest is indicative of an oestrogenic stimulus. However, in a more recent pilot study with 17 surgical patients, after two week isoflavone

supplementation, there appeared to be a statistically non-significant trend towards cancer growth inhibition in the isoflavone treatment group, as manifested by higher apoptosis/mitosis ratios in tumour biopsies compared with those from the control untreated group (Sartippour et al., 2004).

1.3 Insulin like Growth Factor -1 and binding proteins

Insulin like growth factor –1 (IGF-1) originally termed somatomedin (Daughaday et al., 1972) (Van Wyk et al., 1980) is structurally homologous to the hormone insulin and, like insulin, is involved in stimulating the anabolic processes needed for growth during puberty and adult life (Juul, 2003; Ranke et al., 1999). Much of the IGF-1 found in plasma is hepatically produced upon the stimulation of growth hormone released by the pituitary. Like the growth factor itself, the receptors for insulin and IGF-1 are structurally very similar, creating the potential for cross affinity and binding. Indeed, many tissue types, including those of the ovaries and breast, express receptors for both growth hormones. Target tissues also have the capacity to produce IGF-1 locally when needed (D'Ercole et al., 1980) as it is involved in the paracrine control of the tissue's cellular proliferation. This includes normal breast tissue (Paik, 1992), where it is required for mammary development (Ruan & Kleinberg, 1999). Expression of IGF-1 in recent studies assessed by RT-PCR, show that levels tend to be higher in non-neoplastic breast tissue than in adjacent cancerous tissue (Voskuil et al., 2004) (Chong et al., 2006b) (Chong et al., 2006a) providing evidence for a paracrine relationship between IGF-1 producing stromal fibroblasts and neighbouring epithelial cells (Ellis et al., 1994). Concentrations of IGF-1 in plasma are at least a hundred fold greater than that of insulin. However, IGF-1 is bound to a family of six binding proteins (IGFBP-1

to 6) (Baxter et al., 1998), produced by many tissues and found in plasma.

Again growth hormone is responsible for the regulation of these binding proteins and is regulated by the nutritional status of the individual. Binding proteins are also produced locally within the target tissues

More than ninety percent of IGF-1 is bound to IGFBP-3 and an associated acid-labile subunit, forming a high molecular weight complex, which is not available to the tissues (Juul, 2003). Clinical studies suggest that individuals with IGFBP-3 levels in the upper range of normal may have decreased risk of breast cancer while *in vitro* and animal studies show that IGFBP-3 can also regulate cell growth and apoptosis through IGF independent mechanisms (Ali et al., 2003). IGFBP-1, on the other hand, binds only a small fraction of circulating IGF-1, with hepatic production regulated through the portal supply of insulin, with insulin inhibiting its production in the liver (Brismar et al., 1994). Thus, regulation of this binding protein is sensitive to food intake and has been shown to correlate inversely with the secretion of insulin (Kaaks & Lukanova, 2001; Lee et al., 1997b).

1.3.1 IGF-1 and breast cancer risk

Next to oestrogen, Insulin-like growth factor-1 is one of the most potent mitogen for breast cancer cell lines *in vitro* (Karey & Sirbasku, 1988). It can rescue cells from cytotoxic chemotherapy through induction of proliferation and inhibition of apoptosis, stimulating both MAPKinase and PI3Kinase pathways (Gooch et al., 1999; Laban et al., 2003).

In Vivo, antibody blockade of the IGF-1 Receptor inhibits growth of human breast cancer cells in mice (Arteaga et al., 1989) while liver specific gene deletion knockout of IGF-1 gene results in mice with reduced circulating IGF-1

levels and reduced rates of cancer growth and metastasis (Yakar et al., 2005). Lower circulating levels of IGF-1 also delay the onset of induced mammary tumours *in vivo* (Wu et al., 2003).

In humans, significantly elevated concentrations of IGF-1 have been found in breast cancer patients when compared with healthy controls (Bruning et al., 1995; Peyrat et al., 1993). Healthy subjects with high IGF-1 concentrations, when controlled for binding proteins, have been shown to have an elevated risk of breast cancer (RR 7.28, highest vs. lowest quintile (Hankinson et al., 1998). High concentrations of IGF-1 in serum have now been associated with an increased risk of breast cancer, in case control studies (Bohlke et al., 1998; Bruning et al., 1995; Del Giudice et al., 1998; Johansson et al., 2004; Li et al., 2001; Peyrat et al., 1993; Yu et al., 2002) and more recently in prospective studies (Allen et al., 2005; Hankinson et al., 1998; Krajcik et al., 2002; Muti et al., 2002; Rinaldi et al., 2005a; Rinaldi et al., 2006; Schernhammer et al., 2005; Toniolo et al., 2000). It is therefore now accepted that elevated levels of IGF-1 are an independent risk factor for pre-menopausal breast cancer, and this has been confirmed by meta-analysis (Renehan et al., 2004) while a review of recent cohort studies show that associations with IGFBP-3 are less clear (Renehan et al., 2006).

The first study to show a difference in IGF-1 between breast cancer cases and controls compared 44 primary breast cancer patients with 92 controls all over the age of 35. This showed significantly higher levels in the patients, with median concentration of IGF-1 of 152 ng/ml as apposed to 115ng/ml (Peyrat et al., 1993). In a larger study done by (Bruning et al., 1995), 150 Stage I- II breast cancer patients between the ages of 38 to 75 had serum levels analysed for IGF-1 and BP3 just prior to or one month after surgery. These were compared

with 441 women from a population base control group. Pre-menopausal cases showed increased IGF-1, decreased IGFBP-3 and increased IGF-1/BP-3 ratios. The relative risk was 7.34 for the highest compared with the lowest quintile of IGF-1/BP-3.

A more recent study examined levels in 94 patients with pre-menopausal ductal carcinoma-in-situ vs 76 controls and found that women in the two highest tertiles of IGF-1 and the lowest tertile of IGFBP-3 were at 3.7-fold higher risk than women in the lowest tertile of IGF-1 and highest two tertiles of IGFBP-3 (Bohlke et al., 1998).

A prospective study conducted with data obtained in the Nurses Health Study cohort, analysed plasma concentrations of IGF-1 and IGFBP-3 from samples collected in 1989-1990 in women who went on to develop breast cancer after this date up to 1994. They compared these 397 women with 620 aged matched controls. Among pre-menopausal women younger than 50 years, the mean IGF-1 concentration was significantly higher in the cases than in the controls ($p=0.009$) with a relative risk of 4.58 (Hankinson et al., 1998).

Most reviews agree that this relationship is more prominent in pre-menopausal women (Fletcher et al., 2005; Renehan et al., 2004; Schernhammer et al., 2005; Shi et al., 2004; Sugumar et al., 2004) but not all studies have found this (Kaaks et al., 2002; Rinaldi et al., 2006; Rinaldi et al., 2005b), with a recent prospective analysis of the Nurses Health Study II cohort study finding no associations in pre or post-menopausal women (Schernhammer et al., 2006).

The reasons for this discrepancy may be related to the age of participants at recruitment (Rollison et al., 2006) or genetic differences in subpopulations (Al-Zahrani et al., 2006). Polymorphisms of IGF-1 and BP genes have been shown to change risk of breast cancer although authors conclude they do not play a

major role in altering breast cancer risk in Caucasian populations (Canzian et al., 2005) (Canzian et al., 2006).

However the effect of differences in time between blood collection and tumour diagnosis cannot be ruled out given the possibility of tumour influence on IGF levels (Schernhammer et al., 2006). A recent updated meta-analysis of all these studies concludes that there is a modest association between IGF-1 and breast cancer risk in pre-menopausal and young women but no discernible associations between IGF-1, IGFBP-3 and post-menopausal breast cancer risk (Renehan et al., 2006).

1.3.2 IGF-1 and oestradiol

Oestradiol and IGF-1 appear to work together in the proliferation of breast cancer cells with increasing support for cross talking between the ER and IGF-1R signalling pathway (Kato et al., 2000; Laban et al., 2003). Oestradiol has been shown to stimulate expression of IGF-1, the IGF-1R, IRS-1 and down-regulate expression of IGF binding proteins (Huynh et al., 1996b; Lee et al., 1999; Ruan et al., 1995; Stewart et al., 1990), while genistein has been shown to enhance the IGF-1 signalling pathway in proliferation of MCF-7 cells at low concentrations (Chen & Wong, 2004).

1.3.3 IGF-1 and Tamoxifen

The cytotoxic effects of tamoxifen are also partially mediated through IGF-1R signalling (Guvakova & Surmacz, 1997). Tamoxifen has been shown to lower circulating IGF-1 levels by approximately 30% in breast cancer patients (Campbell et al., 2001; Colletti et al., 1989; Helle et al., 1996; Lonning et al.,

1992; Pollak et al., 1990) and healthy women (Bonanni et al., 2001; Decensi et al., 1998; Decensi et al., 2003).

The mechanism by which tamoxifen alters IGF status remains to be fully elucidated. However, tamoxifen can alter IGF-1 levels by reducing the production of GH from the pituitary (Tannenbaum et al., 1992) thereby lowering the amount of IGF-1 released into circulation (Bonanni et al., 2001; Pollak et al., 1990) and in *in vivo* models, directly inhibits the expression of the IGF-I gene in common target organs for breast cancer metastasis (Huynh et al., 1993). It also has direct action as an antiestrogen on breast cancer cells and appears to reduce the number of IGF-1 receptors (Huynh et al., 1996a) and up-regulates expression of binding proteins released by the cells themselves (Huynh et al., 1996b; Winston et al., 1994).

1.3.4 IGF-1 and diet

Although numerous epidemiological and *in vitro* studies positively associate IGF-1 and its binding proteins with increased cancer risk (Renehan et al., 2004) the major determinants of inter-individual variability in circulating IGF-1 levels are not yet fully understood. Studies show that genetic (Fletcher et al., 2005) and nutritional factors, such as energy and protein restriction, play an important role in the regulation of IGF-1 (Thissen et al., 1994).

Reports examining the IGF and diet relationship have mostly been observational and cross sectional human studies are summarised in table 1.3.4.1. These studies have reported associations between IGF-1, BP-3 and BP-1 levels and a range of dietary variables including calcium/dairy foods, fibre, vegetables, fat, carbohydrates and micronutrients, and are discussed in further detail in chapter 5.

1.3.5 IGF-1 and isoflavone/soy intake

A few of the more recent epidemiological studies have also examined the association between IGF-1 and intake of soy/phytoestrogens (Nagata et al., 2003; Probst-Hensch et al., 2003; Vrieling et al., 2004). No correlations were seen between soy product intake and IGF-1 or BP-3 levels in 261 Japanese pre-menopausal women (Nagata et al., 2003) or 386 pre and post-menopausal Dutch women (Vrieling et al., 2004) assessed using food frequency questionnaires. However there was a positive association found between soy intake and IGF-1 in 321 Chinese men studied (Probst-Hensch et al., 2003) and a positive association with plant lignans and BP-1 concentrations in 162 postmenopausal Dutch women (Vrieling et al., 2004).

Prior to reports on the cross-sectional associations between isoflavone intake and IGF, two intervention studies were conducted between 1997-9, the results of which are presented in this thesis. These have now been published (Woodside et al., 2006) (Campbell et al., 2004). Since conducting these isoflavone supplementation studies, other intervention studies have also published their findings on the effects of soy or isoflavone supplementation on IGF-1 and binding proteins (these are summarised in Table 1.3.5.1). Of the studies published to date, only our study (Campbell et al., 2004) and to our knowledge one subsequent study published very recently (Vrieling et al., 2007) have looked at the effects of isoflavone supplementation alone. Further discussion of these intervention studies can be found in chapter 5.

1.4 Conclusion

In vitro and animal evidence suggests that dietary isoflavones may be protective against the development of breast cancer. Epidemiologic evidence remains

largely inconclusive, and while case-control studies have produced some convincing evidence that increased dietary isoflavone intake correlates with reduction in breast cancer risk, early intervention studies have produced conflicting results. In light of the emerging role for IGF-1 and its binding proteins in breast cancer aetiology, this thesis aims to examine the effects of Isoflavones on IGFs, to determine whether, like other antiestrogens, this may be a possible mechanism of action, which could further explain the possible protective effects of these compounds in the diet.

We propose to do this by first examining the effects of the SERM tamoxifen on levels of IGF-1 and its binding proteins in breast cancer patients, then by looking at the short-term effects (single dose or one week supplementation) of a phytoestrogen-rich bar and then a placebo-controlled isoflavone supplementation study over one month on IGF-1 and binding proteins, in healthy pre and post-menopausal women. Finally we examine the effects of isoflavones on DNA synthesis in breast cancer cell lines *in vitro* to compare their actions with tamoxifen in the presence or absence of IGF-1 and oestradiol.

Table 1.2.3.1. Studies on the association between phyto-oestrogens in plasma and urine or phyto-oestrogen-rich diet and breast cancer risk.

Table 1.2.4.1: Interventions with soy foods/supplements in premenopausal women-effects on hormones

Table adapted from (Maskarinec et al., 2004)

Table 1.2.4.2: Interventions with soy foods/supplements in postmenopausal women- effects on hormones.

Author	Isoflavone dose; type of food	Length (mo)	Type	n	E1 (%)	E2 (%)	SHBG (%)	LH (%)	FSH (%)
Baird et al. 1995	165mg; soy foods	1	RI RC	66 25	ND	ND	-6.8 -5.7	-2.5 -2.0	-4.4 -2.8
Murkies et al. 1995	45g soy or wheat flour	3	RI (soy) RC (wheat)	23 24	ND	ND	ND	ND	-1.6 -9.8**
Petrakis et al. 1996	75 mg; soy protein isolate	6	CO	10	ND	-0.94	+44.4	ND	ND
Brzezinski et al. 1997	~375 mg; tofu, soy drink, miso and flax seed	3	RI RC	78 36	ND	-22.2 -13.6	+27.6* -7.5	ND	ND
Duncan et al. 1999	10, 65, 129 mg; soy protein isolate	3	CO (10vs129)	18	-5.6	-11.9	+5.6*	+5.5	+3.04
Knight et al. 1999	40 or 160 mg	3	RI (40) (160)	37	ND	ND	-11.7 -16.3	ND	-6.3 -16
Pino et al. 2000	69mg; soy milk	10wks	Interven	20	-8.59	+0.17	+15.7*	ND	-10.2
Berrino et al. 2001	~45 mg; fibre and phtoestrogen rich diet	5	RI RC	49 50	ND	-18.0 -5.5	+25.2* +3.6	ND	ND
Persky et al. 2002	56, 90 mg; isolated soy protein	6	RI RI RC	24 23 25	ND	+40 +10.2 -18.6	-12.8 -5.2 -17.2	ND	-7.3 -0.68 -1.4
Foth et al. 2003	20mg; soy protein	6	Interven	16	ND	-15.2	ND	+3.6	+2.2
Brooks et al. 2004	41.9mg; soy flour in muffin	4	RI RC	16 15	+40 +5.25	+89.6 -24.7	ND	ND	ND
Nicholls et al. 2002	60mg; textured soy protein	14 days	Interven	7	ND	+25.6	ND	-6.9	-3.3
Arjamandi et al. 2005	60mg; 25g protein from soy products or control protein with no soy or isoflavones.	12	RI ^b RC ^b	35 27	+14.3 -2.8	+39.4 -11.1	-14.5* -12.4	ND	-3.6 -11.2

Table 1.3.4.1: Epidemiological studies- Dietary correlations with IGF-1, BP3 and BP1 concentrations

Author	n	Subjects	Results			
			IGF-1	IGFBP-3	IGF-BP-1	other
Devine et al 1998	119	Postmenopausal women	Zinc +ve Protein +ve	Riboflavin +ve	ND	Baseline-prior to controlling for age and weight, IGF-1 correlated with Calcium, Cholesterol, Energy, Iron, potassium, magnesium, sodium, niacin, phosphorous, protein, riboflavin, thiamine and zinc.
Kaklamani et al. 1999	115	73%female and 27% male; age 30-84	Red meat +ve Fat +ve Oils +ve Energy from lipids +ve Energy from carbo +ve	Saturated fat -ve	ND	
Ma et al. 2001	193 vs 318	Men, Cases with colorectal cancer vs controls; age 40-84 (physicians health study)	Dairy food +ve	Dairy food +ve	ND	
Mucci et al 2001	112	Men, control group for liver cancer study. Greek population; all ages.	Alcohol -ve Cooked tomatoes -ve	Not shown	ND	IGF1/BP3 ratio: Cooked tomatoes -ve
Allen et al 2002	292	Women, age 20-70, Meat eaters vs vegetarians vs vegans	Animal protein +ve Animal plus soy protein +ve Nonsoya plant protein -ve NSP -ve	None	Nonsoya plant protein +ve SFA -ve NSP +ve Zinc +ve	IGF1 13% lower in vegan women than Meat eaters of vegetarians (p=0.006). BP1 20-40% higher in vegans than ME and Vegans(p=0.005). Significant increase in IGF-1 in vegan women who drink $\geq \frac{3}{4}$ pint of soya milk daily.
Holmes et al 2002	1037	Women, age 43-69, 526 pre & 456 postmenopausal	B-cryptoxantine +ve Animal protein +ve	Alcohol +ve Sat fat intake -ve Lycopene +ve Vitamin C +ve	ND	Cereal, pasta, milk and total dairy +ve associated with IGF-1

			Vitamin D +ve Vitamin A +ve Calcium +ve Zinc +ve Energy +ve (with BMI <25)	Energy +ve		
Baibas et al 2003	620	225 men & 395 women, age 35-65, Greek EPIC study	Ethanol -ve Saturated lipids +ve	Not shown	ND	
DeLellis et al 2004	1000	Multiethnic; age 45-75; 50% male/female	Men: fat from meat -ve Low fat milk -ve	Men: total fat -ve Saturated fat -ve Fat from meat - ve Low fat milk -ve Women: alcohol +ve	ND	
Giovanucci et al 2003	753	Men; age 40-75; health professionals follow up study	Animal & vegetable protein +ve (milk, fish & poultry) Potassium +ve Zinc +ve Magnesium +ve Calcium +ve Phosphorous +ve Energy intake +ve(men with BMI <25)	Not shown	ND	IGF1/BP3 ratio: Vitamin E Phosphorous had the strongest association with IGF-1 probably because a good marker of both meat and dairy intake.
Gunnell et al 2003	344	Men; age 50-70	PUFA +ve Calcium +ve Dairy products +ve	PUFA +ve	ND	IGF1/BP3 ratio: inversely related to vegetable intake and tomato containing products. Carbohydrate had a +ve trend with IGF-1 levels but not significant.

			Milk +ve			
Heald et al 2003	257	Multiethnic; age 25-78; ~50% men/women	Fat intake +ve Sat fat +ve Protein +ve Carbohydrate +ve Starch +ve Energy +ve	ND	Total fat -ve PUFA -ve Total protein -ve	
Nagata et al 2003*	261	Premenopausal women; age 20-54; Japanese	Vitamin D +ve	Total fat -ve	ND	No correlations seen with soy product intake and IGF-1 or BP-3 levels
Probst-Hensh et al 2003*	338	321 men and 326 postmenopausal women; age >50; Singapore chinese	Soy +ve (men) Calcium +ve Cholesterol +ve LDL +ve Triacylglycerol +ve Total Chol/HDL +ve	Sat Fat -ve Omega 3 PUFA +ve Fibre +ve Cholesterol +ve LDL +ve Triacylglycerol +ve Total Chol/HDL +ve	ND	Inverse trend with Alcohol intake and IGF-1 levels
Vrieling et al 2004*	286	224 pre & 162 postmenopausal women; age 49-69; EPIC study in Netherlands	Alcohol -ve (premen)	Alcohol +ve (premen)	Postmen: alcohol -ve Plant lignans +ve Cereals -ve	No correlations seen with soy intake/phytoestrogens and IGF-1 or BP-3 levels
Larsson et al 2005	226	Men; age 42-76; Sweden	Protein +ve Zinc +ve Red meat +ve Potassium +ve	ND	ND	Fish & seafood +ve trend with IGF-1
Maskarinec et al 2005	258	Premenopausal women; age 34-46; multiethnic	Dietary fibre +ve Vitamin A +ve	Zinc +ve	ND	IGF1/BP3 ratio: Iron +ve Calcium +ve Saturated fat +ve trend. No ethnic differences.

Norat et al 2006	2109	Women; age 35-70; EPIC study in Europe	Protein +ve Calcium +ve Magnesium +ve Phosphorus +ve Potassium +ve Vit B6 +ve Vit B12 +ve Beta- carotene-ve	Calcium +ve Phosphorus +ve	ND	Adjustment of IGF-1 for BP3; PUFA –ve Zinc not analysed as could not standardise estimates across countries. Food correlations with IGF-1 were Milk +ve, Cheese +ve and Vegetables -ve
---------------------	------	---	--	-------------------------------	----	---

Table 1.3.5.1: Interventions with soy foods or supplements – effects on insulin like growth factors and binding proteins

Author	Isoflavone dose; type of food	Length (mo)	Type	n	Results %change			
					IGF-1	IGFBP-3	IGF- BP-1	other
Maskarinec et al 2005	50mg;soy foods	24	RI RC	97 ^a 99 ^a	+0.48 +0.26	-1.47 -0.33	ND	Iso excretion +ve assoc with IGF-1 (p=0.04)
Wangen et al. 2000	10, 65,129mg soy protein isolate	3	CO	14 ^a 17 ^b	+10.8(PO)* 65 -6.54 (129)*	+5(EF)65 -4.5(129)	ND ND	
Adams et al. 2003	3 vs 83mg; soy protein	12	RI (3) RC (83)	58 ^{bc} 65 ^{bc}	+8.1* +7.7*	0 +1.0	ND ND	
Khalil et al. 2002	88mg; soy protein vs milk protein control	3	RI ^c RC ^c	24 22	Inc* Inc (figures not given)	ND	ND	N.B Only Soy protein supp significantly increased IGF-1 from baseline (<0.0001)
Kaaks et al. 2003	~45 mg; fibre and phytoestrogen rich diet	5	RI ^b RC ^b	50 49	-5.9 -4.4	-4.5 -2.2	+12.2* -6.4	
Campbell et al. 2004	86mg; isoflavone supplement (no protein) & placebo	1	CO	16 ^a 7 ^b	+1.5 (s Vs P) +2.77	+2.5 -1.1	+13.2 +12.3	
Woodside & Campbell et al. 2006	80mg; soy and linseed bar (18g protein)	1 week	interven	10 ^{ab}	+71*	18.3*	-40.7	
Arjamandi et al. 2003	88mg; soy protein vs milk protein control	3	RI ^b RC ^b	20 22	+68.6** +36.4**	ND	ND	Increases in IGF-1 greater for women on HRT.
Arjmandi et al. 2005	60mg; 25g protein from soy products or control protein with no soy or isoflavones.	12	RI ^b RC ^b	35 27	+26.3** +12.8**	+1.85 +5.2**	ND	Both significantly increased IGF-1 but IGFBP-3 only sig increased on soy.
Spentoz et al. 2003	114mg; soy protein supp on low fat diet	2	Interven ^c	15	+22	-1	ND	Increase in IGF-1 of 9% after low fat diet then increase by 22 % from baseline on

								soy protein p=0.02
Gann et al 2005	88mg; soy protein powder with or without Isoflavones (subjects on both LFHF and UD)	3	RI ^a RC ^a	153	2.9 1.2	-1.2 -3.4*	7.5 6.3	Small but significant decrease in BP3 of 2.3% when both groups analysed together.
Vrieling et al 2007	84mg; isoflavone supplement (no protein) & placebo	2	CO	37 ^c	-2.5	+0.43	-6.4	Change in IGF-1 and serum Equol were negatively correlated post supplement.

a =premenopausal

b=postmenopausal

c=men

CHAPTER TWO – METHODS

2.1 Collection and preparation of biological samples

Urine samples (24 h) were stabilised with 1-2 g/l ascorbic acid and sodium azide (0.1 g/L) and stored at -80°C until analysis. Spot urine samples were not stabilised but immediately frozen after collection. Blood samples either kept at 4°C for a maximum of 30 min (plasma) or were allowed to clot for 30 minutes in the dark (serum) and then centrifuged at 3000 rpm for 10 min. Serum or plasma was aliquoted and stored at -80°C until analysis. Lymphocytes were collected into a Vacutainer Cell Preparation Tube (Becton Dickinson, New Jersey), extracted and washed according to manufacturer's instructions and stored at -80°C.

2.2 Endpoint measurements

2.2.1 Diet

Food frequency questionnaires (from EPIC) and food diaries were analysed using DietPlan 5 for Windows 98 using standardised composition (McCance et al., 1978) and food portions established by MAAF (Crawley et al., 1988) to assess background diet through out the studies.

2.2.2 Phytoestrogens

Urine samples were analysed for the isoflavonoids genistein and daidzein, the isoflavan equol and the lignans enterolactone and enterodiol levels, using gas chromatography-mass spectrometry following enzyme hydrolysis, extraction into ether and silylation according to Morton et al. (Morton et al., 1994).

2.2.3 Ascorbic acid

Ascorbic acid concentrations were determined as described by Vuillemier & Keck (1989) (Vuillemier & Keck, 1989). The procedure is based on the enzymatic oxidation of ascorbic acid and subsequent quinoxaline formation to generate a fluorescent derivative measured on the Cobas Fara centrifugal analyser. The whole blood sample, stabilised with dithiothreitol (Fisher Scientific), was centrifuged and separated upon arrival at the laboratory. The plasma was precipitated by adding 100 μ l of plasma to 900 μ l of 5% metaphosphoric acid (Fisher Scientific) in a conical microcentrifuge tube and vortex mixed. Once precipitated the sample was stable for 1 year at -70°C . Our samples were assayed within 6 months of collection. Working standards (0, 1, 5, 10, 15, 20 $\mu\text{mol/l}$) were prepared for each run from a freshly made stock standard of 20 $\mu\text{mol/l}$ ascorbic acid in 5% metaphosphoric acid. Approximately 200 μ l of standards and samples were placed in Cobas cups, and oxidising reagent (9.9 ml, 2 mol/l acetate buffer and stock ascorbate oxidase) and coupling reagent (20 mg of 1,2-phenylenediamine dissolved in 20 ml deionised water) added to appropriate reagent wells. The assay was then started. The intra-assay CV was calculated for several different serum samples. At a mean concentration of 4.2 $\mu\text{mol/l}$, the inter-assay CV was 6.80% ($n=9$), at 50.6 $\mu\text{mol/l}$ the inter-assay CV was 0.72%, while at 151.0 $\mu\text{mol/l}$ the inter-assay CV was 1.50%. Antioxidant malondialdehyde (MDA) levels in serum were analysed using HPLC method (Young & Trimble, 1991).

2.2.4 Retinol, α -tocopherol and carotenoids

Concentrations of serum α -tocopherol and β -carotene were measured by HPLC according to the method of Catignani *et al.* (1983)(Catignani & Bieri, 1983) and Thurnham *et al.* (1988) (Thurnham *et al.*, 1988). They were monitored by ultraviolet detection following extraction into heptane. The assay also measures serum concentrations of retinol, lycopene and α -carotene. Serum or standard (200 μ l) was pipetted into 4 ml glass tubes. Internal standard, a mixture of retinol acetate and α -tocopherol acetate in ethanol (400 μ l - Labscan) was added to each tube and the ethanol precipitated the protein from the serum. Heptane (Analar or 99% - 700 μ l (Labscan)) containing 0.5 g/l butylated hydroxytoluene (BHT) was added to the tubes which were then vortexed vigorously for 1 minute. The tubes were then centrifuged for 10 minutes at 1200 g and 500 μ l of the resulting top heptane layer transferred to a set of identically labelled glass tubes. The complete contents (1300 μ l) of the standard tube contents were dried. These tubes were evaporated to dryness in a rotary evaporator under vacuum for 1- 2 hours. The samples were either assayed immediately or stoppered and stored dry in a -70°C freezer. For HPLC analysis, the samples were reconstituted in ethanol (100 μ l), vortexes and assayed using UV detection. A 45% methanol:45% acetonitrile:10% dichloromethane (200 ml, not filtered, degassed for exactly 5 minutes using Romil HPLC grade solvents, (Analab, Dromore, Co. Down) mobile phase was used with a flow rate of 1.8 ml/min. The column used was Hypersil ODS 5 μ m (Alltech), with changing wavelengths of 325 nm, 292 nm and 450 nm to pick up the retinol, tocopherol and carotenoid peaks respectively. A Thermo Separation Products Constametric 4100 pump was coupled with a Spectra-Physics AS1000 autosampler and a Spectra-Physics Analytical UV2000 ultraviolet

detector. The system was controlled using a Spectra-System SN4000 interface and operated using PC1000 software from Thermo Separation Products. A quality control serum sample was assayed at the start of every batch of samples. The inter-assay CV for α -tocopherol was 5.4% (n=7) and 10.0% (n=7) for β -carotene. The CVs for retinol, lycopene and α -carotene were similar to those for β -carotene. The addition of internal standard allowed the calculation of an intra-assay CV of 4.9% (n=29).

2.2.5 Serum Lipids and lipid peroxidation

Serum lipids were assessed, again at the start and end of each supplementation phase, using standard spectrophotometric assays.

Serum total cholesterol, was estimated using an enzymatic CHOD-PAP kit, while serum triglycerides were measured using the Peridochrom GPO-PAP kit (both Boehringer Mannheim). Precipitation for HDL-cholesterol estimation employed phosphotungstic Mg^{2+} reagents (Lopes-Virella *et al.*, 1977). All cholesterol assays were carried out on the Cobas Fara auto-analyser. As well as employing internal quality control procedures, total cholesterol and HDL-cholesterol were subject to external quality control through the World Health Organisation reference laboratory in Prague. LDL-cholesterol was estimated using the Friedewald formula:

$$LDL\ cholesterol\ (mmol/l) = total\ cholesterol - HDL\ cholesterol - 0.45 \times triglyceride$$

Levels of aqueous phase lipid hydroperoxides were assessed in plasma by the FOX 1 (ferrous oxidation in xylene orange) assay according to Wolff (Wolff,

1994). Susceptibility of LDL to oxidation was assessed according to McDowell et al. (McDowell et al., 1995).

2.2.6 COMET Assay

DNA damage was analysed by the COMET assay in both lymphocytes according to Spanswick et al. (Spanswick et al., 1999).

2.2.7 Creatinine

Creatinine concentrations from 24 hour urine collections were measured using a Sigma colorimetric assay.

2.2.8 Hormones

17 β -oestradiol, Progesterone, LH, FSH and SHBG were measured in serum using routine IRMA/RIA methods as follows:

17 β -oestradiol was measured directly using the DiaSorin Clinical Assays™ Oestradiol-2 Radioimmunoassay. The standards, controls and samples were incubated with oestradiol tracer and antiserum raised in rabbits for 2 hours at 37°C in a water bath. After a second incubation for 15 min with a precipitating reagent, containing antibody to rabbit IgG the samples were centrifuged for 15 min at 2000G to separate free tracer from antibody-bound tracer. The tubes were then decanted and the supernatant discarded. The pellet was counted in a gamma counter. A standard curve was then prepared from the standards ranging from 10 to 2000pg/ml. The unknown values were obtained from the standard curve by interpolation.

Progesterone was measured using the DiaSorin Clinical Assays™

GammaCoat™ Progesterone-¹²⁵I RIA Kit for the quantitative determination of

progesterone levels in serum. This was a competitive binding radioimmunoassay whereby the amount of [125 I]-labeled progesterone bound to the antibody on the plastic coated tube was inversely proportional to the concentration of progesterone present in the serum. The samples, standards and controls were added to the Anti-Progesterone pre-coated tubes in duplicate. Progesterone tracer was then added and the tubes were incubated at 37°C for 60 minutes. The tubes were then decanted and counted in a gamma counter. A standard curve was then prepared from the standards ranging from 0.3 to 60ng/ml. The unknown values were obtained from the standard curve by interpolation.

Serum Luteinizing Hormone (LH) and Follicle-stimulating Hormone (FSH) were measured using the NETRIA Immunoradiometric assay (IRMA). The samples, standards and controls were added to the tubes along with the solid phase (sheep) Anti-LH or FSH, the [125 I]-iodine labelled Anti-LH or FSH (monoclonal) and the assay buffer. These were vortexed and placed on a rotary mixer overnight. The next morning, 2ml of wash buffer were added to the tubes and after centrifugation at 1000G for 5 minutes, the supernatant was decanted. This washing step was repeated before the tubes were decanted and counted in a gamma counter. A standard curve was then prepared for LH and FSH from the standards ranging from 1-260 U/L or 0.5-200 U/L respectively. The unknown values were obtained from the standard curve by interpolation.

Sex Hormone Binding Globulin was measured in serum using the [125 I] Immunoradiometric Assay kit from Orion Diagnostica. This procedure was based on the principles of a non-competitive "liquid-phase" immunoradiometric assay. Samples, standards and controls were diluted in assay buffer and added in duplicate to the test tubes. Equal volumes of SHBG Antiserum (Rabbit) and

[¹²⁵I] anti-SHBG (Mouse monoclonal) were added to all tubes. Tubes were vortexed and incubated at room temperature for 60 minutes. The separation reagent, containing goat anti-rabbit IgG covalently bound to solid particles, was then added and tubes were vortexed and incubated for a further 15 minutes at room temperature. Two ml of 0.9% saline was added and the tubes centrifuged at 2000G for 15 minutes. The tubes were decanted and counted in a gamma counter. A standard curve was then prepared from the standards ranging from 6.25, 200 nmol/L. The unknown values were obtained from the standard curve by interpolation.

2.2.9 IGF-1, BP-1 and BP-3

Levels of IGF-1, IGF-BP1, IGF-BP3 were measured in serum using ELISA methods (Diagnostic Systems Laboratories). The DSL-10-5600 IGF-1 ELISA kit was an amplified "one step" sandwich-type immunoassay. This assay included an extraction step in which IGF-1 was separated from its binding protein in serum by acidification followed by ethanol precipitation of the IGFBP fraction. The standards, controls and extracted samples were then incubated with anti IGF-1 antibody labelled with the enzyme horseradish peroxidase (HRP) in microtitration wells coated with another anti-IGF-1 antibody. After incubation and washing, the wells were incubated with the substrate tetramethylbenzidine (TMB). An acidic stopping solution was then added and the degree of enzymatic turnover of the substrate was determined by dual wavelength absorbance measurement at 450nm with background wavelength correction set at 620nm. The absorbance measured was directly proportional to the concentration of IGF-1 present. The IGF-1 standards were used to plot a standard curve of absorbance versus IGF-1 concentration and the IGF-1 concentrations were

then calculated using the line of best fit plotted using the MINITAB statistical package. Controls were assayed at the beginning and end of the plate while samples were assayed in duplicate.

IGFBP-3 and total IGFBP-1 were measured using the DSL-10-6600/7800 ELISA kits. In these "two-step" immunoassays, standards, controls and samples were incubated in microtitration wells which had been coated with an anti-IGFBP-3/1 antibody. After incubation and washing, the wells were treated with another anti-IGF-3/1 antibody labelled with HRP. After a second incubation and washing step, the wells were incubated with TMB. After adding the acidic stopping solution, the enzymatic turnover was measured at 450 and 620nm as above. The absorbance measured was directly proportional to the concentration of BP present. The standards were used to plot a standard curve of absorbance versus BP concentration and the BP concentrations were then calculated using the line of best fit plotted using the MINITAB statistical package.

2.3 Development of an Enzyme Linked Immunosorbant Assay (ELISA) for the quantitative measurement of IGF-BP3 in Human Serum.

Work carried out under supervision of Dr Brian Rafferty at the National Institute for Biological Standards and Control, Blanche Lane, South Mimms, Potters Bar, Hertfordshire EN6 3QG, UK.

2.3.1 Introduction

Prior to commencing the intervention studies described in this thesis, an attempt was made to develop an in-house assay for IGFBP-3. IGFBP-3 is the major binding protein for IGF-1 in human serum, and the best indicator of clinical conditions involving Growth Hormone deficiencies (Kim et al., 2001) and is also potentially useful as an anti cancer molecule/biomarker (Ali et al., 2003).

ELISAs are the preferred method for measuring Insulin Like Growth Factors and their binding proteins (Rutanen & Pekonen, 1991). At the time of this research, ELISA kits were only commercially available from Diagnostics Systems Laboratories (Diamandi et al., 2000). These were expensive, therefore the development of an in-house ELISA would be have been useful. This work would also allow a better theoretical understanding of ELISAs and could hopefully be used as an alternative to commercial kits in future.

2.3.2 Method

An antibody to IGF-BP3 had previously been raised in White Long Eared Rabbits in the Animal House at NIBSC. IGF-BP3 was human recombinant DNA

(non glycosylated) and was from the standardised stock made at NIBSC. Both the Antibody and the Binding protein had been previously assessed using Radio-iodination and Fast Protein Liquid Chromotography (FPLC).

The assay development involved;

- Coating the 96 well microtitre plate with antibody over night (optimal concentration and volume to be determined).
- Coating the Plate with samples and standards (optimal concentration and volume to be determined).
- Addition of Biotinylated Antibody (optimal concentration and volume to be determined).
- Addition of Avidin Horse Radish Peroxidase and Substrate Reagent for colour development (optimal concentration and volume to be determined).
- Incubation (optimal number of steps time and temperature to be determined between each stage).

Assays which attempted to optimise the assay as detailed above are described below in results.

2.3.3 Results


Assay 1:

Aim: To determine the concentration of the coating antibody needed to give adequate binding of the substrate and to establish the dilution of biotinylated antibody needed to give an adequate signal.

Method:

COATING ANTIBODY (FPLC-RBP3/3)	200µl @ 0.5µg/ml 200µl @ 1.0 µg/ml
SAMPLES/ STANDARD NG IGFBP-3 (93/560) (3.5µg/ml) SERUM POOL DSL STD MAELI'S SERUM	100µl @ 100ng/ml □ 1.5625ng/ml (2 fold dilns)
BIOTINYLATED ANTIBODY (FPLC-RBP3#5)	100µl @ 1/1000 diln 100µL @ 1/2000 diln
INCUBATION STEPS	1 step, 3hr

Results:



Findings: Absorbance readings using 1.0g/ml coating antibody and 1/1000 diln of biotinylated antibody are consistently greater as would be expected.


Assay 2 & 3:

Aim: To determine the effect of halving the volumes used in Assay one.

Method:

COATING ANTIBODY (FPLC-RBP3/3)	100 μ L @ 0.5 μ G/ML 100 μ L @ 1.0 μ G/ML
SAMPLES/ STANDARD NG IGFBP-3 (93/560) (3.5 μ g/ml) SERUM POOL DSL STD MAELI'S SERUM	50 μ L @ 100ng/ml □ 1.5625 ng/ml (2 fold dilns)
BIOTINYLATED ANTIBODY (FPLC-RBP3#5)	50 μ L @ 1/1000 diln 50 μ L @ 1/2000 diln
INCUBATION STEPS	1 step, 3hrs

Results:



Findings: Halving the volumes used gives an adequate signal if maximal parameters are used. (See assay one) The control (NSB) values were consequently lower in Assay 2.

Assay 4 & 5


Aim: To assess the effects of altering incubation times from 3 hours to one.

Method

<u>COATING ANTIBODY (FPLC-RBP3/3)</u>	100µL @ 1.0 µG/ML
<u>SAMPLES/ STANDARD</u> NG IGFBP-3 (93/560) (3.5µg/ml) SERUM POOL DSL STD MAELI'S SERUM	50µl @ 20ng/ml □ 0.3125ng/ml (2 fold dilns) 50µl @ 40ng/ml □ 0.6125ng/ml (2 fold dilns)
<u>BIOTINYLATED ANTIBODY (FPLC-RBP3#5)</u>	50µl @ 1/1000 diln 50µl @ 1/2000 diln
<u>INCUBATION STEPS</u>	Assay 4: 1 step, 1 hr Assay 5: 1 step, 3 hr

Results:

Assay 4+5



Findings: The 3 hour incubation gives an appropriately larger signal while the 1 hour incubation lends itself to a loading effect due to differing incubation with the coating antibody.

Assay 6


Aim: To assess the DSL standards and control solutions VS the serum pool and NG IGFBP-3 over one hour incubation.

Method:

COATING ANTIBODY (FPLC-RBP3/3)	100µL @ 1.0µG/ML
SAMPLES/ STANDARD	
NG IGFBP-3 (93/560) (3.5µg/ml)	50µl @ 20ng/ml □ 0.3125ng/ml (2 fold dilns)
SERUM POOL	
DSL STD	50µl @ 40ng/ml □ 0.625ng/ml (2 fold dilns)
MAELI'S SERUM	50µl @ 0.7, 4, 17, 45, 120ng/ml & 4±1.2, 17±5

	50µl @ 1/50 & 1/100 diln
<u>BIOTINYLATED ANTIBODY (FPLC-RBP3#5)</u>	50µl @ 1/1000 diln
<u>INCUBATION STEPS</u>	1 step, 1hr

Results:



Findings: This assay showed quite clearly that the serum responds differently to the NG BP-3 while DSL standards are derived from human serum.

Assay 7


Aim : To assess the effects of a 3 hour incubation on an assay similar to assay 6 and to determine whether time delay in loading a plate will make a significant impact on absorbance readings.

Method:

<u>COATING ANTIBODY (FPLC-RBP3/3)</u>	100µL @ 1.0µG/ML
<u>SAMPLES/ STANDARD</u>	
NG IGFBP-3 (93/560) (3.5µg/ml) beginning and end of plate ~ 15 min	50µl @ 20ng/ml □ 0.3125ng/ml (2 fold dilns)

SERUM POOL DSL STD MAELI'S SERUM	50µl @ 40ng/ml □ 0.625ng/ml (2 fold dilns) 50µl @ 0.7, 4, 17, 45, 120ng/ml & 4±1.2, 17±5 50µl @ 1/50 & 1/100 diln
BIOTINYLATED ANTIBODY (FPLC-RBP3#5)	50µl @ 1/1000 diln
INCUBATION STEPS	1 step, 3 hr

Results:



Findings: Three hours gives a better signal but a plate loading effect is still visible.


Assay 8

Aim: To extend the sensitivity of the assay while using the serum pool as the standard.

Method:

<u>COATING ANTIBODY (FPLC-RBP3/3)</u>	100µL @ 1.0 µG/ML
<u>SAMPLES/ STANDARD</u>	
NG IGFBP-3 (93/560) (3.5µg/ml)	
SERUM POOL	50µl @ 40µg/ml(1/100) □ 1/25600
DSL STD	0.7, 4, 17, 45, 120ng/ml % 4±1.2, 17
MAELI'S SERUM	±5 (x 4)
<u>BIOTINYLATED ANTIBODY (FPLC-RBP3#5)</u>	100µl @ 1/1000 diln
<u>INCUBATION STEPS</u>	1 step, 3 hr

Results:



Findings: This assay appeared to have an effect across the two halves of the plate, however these results are not.

Assay 9


Aim: To assess the effects of spiking the serum pool with known standards to establish recovery values.

Method:

COATING ANTIBODY (FPLC-RBP3/3)	100 μ L @ 1.0 μ G/ML
SAMPLES/ STANDARD NG IGFBP-3 (93/560) (3.5 μ g/ml) SERUM POOL DSL STD MAELI'S SERUM	50 μ l @ 20ng/ml \rightarrow 0.312ng/ml (2 fold dilns) 50 μ l @ 40ng/ml \rightarrow 0.3125ng/ml (2 fold dilns) Serum + DSL 50 μ l @ 56ng/ml \rightarrow 3.5ng/ml (2 fold) Serum + WDB 50 μ l @ 32ng/ml \rightarrow 2.0ng/ml(2 fold)
BIOTINYLATED ANTIBODY (FPLC-RBP3#5)	50 μ l @ 1/1000 diln
INCUBATION STEPS	1 step, 3 hrs

Results:

Assay 9



Findings: The spiked dilution's appear to give readings as would be expected.

These need to be extended.


Assay 10

Aim: To assess the sensitivity of the assay by extending the lower end of the standard serum curve. To determine the IGFBP-3 levels in an 'unknown' serum.

Method:

<u>COATING ANTIBODY (FPLC-RBP3/3)</u>	100µL @ 1.0µG/ML
<u>SAMPLES/ STANDARD</u> NG IGFBP-3 (93/560) (3.5µg/ml) SERUM POOL DSL STD MAELI'S SERUM	50µl @ 10ng/ml → 0.156ng/ml (2 fold dilns) 50µl @ 10ng/ml → 0.078ng/ml (2 fold dilns) 50µl @ 1/500, 1/2500 diln 50µl @ 1/100, 1/50 diln
<u>BIOTINYLATED ANTIBODY (FPLC-RBP3#5)</u>	50µl @ 1/1000 diln
<u>INCUBATION STEPS</u>	1 step, 3 hrs

Results:



Findings: Providing enough dilutions are done for the unknown serum the standard curve appears to give consistent results. Further work needs to be done to further test this.

Assay 11 & 12


Aim: To determine whether there is a difference in signal when the biotinylated antibody is loaded on to the plate before the serum in order to diminish any plate effect obtained through difference in binding times. (see assay 7)

Method:

COATING ANTIBODY (FPLC- RBP3/3)	100µL @ 1.0µG/ML
SAMPLES/ STANDARD	
NG IGFBP-3 (93/560) (3.5µg/ml)	
SERUM POOL	50µl @ 10ng/ml → 0.15625ng/ml (2 fold dilns)(x2 on Assay 11 + 12)
DSL STD	
MAELI'S SERUM	

<u>BIOTINYLATED ANTIBODY (FPLC-RBP3#5)</u>	Assay 11: 50µl @ 1/1000 before serum Assay 12: 50µl @ 1/1000 after serum
<u>INCUBATION STEPS</u>	1 step, 3hrs

Results:



Findings: The results would suggest that although there is no great difference between the curves, the control (NSB) values for assay 12 are significantly lower.

Assay 13


Aim: To assess the affects of a two step incubation on the overall sensitivity and loading effects of the assay.

Methods:

<u>COATING ANTIBODY (FPLC-RBP3/3)</u>	100µL @ 1.0µG/ML
--	------------------

<u>SAMPLES/ STANDARD</u> NG IGFBP-3 (93/560) (3.5µg/ml) SERUM POOL DSL STD MAELI'S SERUM	100µl @ 40ng/ml → 0.3125ng/ml (2 fold dilns) (x4)
<u>BIOTINYLATED ANTIBODY (FPLC-RBP3#5)</u>	100µl @ 1/1000 diln
<u>INCUBATION STEPS</u>	2 step: 2 hrs then 1 hr

Results:



Findings: Although this assay appears to have remedied the plateau effect, later assays would suggest that the concentration of BP-3 may have started at 20ng/ml rather than 40ng/ml.


Assay 14

Aim: To extend the sensitivity of the two step assay and to assess the signal from the non glycosylated and other serum material.

Method:

COATING ANTIBODY (FPLC-RBP3/3)	100 μ L @ 1.0 μ G/ML
SAMPLES/ STANDARD NG IGFBP-3 (93/560) (3.5 μ g/ml) SERUM POOL DSL STD MAELI'S SERUM	100 μ l @ 100ng/ml \rightarrow 0.195ng/ml (2 fold dilns) 100 μ l @ 80ng/ml \rightarrow 0.156ng/ml (2 fold dilns) 100 μ l @ 1/100, 1/500, 1/2500 dilns 100 μ l @ 0.7, 4, 17, 45, 120ng/ml 100 μ l @ 1/20, 1/100, 1/500 dilns
BIOTINYLATED ANTIBODY (FPLC-RBP3#5)	100 μ l @ 1/4000 diln
INCUBATION STEPS	2 step: 2hrs then 1hr

Results:



Findings: The sensitivity of this assay appears to be repeatable down to a concentration of 0.156ng/ml and is highly reproducible.


Assay 15

Aim: To determine how different concentrations of biotinylated antibody will affect the two step assay sensitivity.

Method:

<u>COATING ANTIBODY (FPLC-RBP3/3)</u>	100 μ L @ 1.0NG/ML
<u>SAMPLES/ STANDARD</u> NG IGFBP-3 (93/560) (3.5 μ g/ml) SERUM POOL DSL STD MAELI'S SERUM	100 μ l @ 100ng/ml \rightarrow 0.195ng/ml (2 fold dilns)
<u>BIOTINYLATED ANTIBODY (FPLC-RBP3#5)</u>	100 μ l @ 1/500, 1/1000, 1/2000 dilns
<u>INCUBATION STEPS</u>	2 step: 2hrs then 1 hr

Results:



Findings: The absorbance readings obtained from using the 1/1000 diln of biotinylated antibody give an adequate response.


Assay 16

Aim: To assess the cross reactivity of the antibody with IGF-I and IGF-II.

Method:

<u>COATING ANTIBODY (FPLC-RBP3/3)</u>	100 μ L @ 1.0 μ G/ML
<u>SAMPLES/ STANDARD</u> NG IGFBP-3 (93/560) (3.5 μ g/ml) SERUM POOL IGF-I (87/518) (3.1 μ g/ml) IGF-II (96/538) (5 μ g/ml) DSL STD MAELI'S SERUM	100 μ l @ 40ng/ml \rightarrow 0.0488ng/ml (2 fold dilns) 100 μ l @ 1.0 μ g/ml \rightarrow 0.0156ng/ml (2 fold dilns) 100 μ l @ 1.0 μ g/ml \rightarrow 0.0156ng/ml (2 fold dilns)
<u>BIOTINYLATED ANTIBODY (FPLC-RBP3#5)</u>	100 μ l @ 1/1000 diln
<u>INCUBATION STEPS</u>	2 step: 2hrs then 1hr

Results:



Findings: IGF-I and II did not give a significant signal at any concentration and do not therefore appear to be binding to the antibody.

Assay 17


Aim: To assess possible competitive binding of IGF-I and II with BP-3 by spiking the serum with known amounts of the growth factors.

Method:

<u>COATING ANTIBODY (FPLC-RBP3/3)</u>	100 μ L @ 1.0 μ G/ML
<u>SAMPLES/ STANDARD</u> NG IGFBP-3 (93/560) (3.5 μ g/ml) SERUM POOL SERUM WITH WDB SERUM WITH IGF-I SERUM WITH IGF-II	 100 μ l @ 80ng/ml \rightarrow 1.25ng/ml (2 fold dilns) 100 μ L @ 20ng/ml \rightarrow 0.625ng/ml (2 fold dilns) 100 μ l @ 40ng/ml \rightarrow 0.625ng/ml (2 fold dilns) 100 μ l @ 40ng/ml \rightarrow 0.625 ng/ml (2 fold dilns)
<u>BIOTINYLATED ANTIBODY (FPLC-RBP3#5)</u>	100 μ l @ 1/1000 diln
<u>INCUBATION STEPS</u>	2 step: 2hrs then 1hr

Results:

Assay 17



Findings: IGF-1 and II appear to be competing with the antibody in binding to BP-3 however there does not appear to be a significant effect on absorbance levels and binding appears to be constant.

2.3.4 Discussion

Optimal conditions were found to be with the two-step incubation, whereby the samples were allowed to bind to the walls of the plate for 2 hours before the Biotinylated Antibody was added, this appeared to remedy the hook affect seen with a one step incubation of 3 hours. The standard curve established was found to be highly reproducible. However as serum must be screened before use at NIBSC facility we were unable to assess the assay on more than one individual serum sample.

In order to validate this assay for future use of the clinical assessment of patients, many serum samples would need to be tested and the assay would need to be compared with commercial ELISAs that use human serum for production of standard curves. Although initial results were encouraging this ELISA was left at the experimental stage as there would need to be considerable testing before it could be published as a standard ELISA for IGFBP3 and used for the analysis of samples in subsequent studies.

It was not therefore used for the analysis of IGFBP-3 in this thesis, and represents early stage assay development.

2.4 Cell Culture

MCF-7 and MDA MB -231 cells (kindly donated by Ludwig cancer research group) were maintained in Phenol Red IMEM (GIBCO) supplemented with 5% FCS (GIBCO), 1% L-Glutamine (sigma) and 1% Fungizone (GIBCO) in 5% CO₂ at 37°C. Conditions involved growing cells in phenol red free IMEM supplemented with 5%FCS for 72 hours prior to plating. Cells were then seeded in 96 well plates (NUNC) at 1×10^4 cells per well in 5% DCC-FCS (Dextran charcoal coated foetal calf serum) phenol red free IMEM and incubated for 24 hours. Medium was changed to serum free IMEM for another 24 hours before the compounds were added. Stock solutions of tamoxifen, Genistein (4,5,7-trihydroxyisoflavone), Daidzein (4,7-dihydroxyisoflavone), Equol and 17 β -estradiol (Sigma) in ethanol, were used to produce the required concentrations of compounds in culture medium. Control conditions also contained the appropriate amount of ethanol (0.1%).

2.5 BrdU assay

The principle of the assay was to determine cell proliferation by measuring 5-bromo-2-deoxyuridine (BrdU) incorporation during DNA synthesis in proliferating cells. This assay indicates growth rate as opposed to cumulative cell growth. Conventional methods use thymidine, however this technique is based on the incorporation of the pyrimidine analogue 5-bromo-2'-deoxyuridine (BrdU). The BrdU assay was performed using a colorimetric ELISA kit (Biotrak version 2 from Amersham Pharmacia Biotech).

The assay involved culturing cells in the presence of the respective test substances in a 96-well microtitre plate at 37°C for 1 to 5 days. BrdU was added

to the cells and reincubated (usually 2-24h). During this labelling period, the pyrimidine analogue BrdU was incorporated in place of thymidine into the DNA of proliferating cells. After removing the culture medium, the cells were fixed and the DNA denatured by addition of fixative (the denaturation of the DNA was necessary to improve the accessibility of the incorporated BrdU for detection by the antibody). The peroxidase-labelled anti-BrdU bound to the BrdU incorporated in newly synthesised, cellular DNA. The immune complexes were detected by the subsequent substrate reaction, and the resultant colour read at 450nm in a microtitre plate spectrophotometer. The absorbance values correlated directly to the amount of DNA and thereby to the number of proliferating cells in culture (Amersham-Pharmacia-Biotech, 1998).

2.5.1 BrdU incorporation

After 24 hours of treatment with test compounds, 5-bromo-2-deoxyuridine (BrdU) was added to 200µl of medium to give a final concentration of 10µM per well. Incorporation was allowed to proceed for a further 24 hours at 37°C before the medium was collected and cells were fixed with an ethanol-containing fixative. Cells were blocked then incubated for 30 min at room temperature on a plate shaker with a peroxidase conjugated mouse monoclonal antibody (1:100). After washing, peroxidase substrate (tetramethylbenzidine) was added and the reaction stopped after 10 min with 1 M H₂SO₄. Absorbance was read within 5 min at 450nm using a 96-well plate reader. Results were from five replicate wells with experiments performed two to three times.

2.5.2 Optimal conditions

To develop conditions whereby assay blank and non specific binding control absorbance readings were low and the difference between controls and stimulated cells were optimal, the following parameters were manipulated:

1. Different cell types were used to establish which growth patterns were more suited to the assay. These included T47D and ZR75 breast cancer cell lines which were both unsuitable due to slow and for ZR75 non-monolayered growth which made denaturing and fixing cells more difficult.
2. Seeding density was altered between 1×10^3 cells/well and 5×10^5 cells/well with 1×10^4 cells/well giving optimal results.
3. Percentage of DCC-FCS used for seeding the cells was varied between 1%, 2.5% and 5% with optimal results seen with 5% DCC-FCS.
4. Length of time cells were starved in 0% DCC-FCS was varied between 0, 24 and 48 hours with 24 hours giving optimal results.
5. Percentage of DCC-FCS used after starvation was varied between 0%, 1%, 2.5% and 5% with compounds added to 0% starvation medium giving optimal results.
6. Test compounds were added for 3, 6 24 and 48 hours before BrdU was incorporated, with 24 hour incubations giving optimal results.
7. BrdU incubation was varied to 3, 6 and 24 hours with optimal results seen after 24 hours.
8. Cells were fixed for 30 and 60 minutes with 30 minutes giving optimal results.
9. Incubation with the blocking buffer was varied between 30 and 60 minutes with 30 minutes giving optimal results.

10. Incubation time with the Peroxidase labelled antibody was varied between 30 and 90 minutes with 30 minutes giving optimal results.
11. Incubation with the substrate TMB was varied between 5 and 30 minutes with optimal results seen after 10 minutes.

CHAPTER THREE – THE EFFECT OF TAMOXIFEN ON IGF STATUS IN BREAST CANCER PATIENTS – A CASE CONTROL STUDY

3.1 Introduction

Breast cancer is the most prevalent cancer in women in Western populations.

The risk of breast cancer increases greatly with age, and other established hormone related risk factors include early menarche, late age at menopause, and delayed age at first pregnancy (Bernstein & Ross, 1993). Family history is also important, although established genetic factors only account for about 5% of breast cancers in Western populations (McPherson et al., 2000).

Insulin-like growth factor-1 is a potent mitogen for breast cancer cell lines (Karey & Sirbasku, 1988). Increased concentrations of IGF-1 have been found in patients with breast cancer when compared with healthy controls (Bruning et al., 1995). In addition, healthy subjects with high IGF-1 concentrations, when controlled for binding proteins, have been shown to have an increased risk of breast cancer (RR 7.28, highest vs. lowest quintile) (Hankinson et al., 1998), and this has been confirmed by meta-analysis (Renehan et al., 2004). It is therefore now largely accepted that high circulating levels of IGF-1 and possibly BP-3 are a risk factor for pre-menopausal breast cancer (Del Giudice et al., 1998; Johansson et al., 2004; Krajcik et al., 2002; Li et al., 2001; Renehan et al., 2006; Renehan et al., 2004; Shi et al., 2004; Toniolo et al., 2000; Yu et al., 2002).

Tamoxifen, an anti-oestrogen in use for over 20 years or more, is the most commonly used agent in the treatment of hormone responsive breast cancer

(Katzenellenbogen et al., 1997). Clinical trials have confirmed the benefit of tamoxifen in preventing breast cancer recurrence and improving disease-free survival (1988; Baum, 1998). Tamoxifen has been tested as a preventative agent in healthy women and appears to significantly reduce both invasive and non-invasive breast cancer in those at increased risk of the disease (Fisher et al., 1998; Powles et al., 1998).

Apart from its main action via oestrogen receptors, tamoxifen has been found to possess numerous other plausible mechanisms in controlling tumour growth, such as binding to protein kinase C (O'Brian et al., 1985), inhibiting ornithine decarboxylase activity (Thomas et al., 1989) and inhibiting angiogenesis (Gagliardi & Collins, 1993). Tamoxifen can also act as an antioxidant (Wei et al., 1998), and can suppress the production of TGF α and stimulate the production of TGF β (Carmichael et al., 2000; Winston et al., 1994).

In addition, tamoxifen initially seems to reduce IGF-1 concentrations in breast cancer patients (Pollak et al., 1990) and increase concentrations of its binding proteins, but studies with longitudinal sampling and long-term follow-up have yet to be carried out.

The aim of the study was to compare concentrations of IGF-1 and its binding proteins in patients with breast cancer (n=14) with those found in age and sex matched control subjects (n=23). A further aim was to assess the longitudinal effect of tamoxifen on IGF status in patients with breast cancer.

Study published in (Campbell et al., 2001).

3.2 Materials and methods

Non-fasting serum samples were collected before surgery from patients with stage I-III invasive breast cancer, both pre-menopausal and post-menopausal. Serum samples were collected, again non-fasting, after <9, <18 and <27 months of tamoxifen treatment at routine surgical outpatients appointments. These 14 patients were part of a larger prospective study looking at the role of diet and lifestyle in survival after breast cancer. They formed a subset of women where blood samples had been collected prior to surgery, before tamoxifen treatment had begun. The baseline concentrations were compared with age- and sex-matched control subjects (n=23) taken from the general population. In the patients with breast cancer, no patient showed evidence of metastatic disease during the treatment period. None of the patients suffered from any other endocrine or malignant disorder during the treatment period. All patients had been prescribed 20mg of tamoxifen daily and gave written informed consent to participate. The study was approved by the Ethical Committee of the Royal Free and University College London Medical School.

3.2.1 Laboratory methods

Serum was stored at -80°C until analysis. Concentrations of IGF-1, IGF binding protein (IGFBP-1), and IGFBP-3 were measured in serum by means of enzyme linked immuno-sorbent assay kits (ELISA) kits purchased from Diagnostics Systems Laboratories (London, UK).

See Chapter two for further detail of Laboratory Methods.

3.2.2 Statistical methods

Owing to the small numbers of subjects involved, non-parametric tests were used throughout. The IGF status of patients with breast cancer compared with healthy controls was analysed using a Mann Whitney-U test. Analysis of the change in IGF status at particular time points compared with baseline values was carried out using Wilcoxon signed rank tests. Test results were considered to be significant when $p < 0.05$.

3.3 Results

The mean (SE) patient age was 51.0 (2.7) years (range, 38-70). Five of the 14 patients were over 50 y.

Concentrations of IGF-1, IGFBP-1 and IGFBP-3 at baseline did not differ between cases and controls (**Table 3.3.1**). Patient concentrations at baseline and during tamoxifen treatment can be seen in (**Table 3.3.2**).

The concentrations of the three variables measured were within the expected range reported in the literature. IGF-BP1 was increased significantly after 18 and 27 months of tamoxifen treatment (**fig 3.3.1**). IGFBP-3 was significantly increased from baseline at nine, 18 and 27 months (**fig 3.3.2**). IGF-1 was not decreased significantly (**fig 3.3.3**), but the IGF-1 to IGFBP-3 ratio, a commonly used measure of bioavailable IGF-1 (Hankinson et al., 1998), was significantly decreased both at nine and 18 months (**fig 3.3.4**). The IGF-1 to IGFBP-3 ratio was not significantly different from baseline at 27 months.

Table 3.3.1: Baseline concentrations of insulin-like growth factor 1 (IGF-1), IGF binding protein 1 (IGFBP-1), and IGFBP-3 in patients and controls

	Patients (n=14)	Controls (n=23)	P value
IGF-1 (ng/ml)	166.6(42.0)	194.2 (63.4)	NS
IGFBP-1 (ng/ml)	13.9 (2.9)	17.8 (3.9)	NS
IGFBP-3(ng/ml)	3119 (507)	3211 (598)	NS

NS, not significant. Mann Whitney-U test

Table 3.3.2: Patient concentrations of insulin-like growth factor 1 (IGF-1), IGF binding protein 1 (IGFBP-1), and IGFBP-3 at baseline and during tamoxifen treatment

	Pre-Tamoxifen baseline (n=14)	<9 months Tamoxifen (n=11)	<18 months Tamoxifen (n=12)	<27 months Tamoxifen (n=5)
IGF-1 (ng/ml)	166.6 (42)	144.3 (35.8)	149.4 (44.7)	181 (26.9)
IGFBP-1 (ng/ml)	21.6 (16.6)	38.9 (23.4)	52.0 (41.8)*	40.7 (24.9)*
IGFBP-3 (ng/ml)	3119 (507)	3673 (476)**	3445 (634)*	3409 (501)*

* P<0.05, **P<0.01 using Wilcoxon signed rank test compared with pre-tamoxifen baseline concentrations




Figure 3.3.1: Effect of tamoxifen supplementation on insulin-like growth factor binding protein 1 (IGFBP-1).

Values are median (black line) with interquartile ranges (boxed area) and minimum-maximum values (bars). * $p < 0.05$ for differences from baseline;

Wilcoxon signed rank test.




Figure 3.3.2. Effect of tamoxifen supplementation on insulin-like growth factor binding protein 3 (IGFBP-3).

Values are median (black line) with interquartile ranges (boxed area) and minimum-maximum values (bars). * $p < 0.05$ for differences from baseline; Wilcoxon signed rank test.




Figure 3.3.3: Effect of tamoxifen supplementation on insulin-like growth factor 1 (IGF-1).

Values are median (black line) with interquartile ranges (boxed area) and minimum-maximum values (bars). * $p < 0.05$ for differences from baseline; Wilcoxon signed rank test.




Figure 3.3.4: Effect of tamoxifen supplementation on insulin-like growth factor 1 (IGF-1) to IGF binding protein 3 (IGFBP-3) ratio.

Values are median (black line) with interquartile ranges (boxed area) and minimum-maximum values (bars). * $p < 0.05$ for differences from baseline; Wilcoxon signed rank test.

3.4 Discussion

We found that tamoxifen has the ability to lower bioavailable IGF-1 (calculated as the IGF-1 to IGFBP-3 ratio) for at least 18 months. Although concentrations of IGF-1 were not reduced significantly, concentrations of its major binding protein IGFBP-3 increased significantly, thereby reducing the amount of IGF-1 available. However, this ratio of IGF-1 to IGFBP-3 was not significantly reduced from baseline at 27 months, so that the effect of longer treatment remains to be elucidated. Tamoxifen also significantly increased concentrations of IGFBP-1 from baseline after 18 months of treatment. This increase has also been seen in other studies (Helle et al., 1996; Lahti et al., 1994).

Our finding of a lack of effect on IGF-1 concentrations is surprising because other studies have shown a reduction in IGF-1 by tamoxifen (Helle et al., 1996; Pollak et al., 1990). This might be the result of the small numbers of patients in our study group or population selection. However, another study (Lahti et al., 1994) also failed to show an effect on IGF-1 at a median follow up of 29 months. These investigators had seen a significant decrease in IGF-1 values after six months of tamoxifen treatment and their data point to a limited effect after long term treatment. Our data also indicate an initial (although non-significant) reduction in IGF-1, which is lost as follow up time increases. This indicates a potentially important effect of duration of treatment on outcome and highlights the need for further longitudinal studies with strictly timed follow up periods.

We also found no differences between cases and controls in terms of IGF-1, IGFBP-1, and IGFBP-3 concentrations. Again, this is surprising because earlier studies have revealed differences in similar groups. A lack of such a finding may again simply reflect a lack of numbers. We were unable to divide our

subjects into pre-menopausal and post-menopausal groups, again owing to the small sample size, and this may have been reflected in our results because IGF-1 values have been shown to be a stronger predictor of breast cancer risk in pre-menopausal than in postmenopausal women (Hankinson et al., 1998).

Our study was not placebo-controlled, and therefore subjects were not randomised to a treatment regimen; a placebo-controlled study would have distinct design advantages over the one described here. However, placebo-controlled trials of tamoxifen would now be considered ethically unsound (Fisher et al., 2000). An earlier placebo controlled, long term study showed a significant reduction in IGF-1 after a mean follow up of 27 months (minimum follow up three months) but longitudinal samples were not taken (Pollak et al., 1990). The samples used in our study were non-fasting and this may be important because IGF values may fluctuate according to nutrient intake (Clemmons & Underwood, 1991).

Changes in serum concentrations of IGF-1 have been shown to correlate positively with serum oestradiol concentrations in menstruating women (Massa et al., 1993). This agrees with findings that IGF-1 can vary over the menstrual cycle, with significantly lower values being found during the menstrual phase than the follicular and luteal phases (Helle et al., 1998). Although other studies found that IGF-1 concentrations were dependent on the menstrual cycle phase they considered this to be a fairly minor variation (Juul et al., 1997). Our study did not take into account the effect of the menstrual cycle and it may be important to control for this in future work.

The mechanism by which tamoxifen alters IGF status has not been fully elucidated. However, tamoxifen is thought to alter IGF-1 values by reducing the production of growth hormone from the pituitary, thereby lowering the amount of

IGF-1 that is produced by the liver and released into the circulation. Tamoxifen also has direct action as an antioestrogen on breast cancer cells and appears to alter the amount of IGF-1 and binding proteins released by the cells themselves (Pollak et al., 1990; Winston et al., 1994).

Since this work was published, there have been significant developments in the field, both in the use of SERMS and other treatments for breast cancer, and in how these treatments affect IGF concentrations. These will now be summarised. Tamoxifen has been shown to reduce breast cancer incidence by 30-40% in high risk subjects, in large phase III Trials (Cuzick et al., 2002; Fisher et al., 1998; Gasco et al., 2005; Veronesi et al., 2003). However, with the increased risk of endometrial cancer and venous thrombotic events attributed to tamoxifen other compounds including lower dose tamoxifen, other SERMS, aromatase inhibitors, and retinoids are now being assessed. Some of these trials have included analysis of IGF-1 and its binding proteins as cancer risk biomarkers.

Low dose tamoxifen studies have shown that 10mg/day gives comparable reductions in IGF-1 levels to standard 20mg/day doses (Bonanni et al., 2001; Decensi et al., 1998; Decensi et al., 1999; Decensi et al., 2003) with possible dose-response relationships seen in IGF-1/BP-3 ratios (Bonanni et al., 2001). A recent study by (Guerrieri-Gonzaga et al., 2006) has shown a 13% reduction in IGF-1 levels after 40 months of 5mg/day tamoxifen in 235 pre-menopausal women at risk of breast cancer, while the use of low dose tamoxifen for combination hormone replacement therapy users has shown a significant reduction in IGF-1 and increase in BP-3 for those who received 5mg/day tamoxifen over 12 months (Decensi et al., 2007).

Other SERMS including Raloxifene have been trialed in large studies and show 76% reduced risk of breast cancer with no increase risk in endometrial cancer (Cauley et al., 2001; Martino et al., 2004) with the National Surgical Adjuvant Breast and Bowel Project Study of tamoxifen and Raloxifene trial (STAR) showing that Raloxifene is as effective as tamoxifen in reducing the risk of invasive breast cancer and has a lower risk of thromboembolic events and cataracts but a nonstatistically significant higher risk of non-invasive breast cancer (Vogel et al., 2006). Raloxifene has also been shown to reduce IGF-1 and increase BP-3 when compared with placebo after 3 months (Andersson et al., 2002; Eng-Wong et al., 2003) and 2 years in post-menopausal type II diabetes patients (Duschek et al., 2004).

Recent trials have also shown improved efficacy and decreased side effects of aromatase inhibitors such as anastrozole and letrozole compared to tamoxifen (Baum et al., 2002; Ellis et al., 2001; Howell et al., 2005; Mouridsen et al., 2001; Nabholz et al., 2000). However, the effects of Vorozole on IGF-1 levels (Harper-Wynne et al., 2002) showed a non significant increase in IGF-1 and are in line with earlier studies showing increases in IGF-1 levels with other aromatase inhibitors (Bajetta et al., 1997) (Frost et al., 1996).

A case control study found that one year of the retinoid fenretinide reduced IGF-1 by 13% in women aged <50 and by 3% in women over 50 with changes maintained for up to 5 years (Decensi et al., 2001). However, other trials comparing the retinoid fenretinide with tamoxifen found only 2% lowering in IGF-1 with Fenretinide vs 15% in tamoxifen and no synergistic effects between the two treatments were seen (Guerrieri-Gonzaga et al., 2006).

3.5 Conclusion

Patients with breast cancer had similar concentrations of IGF-1, IGFBP-1, and IGFBP-3 to those seen in healthy control subjects in this pilot study. Tamoxifen can increase IGFBP-1, IGFBP-3, and reduce the IGF-1 to IGFBP-3 ratio, and these alterations may contribute to the therapeutic benefits of tamoxifen. The long term effects of tamoxifen use on IGF status remain to be established.

Longer follow up with larger numbers of patients would determine when, and for how long, tamoxifen can reduce circulating IGF-1. The effect of other treatments for breast cancer, including lower dose tamoxifen and other SERMS is now being tested. Given the increase in risk of breast cancer associated with higher levels of IGF-1 in pre-menopausal women and the ability of phytoestrogens to act as SERMS, the potential of these dietary compounds to alter IGF status like tamoxifen and other SERMs is investigated in the following chapters.

CHAPTER FOUR – SHORT-TERM PHYTOESTROGEN

SUPPLEMENTATION IN HEALTHY FEMALE VOLUNTEERS –

PILOT STUDY

4.1 Introduction

Phytoestrogens have been proposed to protect against many diseases including hormone-dependent cancers and coronary heart disease. Found in many plant products, but particularly soy, legumes, and vegetables, they have a wide variety of proposed biochemical effects including estrogenic and antioxidant effects (Bingham et al., 1998).

Many studies have examined the effect of phytoestrogen supplementation *in vitro*, but fewer have looked at their physiological effects *in vivo*. A summary of human intervention studies examining effects of phytoestrogen supplementation on IGF-1 levels can be found in table 1.3.5.1. As a preliminary pilot *in vivo* study, we have examined the effects of a single intake of phytoestrogens and a week-long intervention. We have assessed a variety of biochemical endpoints, including phytoestrogen metabolites in urine, antioxidant status and lipid status. These are all potentially altered by phytoestrogen consumption (Bingham et al., 1998). We have also examined the effects of phytoestrogen supplementation on insulin-like growth factor (IGF) profile as there is increasing evidence that high circulating levels of IGF-1 and possibly BP3 are a risk factor for pre-menopausal breast cancer (Del Giudice et al., 1998; Hankinson et al., 1998; Johansson et al., 2004; Krajcik et al., 2002; Li et al., 2001; Renehan et al., 2006; Renehan et al., 2004; Shi et al., 2004; Toniolo et al., 2000; Yu et al., 2002). As shown in the previous chapter, levels of IGF-1 and its related binding proteins are altered by

tamoxifen (Campbell et al., 2001), and phytoestrogens have a similar structure to tamoxifen (Bingham et al., 1998).

We have also examined the effects of phytoestrogens on DNA damage, as assessed by the COMET assay. Phytoestrogens have antioxidant properties (Bingham et al., 1998) and may therefore protect against oxidative DNA damage.

The aim of this pilot feeding study was therefore to compare the effects of acute (single 80 mg load) versus chronic (80 mg/day for 7 days) administration of phytoestrogens on a number of physiological endpoints including urinary phytoestrogen metabolites, lipids, antioxidant status, DNA damage and IGF-1, IGFBP-1 and IGFBP-3, in healthy female volunteers (n=10).

This was a feasibility study used to establish the assays for analyses of serum IGF-1, BP1 and BP3 concentrations and GC-MS analysis of phytoestrogens in urine and to determine the practicalities of use of the phytoestrogen-rich 'energiser bar' for future feeding studies.

Study published in (Woodside et al., 2006)

4.2 Materials and methods

4.2.1 Study Design

Subjects were healthy female volunteers (n=10), employed within University College London. The study was fully explained to the subjects, and written informed consent obtained. This study received ethical approval from the Royal Free and University College London Medical School Research Ethics Committee.

The supplement used was a soy and linseed bar, each bar containing a total of 20mg of isoflavones and lignans. The nutritional information is shown in **Table 4.2.1.1.**

The ingredients list was as follows: Dehulled Split Soya Beans, Glucose Syrup, Sugar, Golden Linseeds, Rye Flour, Crisped Rice, Dried Apple, Rolled Oats, Fructose, Corn Syrup Solids, Water, Flavourings and Pectin.

Table 4.2.1.1: Estimated nutritional content of phytoestrogen supplement

	per 100g	per bar	Per day
Energy (kJ)	1656	497	1988
Energy (Kcal)	395	119	476
Protein (g)	15.1	4.5	18
Carbohydrate (g)	56.7	17.0	68.0
of which sugars (g)	29.2	8.8	35.2
Fat (g)	12.0	3.6	14.4
of which saturates (g)	1.9	0.6	2.4
Fibre (g)	4.2	1.3	5.2
Sodium (g)	0.1	Trace	0.03

The study was divided into two phases with a wash-out period of 6 days between phases. Subjects were instructed to follow a soy-free diet for 1 week before the study began.

4.2.2 Phase 1

Subjects were asked to provide a 24-h urine collection prior to the study. After an overnight fast, subjects gave a baseline blood sample. They were then given a breakfast containing 80 mg of phytoestrogens (4x supplement bars). Blood samples were taken at 8 and 24 h (fasting) after the meal with spot urine samples collected during the hours 1-2, 2-4, 4-6, 6-8 and 24 h. Meals or snacks, excluding soy products, were permitted throughout the 8 h post meal period, but only liquids were consumed until 2 h after the meal.

4.2.3 Phase 2

Six days later, subjects collected a further 24-h urine sample and had a further fasting baseline blood sample prior to the study. They were then asked to take 80 mg Phytoestrogen (4x supplement bars) daily for a period of 7 days, after which they collected a final 24-h urine sample and had a final fasting blood sample taken.

4.2.4 Collection and preparation of biological samples

Urine samples (24 h) were stabilised with 1-2 g/l ascorbic acid and sodium azide (0.1 g/L) and stored at -80°C until analysis. Spot urine samples were not stabilised but immediately frozen after collection. Blood samples either kept at 4°C for a maximum of 30 min (plasma) or were allowed to clot for 30 minutes in the dark (serum) and then centrifuged at 3000 rpm for 10 min. Serum or plasma was aliquoted and stored at -80°C until analysis. Lymphocytes were collected into a Vacutainer Cell Preparation Tube (Becton Dickinson, New Jersey), extracted and washed according to manufacturer's instructions and stored at -80°C.

4.2.5 Endpoint measurements

Urine samples were analysed for the isoflavonoids genistein and daidzein, the isoflavan equol and the lignans enterolactone and enterodiol levels, using gas chromatography-mass spectrometry following enzyme hydrolysis, extraction into ether and silylation according to Morton et al. (Morton et al., 1994).

Ascorbic acid concentrations were determined in EDTA plasma as described by Vuillemier & Keck (Vuillemier & Keck, 1989). Concentrations of lipid-soluble antioxidants were measured by HPLC in serum according to the method of

Thurnham et al. (Thurnham et al., 1988). Levels of aqueous phase lipid hydroperoxides were assessed in plasma by the FOX 1 (ferrous oxidation in xylenol orange) assay according to Wolff (Wolff, 1994). Susceptibility of LDL to oxidation was assessed in Phase 2 according to McDowell et al. (McDowell et al., 1995). DNA damage was analysed by the COMET assay in both lymphocytes according to Spanswick et al. (Spanswick et al., 1999). Serum total cholesterol was estimated using an enzymatic CHOD-PAP kit, while serum triglycerides were measured using the Peridochrom GPO-PAP kit (both Boehringer Mannheim). Precipitation for HDL-cholesterol estimation employs phosphotungstic Mg^{2+} reagents. Serum samples were analysed for IGF-1 and IGFBP-1 and IGFBP-3 using ELISA kits from Diagnostic System Laboratories (London, UK). For all laboratory methods, daily quality controls were utilised to ensure both within- and between-assay reproducibility.

See Chapter two for further detail on Methods.

4.2.6 Statistical methods

Comparisons between subjects pre- and post-intervention were carried out using a non-parametric Wilcoxon signed rank test due to the small numbers of subjects and the non-normal distribution of the data. Associations between continuous variables were tested using Spearman correlation coefficients. All analyses were carried out using SPSS.

4.3 Results

Subjects had a mean age of 33.1 years ranging from 23 to 50. Eight were premenopausal, of whom three were on the oral contraceptive pill. One of the

subjects was post-menopausal and was also on hormone replacement therapy. All subjects were non-smokers.

Concentrations of genistein, daidzein, equol, enterolactone and enterodiol were elevated, both after a single 80 mg dose (**Table 4.3.1**) and after 1 week of supplementation (**Table 4.3.2**).

The effect of phytoestrogen supplementation on lipid and antioxidant-related endpoints is shown in **Table 4.3.3**. Neither acute nor chronic phytoestrogen supplementation had any effect on any of the lipid or antioxidant-related endpoints assessed.

All IGF-1 and IGFBP-3 levels fell within the range of expected values given for healthy females in this age group. As fasting samples are required for these assays, measures of IGF-1 and IGFBP-3 status were limited to baseline and 24-hour post supplement in the first phase of the study.

Levels of IGF-1 were unchanged following the single 80 mg load. However, in Phase 2, IGF-1 and IGFBP-3 concentrations were significantly elevated after a week of supplementation. Levels of IGFBP-1 were not altered by phytoestrogens in either phase of the study (**Table 4.3.4**).

The change in IGF-1 in Phase 2 was significantly associated with the change in IGFBP-3 ($r=0.721$, $P<0.05$; Spearman correlation coefficient) (**Figure 4.3.1**).

However, if the ratio of IGF-1/IGFBP-3 was compared pre- and post-supplementation, the ratio was also significantly increased post-supplementation, suggesting that IGF-1 has increased independently of the increase seen in IGFBP-3 (**Table 4.3.4**).

Table 4.3.1: Effect of one phytoestrogen meal on urinary phytoestrogen excretion- Phase 1

Baseline		1–2 h	2– 4 h	4– 6 h	6– 8 h	24 h
[median (IQ range)]		[median (IQ range)]	[median (IQ range)]	[median (IQ range)]	[median (IQ range)]	[median (IQ range)]
Genistein (ng/ml)	196 (146, 308)	217 (169, 377)	313 (201, 416)*	392 (344, 558)*	1257 (1013, 2711)*	289 (139, 417)
Daidzein (ng/ml)	428 (236, 593)	484 (388, 714)*	642 (375, 854)*	952 (681, 1837)*	3283 (2463, 4365)*	408 (252, 634)
Equol (ng/ml)	91 (48, 192)	104 (31, 268)	191 (25, 328)	356 (36, 523)*	757 (207, 915)*	136 (42, 264)
Enterolactone (ng/ml)	284 (105, 334)	280 (149, 307)	226 (153, 308)	242 (146, 305)	301 (206, 335)	284 (169, 461)
Enterodiol (ng/ml)	69 (22, 101)	66 (42, 88)	79 (47, 88)	77 (40, 95)	75 (48, 95)	77 (46, 145)

* P<.05 when compared to baseline, Wilcoxon signed rank test.

Table 4.3.2: Effect of 1 week of phytoestrogen supplementation on urinary phytoestrogen excretion – Phase 2

Baseline		Postweek
[median (IQ range)]		[median (IQ range)]
Genistein (ng/ml)	206 (88, 306)	2775 (1818, 3101)*
Daidzein (ng/ml)	385 (195, 599)	4571 (3934, 5803)*
Equol (ng/ml)	96 (53, 131)	899 (501, 1540)*
Enterolactone (ng/ml)	224 (95, 403)	392 (181, 565)*
Enterodiol (ng/ml)	46 (19, 81)	83 (58, 114)*

*P<.05 when compared to baseline 2, Wilcoxon signed rank test.

Table 4.3.3: Effect of phytoestrogen supplementation on lipids and antioxidant status

	Phase 1 [median (IQ range)]			Phase 2 [median (IQ range)]	
	Baseline	8 h	24 h	Baseline	postweek
Total cholesterol (mmol/L)	4.59 (4.07, 5.31)	4.47 (4.05, 4.82)	4.19 (3.89, 4.79)	4.63 (4.16, 5.54)	5.00 (4.32, 5.27)
HDL cholesterol (mmol/L)	1.49 (0.91, 1.68)	1.43 (0.96, 1.62)	1.43 (0.93, 1.87)	1.48 (1.16, 1.58)	1.57 (1.10, 1.73)
Triacylglycerol (mmol/L)	1.00 (0.83, 1.58)	1.44 (1.31, 2.39)	0.92 (0.66, 1.20)	1.19 (0.84, 1.78)	1.19 (0.86, 1.45)
LDL cholesterol (mmol/L)	2.59 (2.24, 3.39)	2.27 (2.13, 2.86)	2.36 (2.16, 2.51)	2.86 (2.29, 3.41)	2.96 (2.70, 3.31)
FOX1 (μmol/L)	1.65 (1.15, 2.57)	1.42 (1.37, 1.64)	1.20 (1.10, 1.37)	1.35 (1.19, 1.45)	1.34 (1.29, 1.46)
Retinol (μmol/L)	1.00 (0.74, 1.49)	0.79 (0.48, 1.47)	1.01 (0.69, 1.30)	0.99 (0.61, 1.10)	0.91 (0.52, 1.19)
α-Tocopherol (μmol/L)	20.9 (17.6, 23.9)	19.1 (14.8, 22.8)	21.4 (16.3, 24.8)	18.9 (15.5, 24.2)	21.0 (12.6, 23.3)
Lycopene (μmol/L)	0.09 (0.04, 0.16)	0.09 (0.06, 0.11)	0.09 (0.06, 0.11)	0.06 (0.04, 0.09)	0.07 (0.03, 0.16)
α-Carotene (μmol/L)	0.14 (0.06, 0.29)	0.12 (0.06, 0.21)	0.15 (0.09, 0.27)	0.08 (0.04, 0.11)	0.07 (0.05, 0.41)
β-Carotene (μmol/L)	0.14 (0.09, 0.20)	0.13 (0.09, 0.18)	0.16 (0.08, 0.19)	0.12 (0.09, 0.16)	0.10 (0.07, 0.17)
DNA damage (olive tail moment)	—	—	—	4.21 (1.50, 5.07)	2.11 (1.31, 2.74)
Lag time (min)	—	—	—	44.2 (37.4, 53.6)	44.5 (38.8, 48.4)

Table 4.3.4: Effect of phytoestrogen supplementation on insulin-like growth factor and binding protein concentrations

Phase 1 [median (IQ range)]			Phase 2 [median (IQ range)]	
	Baseline	24 h	Baseline 2	Postweek
IGF-1 (ng/ml)	218 (151, 293)	250 (179, 310)	155 (123, 258)	265 (228, 360)*
IGFBP-1 (ng/ml)	22.6 (19.0, 36.8)	17.4 (15.8, 34.6)	24.8 (14.6, 57.7)	14.7 (10.0, 36.0)
IGFBP-3 (ng/ml)	4564 (3954, 5306)	4110 (3887, 4533)	3735 (3631, 4196)	4420 (4192, 4935)*
IGF-1/IGFBP-3 ratio	0.05 (0.04, 0.06)	0.06 (0.04, 0.07)	0.04 (0.03, 0.06)	0.06 (0.05, 0.07)*

*P<.05 when compared to baseline, Wilcoxon rank signed test.




Figure 4.3.1: Association between change in IGF-1 and IGFBP-3 over 1 week of supplementation ($r=0.721$, $P<0.05$; Spearman correlation coefficient).

4.4 Discussion

This study has assessed the effect of phytoestrogen supplementation, in the form of a soy, rye and linseed bar, on a variety of biochemical endpoints.

Concentrations of phytoestrogen metabolites in urine reached their highest levels 6-8 h after supplementation for isoflavones, and 24 h for lignans in phase

1. This is generally in agreement with other studies examining the effect of supplementation on urinary excretion of phytoestrogens (King & Bursill, 1998); (Rowland et al., 2003; Setchell et al., 2003). This study, however, was not designed to be a rigorous study of phytoestrogen pharmacokinetics.

Phytoestrogens can act as antioxidants (Bingham et al., 1998) and may inhibit the oxidation of LDL (Lissin & Cooke, 2000) although the data from this study has not confirmed this in terms of lag time to oxidation. A protective/antioxidant effect has been shown by phytoestrogen supplementation on the susceptibility of LDL to oxidation in another study (Tikkanen et al., 1998). Another study, however, in common with this one, has failed to show an inhibition of lipid peroxidation after 8 weeks of isoflavonoid supplementation in subjects with high-normal blood pressure (Hodgson et al., 1999), measuring urinary F-2-isoprostanes a reliable indicator of oxidative stress. The study measuring F-2-isoprostanes was randomised, double-blind and placebo-controlled, in contrast to the present study and was carried out in relatively large numbers (n=59). Our subjects were healthy and therefore had normal antioxidant status. Therefore the lag times of these subjects may not have been easily affected by this supplement.

We have also observed an apparent reduction in DNA damage in lymphocytes (Olive tail moment from 4.21 to 2.11) after 1 week of phytoestrogen supplementation, although this was not significant. We have previously shown

that genistein and equol within physiological range can reduce oxidative DNA damage *in vitro* in lymphocytes (Sierens et al., 2001). Mitchell & Collins (Mitchell & Collins, 1999) carried out a 4 week feeding study in healthy men to assess the effects of a phytoestrogen supplement in the form of soy milk on levels of DNA damage. There was no significant effect on H₂O₂-induced DNA damage in lymphocytes in those receiving the soy milk supplement as assessed by the comet assay. However, the levels of oxidised pyrimidine base damage over the 4-week period progressively decreased. A recent study found no effect of rye crispbread (rich in lignans secoisolariciresinol and matairesinol) over 2 weeks on oxidative DNA damage (Pool-Zobel et al., 2000). We have only evaluated endogenous DNA damage in this study, where the damage may or may not have been oxidative in origin, and our intervention period was shorter than that which previously showed an effect. The antioxidant potential of these foods and compounds *in vivo* clearly requires further investigation.

This study found no effect of either a single dose or week's supplementation with 80 mg phytoestrogens daily on serum lipids. This is in contrast to a meta-analysis of studies published in 1995 (Anderson et al., 1995). Of the 38 studies reviewed, 89% reported a net decrease in plasma cholesterol that averaged around 9% for an intake of around 47 g soy protein daily. However, it is notable that 77% of the studies included in the meta-analysis had 95% confidence intervals that included zero.

Since the meta-analysis, several well-controlled studies have been carried out, and the results of these have been variable (Lichtenstein, 2001). There are several potential explanations for this variability. Merz-Demlow et al. (Merz-Demlow et al., 2000) suggest that the effect of isoflavones on lipids may be related to menstrual cycle phase. The food matrix may also be important, as

Lichtenstein (Lichtenstein, 2001) suggests that the effect of isoflavones appears to be somewhat dependent on whether the isoflavones are ingested in isolation or as an enriched preparation of those endogenously present in soy.

A recent study observed a reduction in LDL cholesterol in 156 men and women by an average of 6%. This was due entirely, however, to a mean reduction of 9% among those with a raised baseline LDL cholesterol (>4.29 mmol/l) and was only observed in men and post-menopausal women (Crouse et al., 1999). The original meta-analysis also found a greater lowering effect in those with initially raised total cholesterol levels (Anderson et al., 1995). It may be that we observed no difference in our largely pre-menopausal sample set because they had low-normal lipid status initially, but it is more likely that one week of supplementation was too short to demonstrate an effect.

This study has shown that 1 week of phytoestrogen supplementation can alter IGF status, producing a significant increase in both IGF-1 and IGFBP-3 levels. The vast majority of IGF-1 is carried through the circulation bound to a complex containing IGFBP-3 and, until released by proteolysis, is relatively unreactive (Rajaram et al., 1997). Thus the strong correlation between increasing levels of IGF-1 and IGFBP-3 was as expected. However, when the ratio of IGF-1 to BP-3 was considered, this ratio was significantly increased post-supplement, and this finding requires further elucidation. Tamoxifen, a selective estrogen receptor modulator (SERM) has been shown to lower IGF-1 and increase IGFBP-3 concentrations (Campbell et al., 2001; Helle et al., 1996); therefore the phytoestrogen supplement appears to be behaving differently to tamoxifen. Oral estrogen treatment for menopausal women has also been shown to reduce IGF-1 (Kelly et al., 1993) through inhibition of hepatic IGF-1 synthesis. The mechanism by which this supplement has increased IGF-1 requires further

study. Since conducting this study in 1998, other intervention studies have also published their findings on the effects of soy supplementation on IGF status **(Table 1.3.5.1)**.

Of the twelve studies published to date, half have seen a significant increase in serum IGF-1 concentrations while the others have seen no effect **(Table 1.3.5.1)**.

Two studies were conducted using soy protein with a milk protein control. In men, serum IGF-1 was higher in those supplemented with soy protein than those consuming milk protein (Khalil et al., 2002). The same supplements given to post-menopausal women (Arjmandi et al., 2003) resulted in significant increases in IGF-1 on both soy protein ($p < 0.0001$) and the milk protein control ($p = 0.0045$). However like the men, the greatest increases in IGF-1 (97%) was seen in women on the soy protein especially those who were not on HRT. These two studies would suggest that protein itself does increase IGF-1 but that soy protein had an effect on IGF-1 independent of its protein content. Calcium content did not explain the findings as both supplements had similar calcium content.

By contrast, in a three month study looking at markers of bone turnover (Wangen et al., 2000) in pre-menopausal women, IGF-1 and IGF-BP3 were increased by a low isoflavone diet, while in post-menopausal women, there were trends towards decreased IGF-1 and IGF-BP3 concentrations with increasing isoflavone concentration. The authors concluded that, although soy isoflavones do affect markers of bone turnover, the changes observed were of small magnitude and were not likely to be clinically relevant. In a study looking at soy protein supplementation with or without 83 mg isoflavones (Adams et al., 2003), it was found that changes in serum IGF-1 and BP3 were similar in both

groups and there appeared to be no effect of soy isoflavones on IGF-1 when compared to the protein supplement alone. Similarly, a one year soy protein supplementation in post-menopausal women (n=62) (Arjmandi et al., 2005), found that both soy and dietary protein supplements significantly increased serum IGF-1, however only those consuming soy protein had a significant increase in IGF-BP3.

In a 2 year soya intervention (~30mg isoflavones) in pre-menopausal women (Maskarinec et al., 2005b), there was no significant change in serum IGF-1 or BP3 throughout. They did see a positive correlation between total isoflavone excretion and IGF-1 ($p=0.004$). The author concluded that adding soya foods to the diet of pre-menopausal women does not appear to lower serum levels of IGF-1 and BP-3, if anything the greater protein intake from soy may lead to a small increase in IGF-1 serum levels.

In all of the studies mentioned above it is difficult to disentangle the effects of isoflavones from soy protein (Voskuil et al., 2005) especially in those studies which have looked at soy supplementation as part of an overall healthy dietary intervention (Gann et al., 2005; Kaaks et al., 2003; Spentzos et al., 2003).

Only our study (discussed in chapter four) (Campbell et al., 2004) and, to our knowledge, one subsequent study published recently (Vrieling et al., 2007), has looked at the effects of isoflavone supplementation on its own.

The results from this study and the intervention studies above are inconclusive but possibly show opposing effects of isoflavones and soy protein, with IGF-1 decreased by isoflavones but increased by soy protein and possibly protein in general. Certainly feeding studies (Clemmons et al., 1985; Smith et al., 1995) have shown that a diet restricted of certain essential amino acids reduces IGF-1 levels possibly by decreasing production of GH which in turn reduces hepatic

production of IGF-1 (Ketelslegers et al., 1995) and increases BP-1 (Jousse et al., 1998). It has been observed previously that IGF is closely linked to nutritional status (Clemmons & Underwood, 1991). Our observation of an increase in both IGF-1 and IGF-BP3 could well be attributed to the increase in kilocalories and protein consumed in the phytoestrogen bar (476 Kcal daily; 18 g protein daily; Table 1), as IGF-1 appears to be closely related to protein and energy metabolism (Clemmons & Underwood, 1991). No dietary intake data were collected in the present study so it is unclear whether total protein and energy intake increased or whether subjects unconsciously decreased their intake to compensate for the extra calories over the week. Intervention studies using a phytoestrogen extract or, as a control, a food supplement from which phytoestrogens have been removed, should be used in future to exclude the possibility that an increase in protein and/or energy intake produced the observed alteration in IGF-1 and IGF-BP3.

The status of endogenous oestrogen concentrations may also contribute to the differences seen with IGF-1 – this was a heterogeneous study population including pre and post-menopausal women, HRT vs none, those on the oral contraceptive pill, etc. Thus it may also be important to take into account the effects of the menstrual cycle in future studies as it has been shown that IGF-1 can vary throughout the menstrual cycle (Caufriez et al., 1986; Helle et al., 1998; Juul et al., 1997; Lavigne et al., 2004).

Oestrogenic, and therefore confounding, effects of oral contraceptives and hormone replacement therapy should also be considered as numbers were not large enough in this study to examine their effect.

The supplement bar we used contained two classes of phytoestrogen – isoflavones (found predominantly in soy) and lignans (found predominantly in

linseed)(Bingham et al., 1998). The biological effects of these two classes may be different, and research has, to date, concentrated on the isoflavones.

However, some recent studies have assessed lignan intake and would appear to show a protective effect on breast cancer risk (Keinan-Boker et al., 2004; Thanos et al., 2006; Touillaud et al., 2007). From this study, we cannot distinguish the effect of the two different classes, or indeed, conclusively attribute the biological effects we have observed to the phytoestrogens independent of any other supplement constituent.

Nevertheless these findings may have future implications for health and disease. Phytoestrogens have been proposed to protect against breast cancer, as Japanese women, who consume their traditional diet with a high intake of phytoestrogens, have a very low incidence of breast cancer (Woodside & Campbell, 2001). IGF-1 has been shown to be a risk factor for breast cancer in pre-menopausal women (Del Giudice et al., 1998; Hankinson et al., 1998; Johansson et al., 2004; Krajcik et al., 2002; Li et al., 2001; Renehan et al., 2006; Renehan et al., 2004; Shi et al., 2004; Toniolo et al., 2000; Yu et al., 2002) and, if it can be confirmed that phytoestrogens increase circulating IGF-1 concentrations, then this would actually indicate an increased risk of breast cancer in those consuming high-phytoestrogen diets. Obviously further work would be required before such conclusions can be drawn, but the data presented here do highlight the need for further, well-designed trials. Some of the issues raised above have been taken into consideration in the study design described and tested in chapter 5.

In conclusion, this small study suggests that one week of supplementation with a phytoestrogen-rich bar alters concentrations of IGF-1 and IGF-BP3. The supplement also seemed to protect DNA against damage in lymphocytes, but

had no effect on other antioxidant parameters or lipid status in healthy female subjects. Further larger and long-term intervention studies using isolated isoflavone supplementation, which therefore allow the inclusion of a placebo control, should determine whether the isoflavone content of soy has an effect on IGF status.

CHAPTER FIVE – THE EFFECT OF ISOFLAVONE
SUPPLEMENTATION ON INSULIN-LIKE GROWTH FACTOR,
LIPID AND ANTIOXIDANT STATUS IN HEALTHY FEMALE
VOLUNTEERS

5.1 Introduction

High concentrations of IGF-1 in serum have been associated with an increased risk of breast cancer, in case control studies (Bohlke et al., 1998; Bruning et al., 1995; Del Giudice et al., 1998; Johansson et al., 2004; Li et al., 2001; Peyrat et al., 1993; Yu et al., 2002) and more recently in prospective studies (Hankinson et al., 1998; Krajcik et al., 2002; Muti et al., 2002; Rinaldi et al., 2006; Toniolo et al., 2000). Most reviews agree that this relationship is more prominent in pre-menopausal women (Fletcher et al., 2005; Renehan et al., 2004; Schernhammer et al., 2005; Shi et al., 2004; Sugumar et al., 2004) but not all studies comply (Kaaks et al., 2002; Rinaldi et al., 2006; Rinaldi et al., 2005b), although a recent prospective analysis of the Nurses Health study II cohort study found no associations in either pre or post-menopausal women (Schernhammer et al., 2006).

Tamoxifen has been shown to lower circulating IGF-1 levels by approximately 30% in breast cancer patients (Campbell et al., 2001; Pollak et al., 1990), (Colletti et al., 1989; Helle et al., 1996; Lonning et al., 1992) and in healthy women (Bonanni et al., 2001; Decensi et al., 1998; Decensi et al., 2003).

Isoflavones are a class of phytoestrogens which may have a variety of potential health benefits in endocrine-responsive cancers (Adlercreutz, 2002b), (Adlercreutz, 2002a) and CVD (Cassidy & Hooper, 2006), (Bingham et al.,

1998); (This et al., 2001). They are obtained in the diet predominantly from soy products (Bingham et al., 1998). Isoflavones are structurally similar to tamoxifen and, like tamoxifen, can act as selective oestrogen receptor modulators (Brzezinski & Debi, 1999). They may therefore be able to alter concentrations of IGF-1 and its binding proteins.

The aim of this study was to determine whether one month of isoflavone supplementation could alter serum concentrations of IGF-1 and its related binding proteins. In light of the results in the previous chapter, where a phytoestrogen-rich bar providing 80 mg phytoestrogens/day over 7 days produced an increase in IGF-1 and IGFBP-3, this study used a phytoestrogen extract, which allowed the use of a placebo control. The effect of the menstrual cycle on insulin-like growth factor profile was accounted for in the study design. We also assessed whether the isoflavone supplement produced an alteration in hormone, lipid or antioxidant status as effects on these parameters have been demonstrated in some studies (Merz-Demlow et al., 2000; Takatsuka et al., 2000), but not in others (Howes et al., 2000; Jenkins et al., 2000; Nestel et al., 1999; Simons et al., 2000). The inclusion of a food frequency questionnaire and a 7-day food diary completed in each phase allowed the assessment of nutritional intake.

Study published in (Campbell et al., 2004).

5.2 Materials and methods

5.2.1 Study design

The study was based on a randomised, double-blind, placebo-controlled cross-over design (Figure 4.2.1). Pre- and post-menopausal subjects were randomised to receive either two placebo or two supplement tablets containing

a total of 80 mg isoflavones daily. Each phase was followed by a minimum 2-month washout period and the alternative intervention.




Figure 5.2.1 Study design – for pre-menopausal women blood samples were taken throughout the menstrual cycle as shown. For post-menopausal women, fasting blood and urine were only collected at the start and end of the 28 day intervention period.

5.2.2 Eligibility Criteria

Pre-menopausal - not on oral contraceptives, no antibiotics within previous 4 months, regular menstrual cycle of approximately 28 days.

Post-menopausal - not on hormone replacement therapy, no antibiotics within previous 4 months, had an oophorectomy or amenorrhoea for at least 12 months.

5.2.3 Sample Collection

Fasting blood samples were obtained from post-menopausal subjects at baseline and at 28 days in each phase of the study. Fasting blood samples

were collected at baseline, day 1-3, day 6-8, day 12-15, day 21-23 and day 26-28 from pre-menopausal subjects to assess the effects of the menstrual cycle on the endpoints measured. For all subjects, a 24-h urine sample was collected at baseline and 28 days for each phase. In addition, a fully validated food frequency questionnaire was filled in at the baseline stage of each phase and a food diary completed over the last seven days of each phase. All biological samples were stored at -80°C until analysis. The study was approved by the Royal Free and University College London Ethical Committee.

5.2.4 Endpoint measurements

A variety of endpoint measures were assessed. Levels of IGF-1, IGF-BP1, IGF-BP3 were measured in serum using ELISA methods (Diagnostic Systems Laboratories). Oestradiol, progesterone, LH, FSH, SHBG were measured in serum using routine IRMA/RIA methods. Isoflavones were assessed in urine at the start and end of each supplementation period by GC-MS according to (Morton et al., 1994). Serum lipids were assessed, again at the start and end of each supplementation phase, using standard spectrophotometric assays.

Antioxidant status was assessed by the measurement of vitamins A and E and carotenoids (Craft, 1992), malondialdehyde (Young & Trimble, 1991), vitamin C (Vuillemier & Keck, 1989), and aqueous phase lipid hydroperoxides (Wolff, 1994).

Fully validated EPIC food frequency questionnaires (FFQ) and 7-day food diaries (FD) (McKeown et al., 2001) were analysed using DietPlan 5 for Windows 98.

See Chapter two for further detail on Methods.

5.2.5 Supplement

The supplement used was a proprietary product called Promensil donated by Novogen. Each 500 mg supplement tablet contained 215 mg of an extract of *Trifolium pratense* (red clover) leaf. This extract contained 40 mg/tablet of total isoflavones present as hydrolysed aglycones. Each tablet contained 4.0 mg genistein (4',5,7-trihydroxyisoflavone), 3.5 mg daidzein (4',7-dihydroxyisoflavone), and their methylated precursors 24.5 mg biochanin A (4'-methoxy-5,7-dihydroxyisoflavone) and 8.0mg formononetin (4'-methoxy-7-hydroxyisoflavone). The residual material in each tablet was predominantly flavonoids, principally chlorophylls a and b. Other ingredients were as follows: Calcium hydrogen phosphate, Cellulose-microcrystalline, Soy polysaccharide, Natural vitamin E, Magnesium stearate, Silica, hydroxylpropyl methyl cellulose, Rosemary extract and iron oxide. The placebo had the same physical properties (appearance and taste) and excipient ingredients but did not contain the isoflavone extracts.

5.2.6 Statistical analyses

The intervention was double-blinded and was not un-blinded until all analyses, including statistical analyses had been completed.

Pre- and post-menopausal women were considered separately in the analyses.

Changes from baseline were compared between treatment groups using a paired samples t-test (for two time-point comparisons) or repeated measures ANOVA (for multiple time-point comparisons). The association of continuous variables was assessed using Pearson correlation coefficients. For the pre-menopausal subjects, at the 5% level of significance, the study had 80% power

to detect as statistically significant a 35% difference in change in IGF-1 between the treatment groups over the supplementation period.

5.3 Results

A total of 16 pre- and 7 post-menopausal subjects completed both phases of the study. The baseline characteristics of the subjects are shown in **Table**

5.3.1. Among pre-menopausal subjects, three out of 16 were parous and among post-menopausal subjects, five out of six were parous. The mean (range) number of wash-out days was 95 (56, 206) in pre-menopausal subjects and 70 (62, 91) in post-menopausal subjects.

Table 5.3.1 Baseline characteristics of study subjects

	Pre-menopausal (n=16)	Post-menopausal (n=7)
Age	34 (9)	57 (6)
BMI	23.3 (4.3)	24.4 (5.0)
Waist to hip ratio	0.75 (0.04)	0.77 (0.05)
Protein (E%)	14.1 (2.5)	14.3 (1.6)
Carbohydrate (E%)	41.8 (6.8)	47.0 (7.6)
Fat (E%)	38.8 (4.9)	35.8 (3.8)
Alcohol (E%)	5.3 (5.1)	3.0 (3.9)

Data presented as mean (SD). E% = percentage of total calories

5.3.1 Isoflavones

The change in concentration of genistein, daidzein and equol over the supplementation period was significantly larger during the supplement phase compared to placebo in both pre- and post-menopausal women (**Table 5.3.1.1**)

Concentrations excreted on the placebo were comparable to normal ranges in other studies (Cassidy, 2006). Concentrations of genistein, daidzein and equol excreted in urine, in pre-menopausal subjects, did not change during the placebo phase, but increased significantly during the supplement phase (**Fig 5.3.1.1**). This was also the case in post-menopausal subjects (**Fig 5.3.1.2**).

However, the concentrations of genistein in postmenopausal subjects (**Table 5.3.1.2**) increased both on placebo (Mean±SD placebo (baseline 258.6±174.6 versus 28 days 659.2±388.5 ng/ml; $p<0.05$)) and on supplement (Mean±SD supplement baseline 749.9±374.7 ng/ml versus 28 days 3959.5±2194.4; $p<0.01$)). However, the magnitude of increase in genistein was much higher on the supplement (5.3-fold) than when on placebo (2.5-fold) and would appear to be because of a very low baseline level for genistein during the placebo phase, compared to the baseline level at the start of the supplement phase.

The placebo increase seen in post-menopausal women may have been caused by a carry over effect. Subjects who completed the supplement phase first did show higher genistein concentrations in the placebo phase nevertheless there was no statistical difference between the two groups when compared with the Mann Whitney U test (Mean±SD placebo first (placebo baseline 170.93± 65.05) and on supplement first (placebo baseline 375.6± 221.92)).

Concentrations of the two lignans, enterolactone and enterodiol, did not change during either phase, which is to be expected as neither of these compounds were present in the supplement.

The metabolite equol, was significantly elevated on the supplement though the increase varied greatly between individuals ranging from 0.0 - 4172 ng/ml. A comparison of mean urine isoflavone concentrations post-supplement with other studies, shows that equol excretion varies considerably more than the other isoflavones (between 8 and 93 fold difference) (**Table 5.3.1.3**).

Subjects with urinary equol concentrations post-isoflavone supplement of >1µg/ml were defined as equol excretors. As defined above, 23% (3/14) pre-menopausal subjects were equol excretors compared to 20% (1/5) of post-menopausal subjects. Post-supplementation, Equol excretion ranged from 0-946 ng/ml in the non-excretors to 1197-4172 ng/ml in the excretors. When the four excretors were compared to the 15 non-excretors using an independent samples t test, levels of MDA, triglycerides, γ-tocopherol and α-tocopherol (at baseline) were significantly higher in the excretors. There were no significant differences seen in the FFQ or the food diaries of these subjects. There were also no significant differences in alterations of IGF-1 and BP-3 or BP-1 concentrations in these two groups when comparing mean changes on placebo or supplement (**Table 5.3.1.4**).

Table 5.3.1.1. Effect of supplementation on urinary excretion of isoflavone metabolites in pre-menopausal subjects.

	Placebo		Supplement	
	Day 1-3	Day 26-28	Day 1-3	Day 26-28
Genistein ng/ml	254.8(212.8)	487.9(386.6)	567.2(458.2)	2756.1(2248)*
µg/day	436.5(375.3)	967.3(1064.7)	1097.4(999.5)	4641.9(3599.7)*
ng/µmol creatinine	41.6(38.2)	86.4(81.4)	106.2(93.7)	445.4(282.2)**
µmol/mol creatinine median (interquartile range)	127.2 (45.4,266.9)	186.0 (137.4, 476.5)	319.1 (39.5,731.5)	15323.1(793.7, 2254.5)**
Daidzein ng/ml	411.5(410.6)	228.5(187.5)	216.6(147.0)	3582.5(2107)**
µg/day	740.7(1032.8)	384.8(280.8)	416.9(453.5)	6362.6(5273)**
ng/µmol creatinine	64.6(79.2)	40.3(37.2)	34.9(25.2)	607.3(455.6)**
µmol/mol creatinine median (interquartile range)	147.8 (48.0, 308.7)	102.5 (63.7, 194.2)	94.0 (66.9, 231.4)	1785.7 (925.1, 3416.1)**
Equol ng/ml	94.9(108.9)	68.0(52.4)	162.2(404.6)	911.9(1133.2)
µg/day	118.4(134.1)	121.1(110.4)	167.5(286.4)	1363.4(1726)*
ng/µmol creatinine	9.22(8.8)	12.5(13.0)	16.27(28.9)	132.7(131.3)*
µmol/mol creatinine median (interquartile range)	20.8 (13.3, 56.9)	49.9 (17.8, 57.5)	32.2 (2.6, 82.8)	443.3 (152.2, 760.9)*
Enterolactone ng/ml	336.8(888.5)	198.1(274.5)	203.0(185.7)	188.3(377.8)
µg/day	542.0(1286)	332.8(474.9)	337.7(312.3)	289.0(597.7)
ng/µmol creatinine	65.1(180.7)	39.7(76.8)	32.3(30.4)	28.4(60.1)
µmol/mol creatinine median (interquartile range)	32.7 (14.4, 131.9)	66.0 (31.4, 110.7)	85.8 (24.1, 168.9)	34.9 (15.9, 86.5)
Enterodiol ng/ml	26.9(30.9)	18.4(18.3)	40.9(62.0)	18.1(18.6)
µg/day	36.7(50.4)	29.6(32.5)	74.6(137.3)	28.0(26.8)
ng/µmol creatinine	3.49(5.7)	3.14(3.54)	6.45(10.8)	2.72(2.33)
µmol/mol creatinine median (interquartile range)	5.8 (4.04, 8.87)	3.6 (1.66, 22.9)	6.17 (3.14, 19.4)	5.31 (4.25, 12.0)

Mean (SD) *p<0.05, **p<0.01 when baseline is compared with post placebo/supplement (paired t test)

Table 5.3.1.2. Effect of supplementation on urinary excretion of isoflavone metabolites in post-menopausal subjects.

	Placebo		Supplement	
	Baseline	Day 28	Baseline	Day 28
Genistein ng/ml	258.6(174.6)	659.2(388.5)*	749.9(374.7)	3959.5(2194)**
µg/day	581.0(376.4)	1362.8(886.3)*	1361.3(579.3)	8696.8(5096)**
ng/µmol creatinine	66.2(56.8)	141.0(89.4)*	131.0(76.8)	1455.5(1610)
µmol/mol creatinine median (interquartile range)	155.3 (104.0, 333.8)	617.0 (141.8, 785.9)*	411.0 (336.9, 517.9)	3538.2(1461.1, 6756.3)*
Daidzein ng/ml	343.7(196.6)	353.4(255.8)	525.5(345.4)	4964.0(3733)*
µg/day	765.4(396.8)	758.5(660.3)	1013.6(801.8)	11083(9432)*
ng/µmol creatinine	85.8(58.7)	77.8(64.6)	102.1(88.4)	1335.6(783.2)**
µmol/mol creatinine median (interquartile range)	259.4 (187.4, 496.6)	195.0 (122.2, 562.3)	301.9 (100.5, 794.2)	5717.0(2464.9, 5932.5)*
Equol ng/ml	63.0(26.7)	78.4(56.4)	86.5(39.4)	705.0(344)*
µg/day	141.6(69.6)	172.5(143.6)	163.2(72.5)	1536.9(823)*
ng/µmol creatinine	13.7(5.47)	17.7(14.2)	12.9(3.08)	256.9(267)
µmol/mol creatinine median (interquartile range)	67.2 (34.5, 73.5)	58.0 (25.5, 127.9)	50.5 (45.3, 63.2)	704.8 (282.3, 2019.6)
Enterolactone ng/ml	166.7(161.3)	142.9(74.8)	169.7(154.2)	167.0(166.5)
µg/day	364.9(334.2)	269.2(128.4)	353.3(361.5)	396.9(410.5)
ng/µmol creatinine	38.14(32.7)	27.9(13.3)	32.8(38.5)	73.7(94.0)
µmol/mol creatinine median (interquartile range)	82.2(38.2, 187.9)	108.8(74.9, 120.6)	59.5(17.2, 216.5)	119.0(21.7, 560.5)
Enterodiol ng/ml	9.95(9.08)	10.0(12.7)	8.94(5.94)	9.19(6.82)
µg/day	22.12(20.9)	22.64(31.2)	17.2(13.8)	20.2(15.4)
ng/µmol creatinine	2.51(2.54)	2.32(3.11)	1.65(1.47)	2.85(2.21)
µmol/mol creatinine median (interquartile range)	7.65(2.56, 9.01)	4.0(2.2, 11.8)	3.90(2.7, 6.25)	6.85(3.05, 18.3)

Mean (SD) *p<0.05, **p<0.01 when baseline is compared with post placebo/supplement (paired t test)




Figure 5.3.1.1 Effect of supplementation on urinary excretion of isoflavone metabolites in pre-menopausal subjects.

Mean \pm SE; * p <0.05, ** p <0.01 when baseline is compared with post-supplement (paired t test)




Figure 5.3.1.2. Effect of supplement on urinary excretion of isoflavone metabolites in post-menopausal subjects.

Mean \pm SE * p <0.05, ** p <0.01 when baseline is compared with post-supplement. (paired t test)

Table 5.3.1.3. Comparison of Isoflavone excretion with other studies

Author		Baseline	Post-supplement	Fold increase
Albertazzi (60g soy powder, 3 months)	Genistein ng/ml	114	2818	25
	Daidzein ng/ml	224	2956	13
	Equol ng/ml	42	989	23
Campbell (80mg Isoflavones, 28 days)	Genistein nmol/d	1478	20561	14
	Daidzein nmol/d	2543	26044	10
	Equol nmol/d	651	5268	8
Karr (23mg Gen, 13mg Daid in soy protein, 9 days)	Genistein nmol/d	225	5139	23
	Daidzein nmol/d	561	9303	17
	Equol nmol/d	449	5900	13
Petrakis (38mg Genistein in soy protein, 5 months)	Genistein µg/d	197	3121	16
	Daidzein µg/d	114	6007	53
Rowland (35mg Gen, 21mg Daid in soy protein, 17days)	Genistein nmol/d	384	10326	27
	Daidzein nmol/d	1060	17969	17
	Equol nmol/d	43	3995	93

Table 5.3.1.4: Comparison of serum lipids, tocopherols and IGF-1, BP-3 and BP-1 concentrations in equol excretors and non excretors.

Nutrients	Non-excretors (n=15)	Excretors (n=4)	P
MDA placebo baseline	1.85 (0.57)	4.13 (1.77)	0.000
Triglycerides placebo baseline	0.665 (0.535)	2.23 (2.13)	0.014
γ -tocopherol placebo baseline	4.00 (0.43)	5.72 (2.30)	0.009
α -tocopherol placebo baseline	21.60 (4.99)	31.44 (12.80)	0.024
IGF-1 Placebo (day 1- day 28)	-27.87 (37.93)	7.32 (19.5)	0.095
IGF-1 Supplement (day 1 – day 28)	-2.95 (49.47)	4.47 (45.51)	0.790
IGFBP-3 Placebo (day 1- day28)	-228.4(344)	-418.6 (225.8)	0.315
BP-3 Supplement (day 1-day 28)	-111.0 (334.2)	-408.2 (257.6)	0.120
IGFBP-1 Placebo (day 1-day 28)	-14.21(30.4)	-5.53 (25.3)	0.609
BP-1 Supplement (day 1-day 28)	-5.64(26.77)	-9.59 (31.02)	0.802

Mean (SD) Independent samples t test

5.3.2 Hormones

Oestradiol, progesterone, LH, FSH and SHBG all followed normal menstrual cycle patterns in pre-menopausal subjects on placebo (**Figures 5.3.2.1-5.3.2.5**), and levels were not significantly different on the isoflavone supplement (**Table 5.3.2.1**) when compared using repeated measures ANOVA. Similarly, there was no significant difference seen between placebo and supplement hormone levels measured in post-menopausal subjects (**Table 5.3.2.2**).

However when hormone concentrations were compared using a paired t test there was a significant increase in SHBG concentrations on day 21-23 in supplement phase vs placebo of pre-menopausal subjects ($P=0.003$).

When hormone levels were compared between groups, pre-menopausal women had higher levels of oestradiol ($P=0.019$) and Progesterone ($P=0.016$) and post-menopausal women had significantly higher levels of LH ($P=0.000$) and FSH ($P=0.001$). No significant difference was seen with SHBG levels in these groups (**Table 5.3.2.3**).




Figure 5.3.2.1: Effect of supplementation on LH levels in pre-menopausal women. Mean (SE)




Figure 5.3.2.2: Effect of supplementation on FSH levels in pre-menopausal women. Mean (SE)




Figure 5.3.2.3: Effect of supplementation on Oestradiol levels in premenopausal women. Mean (SE)




Figure 5.3.2.4: Effect of supplementation on Progesterone levels in premenopausal women. Mean (SE)




Figure 5.3.2.5: Effect of supplementation on SHBG levels in pre-menopausal women. Mean (SE). *P=0.003 paired t test supplement vs placebo (Day 21-23).

Table 5.3.2.1: Effect of isoflavone supplementation on circulating hormone levels in pre-menopausal women. Mean \pm (SD).

		Baseline	Day 6-8	Day 12-14	Day 21-23	Day 26-28
LH (U/L)	Placebo	2.5 (1.0)	2.7 (1.6)	6.6 (4.6)	3.1 (3.0)	2.2 (1.9)
	Supplement	2.5 (1.4)	2.5 (0.8)	9.1 (8.8)	2.5 (2.3)	1.6 (1.0)
FSH (U/L)	Placebo	6.5 (2.6)	6.9 (1.8)	6.6 (1.9)	4.8 (2.2)	4.8 (3.2)
	Supplement	7.0 (2.5)	7.4 (2.4)	8.2 (3.2)	4.4 (1.7)	4.5 (2.6)
Oestradiol (pmol/L)	Placebo	170.7 (167.4)	271.4 (345.2)	801.7 (967.3)	463.9 (334.9)	217.1 (165.7)
	Supplement	105.5 (60.8)	396.5 (669.6)	711.9 (487.4)	425.1 (308.6)	269.1 (183.0)
Progesterone (nmol/L)	Placebo	3.2 (3.1)	1.0 (0.8)	4.3 (8.6)	17.1 (11.5)	11.6 (13.4)
	Supplement	2.3 (2.1)	1.3 (0.9)	1.9 (1.3)	22.5 (13.3)	11.0 (6.7)
SHBG (nmol/L)	Placebo	52.5 (28.2)	52.5 (31.7)	49.1 (23.7)	47.9 (28.9)	53.3 (31.7)
	Supplement	56.9 (35.3)	50.3 (30.1)	51.0 (28.3)	57.4* (36.6)	57.9 (33.0)

*P=0.003 paired t test supplement vs placebo (SHBG Day 21-23).

Table 5.3.2.2: Effect of isoflavone supplementation on circulating hormone levels in post-menopausal women. Mean \pm (SD).

		Baseline	Day 28
LH (U/L)	Placebo	17.3 (4.0)	18.5 (6.5)
	Supplement	16.3 (4.3)	17.7 (6.0)
FSH (U/L)	Placebo	90.9 (39.3)	106.7 (58.6)
	Supplement	86.7 (32.8)	97.6 (48.5)
Oestradiol (pmol/L)	Placebo	49.1 (43.4)	40.1 (13.4)
	Supplement	42.1 (17.6)	47.6 (25.5)
Progesterone (nmol/L)	Placebo	1.05 (0.41)	1.09 (0.62)
	Supplement	0.91 (0.73)	0.99 (0.47)
SHBG (nmol/L)	Placebo	62.8 (35.9)	61.8 (28.4)
	Supplement	61.9 (24.5)	63.0 (24.0)

Table 5.3.2.3: Comparison of IGFs and hormone levels measured by menopausal status.

	Premenopausal (n=16) Mean (SD)	Postmenopausal (n=7) Mean (SD)	Mean Difference (95%CI)	Sig. (2-tailed)
Age (y)	33.9(8.5)	57.4 (5.7)	-23.49 (-29.87, -17.11)	0.000
IGF-1 (ng/ml)	169.6 (51.4)	125.9 (43.5)	43.68 (-1.24, 88.6)	0.056
IGFBP-1 (ng/ml)	53.4 (39.9)	47.9 (16.6)	5.51 (-19.0, 30.0)	NS
IGFBP-3 (ng/ml)	3116 (395)	2874 (418)	241.7 (-168.6, 652.0)	NS
Oestradiol (pmol/l)	170.7 (167.4)	49.1 (43.4)	113.5 (20.7, 206.4)	0.019
Progesterone (nmol/l)	3.16 (3.10)	1.05 (0.41)	2.07 (0.45, 3.70)	0.016
LH (U/l)	2.5 (1.0)	17.3 (4.0)	-14.5 (-18.2, -10.8)	0.000
FSH (U/l)	6.5 (2.6)	90.9 (39.3)	-84.1 (-120.5, -47.7)	0.001
SHBG (nmol/l)	52.5(28.2)	62.8 (35.9)	-11.0 (-45.2, 23.3)	NS

Data taken from placebo baseline levels. Comparisons done with two sample t test and equal variances not assumed.

5.3.3 Insulin-like Growth Factors

Baseline concentrations of IGF-1, BP1 and BP3 fell within the expected ranges given by the manufacturers of the ELISA kits used (Diagnostic Systems Laboratories).

No significant difference was seen in IGF levels between pre and post - menopausal women, however there were slightly higher levels of IGF-1 in pre-menopausal women at baseline ($P=0.056$) (**Table 5.3.2.3**).

For pre-menopausal subjects, while there was an effect of time on IGF-1 ($p=0.005$), IGF-BP1 ($p=0.004$), and IGF-BP3 ($p<0.001$) confirming that IGF profile is influenced by menstrual cycle, this did not differ between placebo and isoflavone supplement (**Table 5.3.3.1**) (**Figures 5.3.3.1-5.3.3.3**).

When the change in IGF-1 over the whole supplementation period was compared between the supplement and placebo phases, there was a non-significant reduction in change in IGF-1 ($p=0.06$) on isoflavone supplement compared to placebo. However, this may be due to the non-significantly higher baseline concentrations of IGF-1 during the supplement phase.

In post-menopausal subjects, there was no effect of isoflavone supplementation in comparison with placebo on IGF-1, IGFBP-1 or IGFBP-3 (**Table 5.3.3.2**).

Table 5.3.3.1: Effect of supplementation on insulin-like growth factor and binding protein concentrations in pre-menopausal women

		Baseline (Day 1-3)	Day 6-8	Day 12-15	Day 21-23	Day 26-28
IGF-1 (ng/ml)	Placebo	169.6 (51.4)	170.3 (55.5)	159.7 (64.4)	192.1 (50.2) ^{ac}	194.5 (49.5) ^a
	Supplement	193.7 (71.1)	175.3 (55.0)	177.5 (57.1)	204.2 (46.0) ^{c*}	197.4 (52.1)
IGFBP-1 (ng/ml)	Placebo	53.4 (39.9)	49.9 (38.2)	43.3 (28.8)	46.6 (32.8)	64.4 (50.0)
	Supplement	59.1 (44.1)	54.5 (36.1)	63.5 (57.5)	54.5 (48.3)	72.9 (52.5)
IGFBP-3 (ng/ml)	Placebo	3116 (395)	3195 (425)	3237 (404)	3319 (446) ^a	3411 (419) ^{a*}
	Supplement	3275 (629)	3254 (478)	3305 (417)	3343 (446)	3497 (484) ^b

Data presented as mean (SD). Repeated measures ANOVA revealed significant changes over the menstrual cycle during placebo phase in IGF-1 ($p=0.005$), IGF-BP1 ($p=0.004$) and IGF-BP3 ($p<0.001$) but no significant differences between treatment regimen in these changes (IGF-1 $p=0.44$; IGF-BP1 $p=0.48$; IGF-BP3 $p=0.81$).

^a(paired *t* test significantly different from placebo baseline)

^b(paired *t* test significantly different from supplement baseline)

^c(paired *t* test significantly different from follicular phase Day 6-8)

* ($p\leq 0.01$)

Table 5.3.3.2: Effect of supplementation on insulin-like growth factor and binding protein concentrations in post-menopausal women

		Baseline Mean (SD)	28 d Mean (SD)	Mean change from baseline (SD)	Difference in mean change from baseline (95% CI)	<i>P</i> value
IGF-1 (ng/ml)	Placebo	125.9 (43.5)	137.3 (40.5)	11.37 (27.17)	0.49 (-36.93, 37.90)	0.98
	Supplement	130.2 (34.9)	141.1 (40.3)	10.89 (28.93)		
IGFBP-1 (ng/ml)	Placebo	47.9 (16.6)	48.7 (18.0)	0.76 (22.60)	-9.61 (-33.68, 14.46)	0.37
	Supplement	44.3 (9.0)	54.7 (10.7) ¹	10.37 (8.92)		
IGFBP-3 (ng/ml)	Placebo	2874 (418)	3005 (390)	92.7 (223.3)	-37.7 (-263.8, 188.3)	0.70
	Supplement	2878 (457)	2971 (516)	130.4 (225.6)		

^bMean change from baseline calculated as (28 days- baseline). Difference in mean change from baseline calculated as (placebo-supplement), and compared using a paired t-test.




Figure 5.3.3.1: IGF-1 levels in pre-menopausal women on placebo or supplement. Mean (SE)




Figure 5.3.3.2: IGFBP-3 levels in pre-menopausal women on placebo or supplement. Mean(SE)




Figure 5.3.3.3: IGFBP-1 levels in pre-menopausal women on placebo or supplement. Mean (SE)

5.3.4 Lipids and antioxidants

Total cholesterol and triacylglycerol were not affected by the supplement in either pre- or post-menopausal women. However, HDL-cholesterol concentrations were significantly elevated by isoflavone supplementation in the post-menopausal subjects. (**Figure 5.3.4.1**). There were no significant differences in any of the antioxidant endpoints assessed in either pre- or post-menopausal women when compared using paired t test (**Table 5.3.4.1**) and (**Table 5.3.4.2**).




Figure 5.3.4.1: HDL-cholesterol (mmol/L) in pre and post-menopausal women after supplementation. Mean(SE).

Table 5.3.4.1: Effect of isoflavone supplementation on antioxidant and lipid status in pre-menopausal subjects

		Baseline Mean (SD)	28 d Mean (SD)	Mean change from baseline (SD)	Difference in mean change from baseline (95% CI)	P value
Vitamin C ($\mu\text{mol/l}$)	Placebo	93.6 (47.1)	68.7 (34.1)	-24.96 (54.24)	-26.01 (-61.43, 9.40)	0.14
	Supplement	69.3 (30.0)	70.4 (41.7)	1.05 (31.90)		
MDA ($\mu\text{mol/l}$)	Placebo	2.30 (1.08)	2.26 (0.69)	-0.04 (1.03)	-0.14 (-0.67, 0.39)	0.59
	Supplement	2.22 (0.98)	2.32 (0.74)	0.10 (0.85)		
Total cholesterol (mmol/l)	Placebo	4.65 (1.10)	4.56 (0.76)	-0.09 (0.57)	-0.09 (-0.58, 0.41)	0.71
	Supplement	4.75 (0.72)	4.75 (0.93)	0.001 (0.47)		
HDL-cholesterol (mmol/l)	Placebo	1.63 (0.58)	1.61 (0.41)	-0.03 (0.26)	-0.12 (-0.28, 0.04)	0.13
	Supplement	1.57 (0.52)	1.66 (0.46)	0.10 (0.18)		
Triacylglycerol (mmol/l)	Placebo	0.73 (0.93)	0.63 (0.31)	-0.10 (0.69)	0.01 (-0.34, 0.36)	0.96
	Supplement	0.72 (0.49)	0.62 (0.30)	-0.11 (0.30)		
LDL- cholesterol (mmol/l)	Placebo	2.68 (0.89)	2.67 (0.67)	-0.02 (0.41)	0.03 (-0.38, 0.44)	0.87
	Supplement	2.86 (0.69)	2.81 (0.92)	-0.05 (0.44)		
α -tocopherol ($\mu\text{mol/l}$)	Placebo	21.0 (7.0)	18.9 (4.1)	-2.06 (8.44)	-2.78 (-8.05, 2.49)	0.28
	Supplement	18.4 (3.3)	19.2 (3.2)	0.72 (4.35)		
γ -tocopherol ($\mu\text{mol/l}$)	Placebo	4.4 (1.4)	3.8 (0.8)	-0.56 (1.79)	-0.32 (-1.42, 0.79)	0.55
	Supplement	4.2 (1.1)	3.9 (0.6)	-0.25 (0.74)		
Retinol ($\mu\text{mol/l}$)	Placebo	2.4 (0.7)	2.1 (0.6)	-0.31 (0.74)	-0.46 (-1.06, 0.13)	0.12
	Supplement	2.2 (0.7)	2.3 (0.5)	0.16 (0.60)		
lutein	Placebo	0.32 (0.15)	0.28 (0.12)	-0.04 (0.14)	-0.05 (-0.17, 0.07)	0.38

($\mu\text{mol/l}$)	Supplement	0.28 (0.12)	0.29 (0.12)	0.01 (0.11)		
zeaxanthin ($\mu\text{mol/l}$)	Placebo	0.089 (0.046)	0.082 (0.035)	-0.007 (0.041)	-0.087 (-0.044, 0.027)	0.61
	Supplement	0.076 (0.036)	0.078 (0.029)	0.002 (0.036)		
β -cryptoxanthin ($\mu\text{mol/l}$)	Placebo	0.22 (0.11)	0.20 (0.13)	-0.02 (0.10)	-0.03 (-0.15, 0.09)	0.62
	Supplement	0.22 (0.15)	0.23 (0.14)	0.01 (0.15)		
α -carotene ($\mu\text{mol/l}$)	Placebo	0.083 (0.062)	0.074 (0.070)	-0.012 (0.043)	-0.022 (-0.070, 0.027)	0.36
	Supplement	0.070 (0.049)	0.083 (0.063)	0.009 (0.058)		
β -carotene ($\mu\text{mol/l}$)	Placebo	0.82 (0.60)	0.64 (0.44)	-0.18 (0.45)	-0.32 (-0.85, 0.21)	0.22
	Supplement	0.60 (0.60)	0.79 (0.61)	0.15 (0.60)		
lycopene ($\mu\text{mol/l}$)	Placebo	0.43 (0.40)	0.36 (0.39)	-0.06 (0.55)	-0.17 (-0.67, 0.33)	0.48
	Supplement	0.39 (0.32)	0.49 (0.44)	0.11 (0.46)		

Mean change from baseline calculated as (28 d – baseline). Difference in mean change from baseline calculated as (placebo – supplement), and compared using a paired t-test.

Table 5.3.4.2: Effect of isoflavone supplementation on antioxidant and lipid status in post-menopausal subjects

		Baseline Mean (SD)	28 d Mean (SD)	Mean change from baseline (SD)	Difference in mean change from baseline (95% CI)	P value
Vitamin C ($\mu\text{mol/l}$)	Placebo	37.9 (8.9)	34.9 (14.3)	1.13 (16.41)	12.30 (-12.18,36.78)	0.25
	Supplement	46.8 (16.2)	33.9 (18.0)	-11.17 (16.77)		
MDA ($\mu\text{mol/l}$)	Placebo	2.47 (1.90)	1.98 (0.99)	-0.49 (1.55)	-0.10 (-1.64, 1.43)	0.87
	Supplement	2.35 (0.51)	1.96 (0.47) ¹	-0.39 (0.28)		
Total cholesterol (mmol/l)	Placebo	6.12 (0.96)	6.04 (1.10)	-0.08 (0.51)	-0.52 (-1.38, 0.34)	0.19
	Supplement	5.57 (1.03)	6.01 (1.14)	0.44 (1.04)		
HDL-cholesterol (mmol/l)	Placebo	1.58 (0.62)	1.54 (0.61)	-0.04 (0.10)	-0.22 (-0.40, -0.04)	0.02
	Supplement	1.54 (0.60)	1.72 (0.64) ²	0.18 (0.22)		
Triacylglycerol (mmol/l)	Placebo	1.42 (1.35)	1.33 (1.28)	-0.09 (0.17)	-0.12 (-0.58, 0.35)	0.56
	Supplement	1.15 (0.95)	1.18 (0.87)	0.03 (0.39)		
LDL- cholesterol (mmol/l)	Placebo	3.90 (1.08)	3.90 (1.20)	-0.00 (0.42)	-0.25 (-1.05, 0.55)	0.48
	Supplement	3.52(1.12)	3.76 (1.04)	0.25 (0.94)		
α -tocopherol ($\mu\text{mol/l}$)	Placebo	28.0 (6.1)	27.4 (5.2)	-0.57 (3.48)	0.37 (-0.48, 1.21)	0.33
	Supplement	27.7 (5.0)	26.8 (4.8)	-0.94 (3.95)		
γ -tocopherol ($\mu\text{mol/l}$)	Placebo	4.39 (0.60)	4.18 (0.54)	-0.21 (0.31)	-0.03 (-0.19, 0.13)	0.69
	Supplement	4.36 (0.45)	4.18 (0.41)	0.25 (0.095)		
Retinol ($\mu\text{mol/l}$)	Placebo	2.89 (0.50)	2.74 (0.33)	-0.15 (0.45)	-0.20 (-0.66, 0.26)	0.33
	Supplement	2.79 (0.48)	2.83 (0.48)	0.04 (0.31)		
lutein	Placebo	0.27 (0.09)	0.26 (0.08)	-0.01(0.07)	-0.043 (-0.14, 0.06)	0.34

($\mu\text{mol/l}$)	Supplement	0.29 (0.10)	0.32 (0.18)	0.03 (0.145)		
zeaxanthin	Placebo	0.08 (0.030)	0.06 (0.016)	-0.01 (0.02)	0.01 (-0.00, 0.02)	0.20
($\mu\text{mol/l}$)	Supplement	.069 (0.033)	0.05 (0.02)	-0.02 (0.02)		
β -cryptoxanthin	Placebo	0.40 (0.29)	0.40 (0.25)	0.01 (0.08)	-0.02 (-0.10, 0.05)	0.49
($\mu\text{mol/l}$)	Supplement	0.29 (0.12)	0.32 (0.15)	0.03 (0.06)		
α -carotene	Placebo	0.10 (0.06)	0.12 (0.08)	0.02 (0.02)	0.01 (-0.02, 0.04)	0.42
($\mu\text{mol/l}$)	Supplement	0.09 (0.07)	0.10 (0.07)	0.01 (0.02)		
β -carotene	Placebo	0.88 (0.56)	1.03 (0.71)	0.16 (0.19)	0.029 (-0.15, 0.21)	0.71
($\mu\text{mol/l}$)	Supplement	0.88 (0.55)	1.01 (0.67)	0.13 (0.17)		
lycopene	Placebo	0.39 (0.31)	0.39 (0.26)	0.00 (0.10)	-0.03 (-0.11, 0.06)	0.48
($\mu\text{mol/l}$)	Supplement	0.31 (0.27)	0.34 (0.23)	0.03 (0.10)		

Mean change from baseline calculated as (28 d – baseline). Difference in mean change from baseline calculated as (placebo – supplement), and compared using a paired t-test.

5.3.5 Nutrients

When nutrient intake was assessed by food frequency questionnaire (FFQ) or 7-day food diary (FD), there was no significant difference in energy, carbohydrate, fat, protein (in grams), SFA, PUFA, MUFA, cholesterol, fibre (Southgate method), vitamin E or carotene intake between placebo or supplement phase, whether assessed by FFQ or food diary in pre- or post-menopausal subjects (**Table 5.3.5.1**). For protein (in g) and vitamin C intake there was no significant difference between phases when assessed by FFQ. However, there was a decrease in the percentage of daily calories derived from protein in questionnaires from the supplement phase compared with the placebo phase in pre-menopausal subjects (15.0 (2.0) vs. 13.8 (1.7) placebo versus supplement, mean (SD) $p < 0.05$), while vitamin C intake was lower during the supplement phase than the placebo phase in post-menopausal women when assessed by FD, (105 (41) vs. 74 (30) mg; placebo versus supplement, mean (SD) $p < 0.05$). When compared with findings from the National Diet and Nutrition Survey 2000-1, percentage contribution of macronutrients to energy intake were similar to those of the national average for women (%total carbohydrate 48.5, %protein 16.6, %total fat 34.9)(Swan, 2004). Although the food diaries and FFQs showed good levels of association between assessed nutrient intake, lower levels were seen in FFQ for some micronutrients and estimated intakes of starch, energy and carbohydrate in pre-menopausal women which would suggest a tendency for under reporting. However, the FFQs appeared to give a good indication of long-term dietary habits as caloric intake was positively correlated with height (supplement, $r = 0.48$, $P = 0.029$), cholesterol intake with hip, weight measurements and BMI (supplement $r = 0.48$, $P = 0.032$) (supplement $r = 0.51$, $P = 0.018$) (supplement,

$r=0.456$, $P=0.038$), MUFA with hip measurements (supplement, $r=0.0454$, $P=0.045$) and percentage of fat negatively correlated with waist/hip ratio (placebo, $r=-0.52$, $P=0.0189$) (supplement, $r=-0.456$, $P=0.043$). Intake of fibre as estimated by the Englyst method was correlated with w/hip ratio ($r=0.45$, $P=0.047$) showing that the greater the fibre intake the smaller the hips. Intake of protein correlated with waist and hip measurements in both phases; waist (placebo $r=0.460$, $P=0.041$), hip (placebo $r=0.489$, $P=0.029$), waist (supplement $r=0.477$, $P=0.034$), hip (supplement $r=0.453$, $P=0.045$).

With FDs only height correlated with caloric intake ($r=0.551$, $P=0.014$) cholesterol ($r=0.518$, $P=0.023$), fat ($r=0.526$, $P=0.021$) MUFA ($r=0.539$, $P=0.017$) and starch ($r=0.570$, $P=0.011$) on the supplement phase.

Table 5.3.5.1: Dietary intake, on placebo and supplement in pre and post-menopausal women. Mean (SD)

Nutrients	Unit / day		Pre-menopausal		Post-menopausal	
			FFQ	7-FD	FFQ	7-FD
Water	gm	Placebo	2266 (848)	1907 (469) ^c	2692 (657)	2093 (600)
		Supp	3240 (3361)	1729 (567)	2601 (1090)	2186 (641)
Protein	gm	Placebo	67 (17)	77 (19)	76 (20)	70 (21)
		Supp	66 (13)	69 (19)	74 (18)	67 (22)
Fat	gm	Placebo	74 (23)	96 (28)	83 (17)	78 (23)
		Supp	83 (21)	83 (22)	84 (23)	76 (33)
Carbohydrate	gm	Placebo	204 (61)	244 (57) ^d	264 (61)	239 (61)
		Supp	211 (64)	218 (50)	264 (69)	233 (61)
Energy (kcal)	Cal	Placebo	1791 (458)	2209 (485) ^d	2105 (417)	1940 (479)
		Supp	2855 (3512)	1929 (431) ^c	2086 (372)	1898 (603)
Energy (kJ)	kJ	Placebo	7514 (1923)	9268 (2025) ^d	8835 (1758)	8173 (2018)
		Supp	11970 (14737)	8098 (1796) ^c	8758 (1560)	8005 (2537)
Total Nitrogen	gm	Placebo	11 (3)	13 (3)	12 (3)	11 (3)
		Supp	17 (23)	11 (3) ^c	12 (3)	11 (4)
Saturated fatty acids	gm	Placebo	25 (9)	34 (12)	30 (7)	29 (10)
		Supp	28 (8)	32 (12)	29 (9)	27 (14)
Mono-unsaturated fatty acids	gm	Placebo	26 (8)	32 (8)	30 (7)	25 (7)
		Supp	28 (7)	28 (8)	29 (9)	25 (12)
Poly-unsaturated fatty acids	gm	Placebo	15 (5)	18 (6)	16 (6)	15 (4)
		Supp	17 (9)	15 (5)	18 (7)	14 (5)
Cholesterol	mgm	Placebo	198 (59)	279 (105)	200 (64)	221 (87)
		Supp	206 (38)	271 (122)	200 (42)	197 (106)
Starch	gm	Placebo	97 (31)	135 (28) ^d	123 (40)	124 (37)
		Supp	98 (28)	125 (29) ^d	124 (36)	121 (45)
Total Sugars	gm	Placebo	103 (52)	103 (37)	139 (34)	112 (42)
		Supp	109 (43)	90 (46) ^c	139 (56)	107 (30)
Dietary fibre, Southgate method	gm	Placebo	21 (7)	19 (4)	29 (8)	21 (5)
		Supp	21 (7)	17 (4)	30 (9)	20 (7)
Dietary fibre, Englyst method	gm	Placebo	16 (6)	14 (3)	24 (7)	16 (4) ^d
		Supp	16 (5)	12 (3) ^b	24 (9)	15 (6)
Sodium (Na)	mgm	Placebo	2166 (722)	3225 (637) ^d	2450 (596)	2821 (742)
		Supp	2261 (611)	2821 (840) ^d	2672 (541)	2732 (976)
Potassium (K)	mgm	Placebo	3340 (1369)	3117 (543)	4045 (879)	3100 (609)
		Supp	4660 (5481)	2748 (501) ^c	3899 (1080)	3081 (895)
Calcium (Ca)	mgm	Placebo	816 (288)	992 (322)	945 (275)	912 (344)
		Supp	929 (255) ^a	861 (300)	896 (328)	874 (358)
Magnesium (Mg)	mgm	Placebo	286 (101)	297 (53)	371 (98)	294 (64)
		Supp	278 (66)	255 (48) ^b	362 (125)	284 (87)
Phosphorus (P)	mgm	Placebo	1168 (294)	1391 (291)	1377 (276)	1278 (341)
		Supp	1203 (224)	1186 (288) ^b	1404 (376)	1241 (445)
Iron (Fe)	mgm	Placebo	10 (3)	13 (3) ^d	12 (3)	12 (3)
		Supp	10 (2)	11 (3) ^b	12 (3)	13 (4)
Copper (Cu)	mgm	Placebo	1 (0.4)	1 (0.3) ^d	1 (0.3)	1 (0.3)
		Supp	2 (2)	1 (0.3)	1 (0.5)	1 (0.4)
Zinc (Zn)	mgm	Placebo	7 (2)	9 (2) ^d	9 (2)	8 (2)
		Supp	10 (12)	8 (2) ^c	9 (2)	8 (3)
Chloride (Cl)	mgm	Placebo	3449 (1117)	4917 (1084) ^d	3946 (943)	4371 (1116)

		Supp	5608 (8026)	4409 (1375) ^{c*}	4191 (826)	4181 (1455)
Manganese (Mn)	mgm	Placebo	4 (2)	3 (1)	6 (1)	4 (1)
		Supp	5 (4)	3 (1)	6(2)	4 (1)
Selenium (Se)	ugm	Placebo	45 (21)	54 (15)	63 (21)	58 (18)
		Supp	66 (98)	50 (15)	68 (18)	58 (31)
Iodine (I)	ugm	Placebo	100 (40)	110 (39)	96 (21)	111 (62)
		Supp	154 (174)	96 (31)	96 (24)	128 (66)
Retinol	ugm	Placebo	334 (213)	668 (731) ^c	531 (245)	562 (347)
		Supp	389 (182)	444 (248)	540 (216)	459 (287) ^c
Carotene	ugm					2157 (1037) ^d
		Placebo	4038 (2282)	2520 (835) ^{cd}	3924 (1494)	
		Supp	4314 (2429)	2749 (1212) ^d	3586 (1007)	2150 (1746)
Vitamin D	ugm	Placebo	2 (1)	2 (2)	2 (1)	3 (1)
		Supp	3 (4)	2 (1)	3 (1)	3 (1)
Vitamin E	mgm	Placebo	9 (3)	9 (3)	11 (6)	9 (3)
		Supp	12 (10)	7 (2)	12 (6)	9 (2)
Thiamin	mgm	Placebo	1 (0.4)	2 (3)	2 (0.5)	2 (0.4)
		Supp	1 (0.3)	1 (0.4)	2 (1)	3 (3)
Riboflavin	mgm	Placebo	2 (0.5)	2 (1)	2 (1)	2 (1)
		Supp	2 (0.4)	2 (0.4) ^b	2 (1)	2 (1)
Niacin	mgm	Placebo	21 (6)	19 (4)	25 (7)	20 (5)
		Supp	19 (5) ^a	16 (5) ^{bc*d}	27 (11)	18 (6)
Tryptophan divided by 60	mgm	Placebo	14 (4)	16 (4)	15 (4)	15 (4)
		Supp	14 (3)	15 (4)	14 (4)	14 (5)
Vitamin B6	mgm	Placebo	2 (1)	2 (0.5)	2 (1)	2 (1)
		Supp	2 (0.4)	2 (0.4) ^d	2 (0.4)	2 (1)
Vitamin B12	ugm	Placebo	2 (1)	3 (1) ^d	2 (1)	3 (2)
		Supp	2 (1)	3 (1) ^d	2 (1)	4 (4)
Folate	ugm	Placebo	354 (129)	268 (84) ^d	393 (123)	266 (83)
		Supp	332 (73)	230 (68) ^d	408 (154)	276 (111)
Pantothenate	mgm	Placebo	4 (1)	4 (1)	6 (1)	4.4 (1.1)
		Supp	4 (1)	4 (1)	6 (1)	3.9(1.4) ^b
Biotin	ugm	Placebo	35 (11)	35 (9) ^c	46 (15)	37 (11)
		Supp	35 (7)	31 (9)	46 (19)	32 (8)
Vitamin C	mgm	Placebo	192 (123)	94 (36) ^{c*d*}	231 (85)	105 (41) ^{d*}
		Supp	168 (69)	93 (53) ^d	232 (99)	74 (30) ^{bd*}
% Calories from						
Protein		Placebo	15 (2.0)	14 (2)	14 (2)	14 (2) ^c
		Supp	13.8 (1.7) ^a	14 (2)	14 (3)	14 (2)
Carbohydrate		Placebo	43 (6)	42 (7) ^c	47 (5)	47 (8)
		Supp	41 (6)	43 (6)	47 (8)	47 (5) ^c
Fat		Placebo	37 (6)	39 (5)	36 (4)	36 (4)
		Supp	3 (4)	38 (5)	36 (7)	35 (5) ^{c*}
Alcohol		Placebo	5 (6)	5 (5) ^{c*}	3 (3)	3 (4) ^{c*}
		Supp	7 (8)	5 (4) ^{c*}	2 (2)	4 (4) ^{c*}

a : mean values of FFQ on supplement significantly different from FFQ on placebo (P<0.05 paired t test)

b : mean values of FD on supplement significantly different from FDs on placebo (P<0.05 paired t test)

c : mean values of FFQ significantly correlated with FD on placebo and supplement.
(P<0.05 Pearsons correlation)

d: mean values of FFQ significantly different from FD (P<0.05 paired t test)

*: P<0.01

5.3.6 Associations between continuous variables

This study was a randomised, placebo-controlled intervention study, designed to determine the effect of isoflavone supplementation on a number of endpoints. A comparison of changes in the endpoints of interest between intervention and control groups therefore represents the main analysis of this study. However, secondary analysis, which due to the small numbers, multiple comparisons, and study design must be considered only as hypothesis-generating, has been carried out to assess cross-sectional associations between variables. These are presented as Tables in Appendix, and results that are consistent across more than one timepoint or method of assessment are presented here.

5.3.6.1 Associations with isoflavone status

When Isoflavone excretion in µg/day was compared with serum lipid/antioxidant levels, dietary intake and anthropometric measurements (**Tables 5.A.1 – 5.A.8**), the following associations were found:

Baseline levels of **equol** excretion correlated positively with energy intake at baseline, assessed by both FFQ ($p=0.019$) and FD ($p=0.022$). This may have largely been due to alcohol intake, as there was also a strong positive correlation with alcohol consumption and equol excretion at baseline on both placebo and supplement of the FFQ ($p<0.001$ and $p<0.001$ respectively) and FD ($p=0.037$ and $p=0.001$ respectively).

Excretion of **genistein** (ug/day) at baseline correlated with serum levels of a number of antioxidants including retinol ($p=0.025$), α -carotene ($p=0.046$), lutein ($p=0.011$) and β -cryptoxanthin ($p=0.034$). Genistein at baseline also correlated with estimated intakes of carotene on placebo FFQ ($p=0.020$) (**Table 5.A.4**), and FD vitamin C ($p=0.023$), suggesting overall an association between genistein status and antioxidant intake and status. However, as presented

earlier, there was no direct effect of isoflavone supplementation, leading to increased genistein excretion, on antioxidant status.

When isoflavone excretion in µg/day in pre-menopausal women was compared with serum levels of IGF-1, BP3 and BP1, there were strong positive associations between enterolactone and BP-1 throughout the cycle on both placebo and supplement (**Table 5.A.7**).

In post-menopausal women there was a strong positive correlation between levels of daidzein post supplement and levels of IGF-1 post-supplement ($p=0.004$), although this was not seen at other timepoints. Enterolactone at baseline was positively associated with IGF-1 both on placebo ($p=0.029$) and supplement ($p=0.034$) (**Table 5.A.8**). These observations suggest some association between status of the lignan enterolactone and the IGF system.

5.3.6.2 Associations with hormonal status

When hormone concentrations were compared with isoflavone excretion in µg/day (Tables 4.A.9-4.A.10) the following associations were found:

In pre-menopausal women (**Table 5.A.10**) baseline levels of **equol** were negatively associated with levels of SHBG throughout the cycle on placebo and on days 1-3, 21-23 and 26-28 of the supplement phase.

Daidzein excretion at baseline, was positively correlated with concentrations of LH, particularly in the follicular ($p<0.001$) and luteal ($p<0.001$) phase of the cycle, and with FSH concentrations throughout the cycle on both placebo and supplement. Baseline excretion of **genistein** was also positively correlated with FSH and LH throughout the cycle.

In post-menopausal women (**Table 5.A.9**) baseline levels of **genistein** excretion correlated negatively with SHBG and FSH levels at baseline whilst on supplement.

Again, however, we saw no effect of supplementation with these compounds on hormonal status.

5.3.6.3 Associations with IGF-1 and BPs

The associations between IGF-1 and binding proteins and hormonal status are presented in Tables **5.A.11-5.A.13**. When pre-menopausal hormone levels were compared with IGF-1 and BP levels (**Table 5.A.11**) there were significant negative correlations between IGF-1 and oestradiol in the Luteal phase (day 21-23) both on placebo and supplement. There were also strong positive correlations between progesterone and BP-1 at baseline and mid cycle. SHBG levels were also significantly positively correlated to BP-1 levels throughout the menstrual cycle with some weaker negative correlations with IGF-1 at baseline (**Table 5.A.12**).

In post-menopausal subjects, there were some correlations with hormone levels (**Table 5.A.13**) with SHBG showing positive correlations with IGF-1 levels, which were not seen in pre-menopausal women.

Correlations between IGF-1, BP-3 and BP-1 levels and Isoflavone concentrations are found in **Table 5.A.6 - 5.A.8**. Correlations between IGF-1, BP1 and BP3 and serum lipids and serum antioxidants can be found in **Tables 5.A.14 - 5.A.17** for premenopausal subjects and **Tables 5.A.19 - 5.A.20** for post-menopausal subjects. The following results are the strongest associations seen in pre and post-menopausal women.

BP-1 was positively associated with HDL on both placebo and supplement in pre-menopausal women.

In pre-menopausal women correlations with concentrations of BP-1 and BP-3 and serum antioxidants were as follows: BP-1 was associated with levels of α -

carotene ($r=0.606$, $p=0.013$, $n=16$) and β -Carotene ($r=0.580$, $p=0.018$, $n=16$) at supplement baseline.

Correlations between concentrations of IGF-1, BP1 and BP3 and dietary intake assessed by FFQ/FD can be found in **Tables 5.A.23-5.A.27**. The following correlations were of note:

In pre-menopausal women BP1 concentrations showed positive correlation with vitamin E intake throughout when assessed by FD, while in post-menopausal women there was a positive association between BP3 and BP1 with vitamin E intake, although this was not consistent at all timepoints, and BP1 and BP3 were not associated with vitamin E status.

5.3.6.4 Associations with lipid and antioxidant status

Pearson correlation coefficients comparing serum lipid levels with the intake of fat in FDs and FQs (**Table 5.A.21**) showed some significant correlations with serum levels of LDL, cholesterol and MDA with estimated intakes of Fat, PUFA, MUFA, cholesterol and % of calories from fat assessed using the FFQ in the placebo phase, although this was not shown for dietary data from food diaries. Pearson correlations comparing serum concentrations of antioxidants with FFQ/FD intake (**Table 5.A.22**) showed strong correlations between serum levels of carotenoids and carotene intake when assessed by both FFQ and FD but especially strong correlations seen when comparing levels of α - and β -carotene with FD estimates of carotene.

5.4 Discussion

5.4.1 Isoflavones

Supplementation for 28 days with 80 mg/day red clover isoflavones significantly increased the excretion of isoflavones although levels were highly variable between individuals. As equol is a product of intestinal bacterial metabolism formed from its isoflavone precursors daidzin and daidzein (Coward et al., 1993; Setchell et al., 2002), the large range of equol's excretion after supplementation (0-27587 nmol/day) may be explained by individual differences in isoflavone metabolic pathways (Rowland et al., 2003). Feeding studies would suggest that ~30-40% of the western population excrete equol in urine when challenged with soy foods (Karr et al., 1997; Kelly et al., 1995; Rowland et al., 2000); (Lampe et al., 1998). *In vitro* and animal *in vivo* studies show equol to be more estrogenic and have greater antioxidant activity than other isoflavones (Kostelac et al., 2003; Setchell et al., 2002), with recent clinical studies finding greater benefits for equol producers, in the areas of lipid profiles, bone density, hormone profiles and IGF-1 profiles (Duncan et al., 2000; Lydeking-Olsen et al., 2004; Meyer et al., 2004; Vrieling et al., 2007; Wu et al., 2006) decreased risk of prostate cancer (Akaza et al., 2004) and inhibition of prostate cancer in animals (Lund et al., 2004).

As the proportion of metabolites excreted may be a result of differential absorption and variation in gut microflora (Atkinson et al., 2004), the influence of habitual dietary intake on excretion of equol has been a subject of considerable interest.

5.4.2 Equol metabolism and diet

The dietary habits that promote equol production still remain unclear. Animal studies have shown that the inability of some subjects to produce equol is a consequence of the lack of specific components of the gut microflora (Bowey et al., 2003). Some studies have suggested that high fibre and low fat diets are conducive to equol production. In a study which classified 30 men and women into "excreters or non- excreters" (Lampe et al., 1998), there was no difference in fat intake but rather increased calories from carbohydrate and greater amounts of plant protein and dietary fibre in the diet of excreters. Another found that equol excreters consumed significantly less fat and more carbohydrate (Rowland et al., 2000).

In theory, this type of diet may promote the growth or activity of certain strains of bacteria shown to convert Daidzein from Dihydrodaidzein (DHD) to equol *in vitro* (Decroos et al., 2005; Ueno, 2001). However, one month supplementation with 16g fibre daily in the form of wheat bran in pre-menopausal women did not increase Equol production (Lampe et al., 2001) and a recent study found that chronic soy consumption does not induce major changes to gut metabolism of isoflavones, suggesting that the bacteria and enzymes responsible for equol or O-DMA production are not inducible (Wiseman et al., 2004). Other results suggest that high fiber intake could actually decrease bioavailability of isoflavones (Tew et al., 1996). Some feeding studies done with soy milk in a metabolic unit have suggested that the consumption of soy itself can induce equol production (Lu & Anderson, 1998; Lu et al., 1996b). However, this finding was not seen in a more recent study of one month exposure to 100mg isoflavones in the form of cereal bars and yoghurts in post-menopausal women (Vedrine et al., 2006).

In this study, we did not see consistent associations between fibre intake and equol excretion, although it was not originally designed to examine this. We did find significantly higher levels of MDA, triglycerides and α - and γ -tocopherol levels in equol producers compared to non producers, and consistent associations between equol excretion and energy intake. We also found some associations between equol excretion and levels of total cholesterol ($p=0.0090$), triglycerides ($p=0.0012$), LDL ($p=0.034$), MDA ($p=0.0015$) concentrations and the fat soluble vitamins α - tocopherol ($p=0.0039$), γ -tocopherol (1.07×10^{-6}) and retinol ($p=0.034$), but this was not consistent at all timepoints and therefore these observations are probably artefactual.

Although a study comparing soy intake assessed through FFQ in 478 post-menopausal Japanese women, found no significant differences in serum lipids between soy intake groups (Somekawa et al., 2001), other studies have not compared equol excretion with serum lipid levels. However, some studies have found correlations between fat intake as assessed through dietary questionnaires and equol excretion. A study of 19 Japanese men and women consuming their traditional diet (Adlercreutz et al., 1991) reported that equol excretion correlated positively with intake of total fat ($p<0.01$), fat-fiber ratio ($p<0.05$) and meat ($p<0.05$) which was not the case with other isoflavonoids. They proposed that the individuals consuming more meat and fat had intestinal flora that metabolised daidzein into equol more readily. Two other more recent and larger studies (Hedlund et al., 2005; Ozasa et al., 2005) have also shown associations with fat intake. A study in 45 caucasian men (Hedlund et al., 2005) showed that those who had consumed ≥ 30 mg soy for at least 2 years and consumed animal meat regularly had a higher probability of producing equol, while a study of western diets by (Lampe et al., 1999) also found isoflavone and

in particular equol excretion had strong associations with high fat and processed meat intake, and found no associations between equol excretion and soy intake. Associations examined in a Japanese population (Ozasa et al., 2005) found equol concentrations were not associated with dietary intake of tofu or miso soup (like genistein and daidzein) but with foods containing fat (butter, margarine, cheese), meat (including pork, ham) and coffee.

However, equol is actually being consumed directly from processed meats that contain soy products (Fletcher, 2003; Murphy et al., 1999) and possibly indirectly through the meat and dairy of animals consuming alfalfa, clover, or soy, supplemented feed (King et al., 1998; Lampe et al., 1999; Nilsson et al., 1967)

Certainly, within the UK population, the main contributors of isoflavones in the non-soy consuming population, were found to be bread/ bread rolls, meat products and then vegetable dishes (Mulligan et al., 2006). There is limited data on Isoflavone content of meat, milk, dairy products and fish products (King et al., 1998), (Antignac et al., 2003; Clarke et al., 2004). However, recent analysis of UK foods has found isoflavones in beers (traces of daidzein, genistein and biochanin A), fish and fish products including Indian foods (Clarke et al., 2004), with a single portion of vindaloo curry containing 11mg of genistein. Certainly, we did find strong consistent positive correlations between excretion of equol and alcohol consumption.

Analysis of Australian milk samples (King et al., 1998) found that, in areas where clover represents a significant potential source of isoflavonoids in cows milk, samples contain equol concentrations of 293 ± 52 ng/ml compared with 90ng/ml in Finish milk (Adlercreutz, 1985). Although the authors concluded these amounts were not clinically significant (Cassidy et al., 1994) large

quantities of milk and dairy products could account for baseline levels of equol excreted in the non soy consuming population and certainly free equol has a far greater binding affinity for SHBG (Martin et al., 1996) and oestrogen receptors alpha and beta (Muthyala et al., 2004) than its precursor daidzein.

Enterodiol and enterolactone are metabolites of plant lignans and their excretion has been associated with higher levels of dietary fibre intake (Adlercreutz, 2002a). Adlercreutz showed differences in lignan excretion, with vegetarians excreting significantly higher amounts than omnivores (Adlercreutz et al., 1986; Adlercreutz et al., 1995b). Lignan excretion in the western diet is also related to fruit intakes of squash, melons, stone fruits and berries (Hutchins et al., 1995; Lampe et al., 1999). However, we were unable to show consistent associations between fibre intake and lignan excretion in this study.

5.4.3 Isoflavones and hormones

It has been proposed that isoflavones exert their cancer preventative effects in part as a result of hormonal actions. To investigate the influence of the isoflavone supplement on regulation of the menstrual cycle and hormonal status, serum concentrations of oestradiol, progesterone, LH, FSH and SHBG were measured in all 16 pre and 7 post-menopausal women.

Hormones in pre-menopausal women followed normal menstrual cycle patterns and serum concentrations were within expected ranges for both age groups, given within kit instruction manuals and laboratory normal ranges. Analysis of pre-menopausal hormones (paired t test) showed SHBG concentrations were higher in the luteal phase on supplement compared with the luteal phase on placebo ($p=0.003$). However using repeated measure ANOVA (for multiple time point comparisons), Isoflavone supplementation had no effect on circulating

hormone concentrations in either pre- or post-menopausal subjects. Other intervention studies that have looked at the hormonal effects of soy have had varying results (Kurzer, 2002) (**Table 1.2.4.1**). For oestradiol concentrations, some have shown increases in the follicular phase (Cassidy et al., 1994), (Duncan et al., 1999a; Kumar et al., 2002) and overall increases throughout (Petrakis et al., 1996) while others have seen decreases in the luteal phase (Lu et al., 2000a; Lu et al., 1996a; Maskarinec et al., 2002; Wu et al., 2000) and follicular phase (Lu et al., 2000a; Lu et al., 1996a; Martini et al., 1999; Nagata et al., 1998) (Nicholls et al., 2002). Only two studies have shown a significant reduction in oestradiol (Lu et al., 1996a; Wu et al., 2000). All of these mentioned studies showed a reduction in total progesterone or concentrations measured in the luteal phase but only one found this to be a significant reduction (Lu et al., 1996a). SHBG levels measured in these intervention studies also varied with only two studies showing increases in the follicular and luteal phase (Nagata et al., 1998; Wu et al., 2000). A smaller number of intervention studies have looked at the effects of soy/isoflavones on FSH and LH levels. Studies done in a metabolic unit on six pre-menopausal women (Cassidy et al., 1995; Cassidy et al., 1994) found significantly decreased FSH and LH levels in response to 45 mg isoflavones (in the form of a textured soybean product) daily, while increasing the follicular phase length of the menstrual cycle. However, in another study, 36 oz of soy milk had no effect on LH or FSH levels in 10 pre-menopausal women with no change in mean menstrual cycle length (Lu et al., 2000a), (Lu et al., 2001). Another study (Maskarinec et al., 2002) saw no significant changes in hormone levels when comparing 17 premenopausal women on 100mg soy isoflavones for a year compared to the same size control group. Studies testing three different doses of soy protein powder in 14 pre-

menopausal women (Duncan et al., 1999a; Duncan et al., 2000), found that only the low Isoflavone dose significantly decreased LH and FSH levels midcycle. A cross-sectional study by (Verkasalo et al., 2001) looked at 636 pre-menopausal women on the EPIC study who were consuming soy milk and separated them into a <142ml or >284ml vs control group. Although FSH and LH concentrations were ~25% higher in <142ml group, they found no statistically significant associations of hormones with soy milk intake in pre or post-menopausal women. When subjects were stratified into equol excreters (n=5) vs non excreters (n=9), they found that over all three different isoflavone doses, excreters had lower levels of oestrone and higher concentrations of SHBG and midluteal progesterone (Duncan et al., 2000). (Setchell et al., 2002) has hypothesised that the effects of soy may lie with an individuals ability to metabolise daidzein to the more estrogenic metabolite equol, but (Maskarinec et al., 2002) did not find a difference in oestrone or oestradiol concentrations in equol excreters after 6 and 12 months of 100mg soy isoflavone intake. Intervention with lactobacillus and birfidobacterium (Bonorden et al., 2004) was also unable to verify this hypothesis as there was no significant difference in hormone concentrations in equol excreters vs non excreters with no significant alteration in equol or hormone concentrations after 2 months probiotic intervention. In a more recent randomised intervention study with 220 pre-menopausal women (Maskarinec et al., 2004), there appeared to be no significant difference in the oestrone, oestradiol, prgesterone and SHBG levels of the intervention group consuming ~50mg isoflavones daily versus the control group in samples taken in the luteal phase (day 19) after 3, 6,12 and 24 months. The author concluded that the effects of soy on the breast, if they exist,

are mediated by mechanisms other than the lowering of circulating oestrogen levels in pre-menopausal women.

A one-month crossover intervention study using soy milk (113-202 mg/day isoflavone content) or an isoflavone-free soymilk (<4.5 mg/day isoflavones), with a 4-month washout between phases, looked at the effect on the metabolism of 17β -oestradiol (Lu et al., 2000b). Higher levels of two putative carcinogenic metabolites of 17β -oestradiol, 4- and 16α -hydroxy oestrogen, and lower amounts of anti-carcinogenic metabolites, 2-hydroxyoestrogens, have been associated with a greater risk of breast cancer. The diet rich in isoflavones increased the cycle mean daily urinary excretion of 2-hydroxyestrone, while the mean daily excretion of 16α -hydroxy-estrone did not change. The ratio of 2-hydroxyestrone to 16α -hydroxy-estrone was higher during the isoflavone-rich diet (2.6 ± 0.34) than during the isoflavone-free diet (2.0 ± 0.32 ; $p=0.01$). These results suggest that isoflavones increase the metabolism of endogenous oestrogens to the protective 2-hydroxylated oestrogens in women, and this may play a role in lowering 17β -oestradiol levels and therefore reducing breast cancer risk (Lu et al., 2000b).

In post-menopausal women most soy intervention studies have shown few differences in levels of oestradiol, LH, or FSH (**Table 1.2.4.2**) (Baird et al., 1995; Berrino et al., 2001; Brooks et al., 2004; Brzezinski, 1997; Duncan et al., 1999b; Foth & Nawroth, 2003; Murkies et al., 1995; Nicholls et al., 2002; Persky et al., 2002; Petrakis et al., 1996; Pino et al., 2000). In the post-menopausal arm of a study with three different doses of soy protein, there was a small decrease in oestrone and oestradiol concentrations with a significant increase in SHBG on the high isoflavone diet containing ~132mg/per day for 93 days (Duncan et al., 1999b). Other studies have also seen an increase in SHBG (Berrino et al.,

2001; Kaaks et al., 2003; Pino et al., 2000). A study containing large quantities of isoflavones (Brzezinski, 1997) saw an increase in SHBG and a decrease in the strength of hot flushes, while (Nicholls et al., 2002) saw a decrease in LH peaks after discontinuation of soy containing 60mg isoflavones suggesting a residual effect post supplementation. In the large EPIC –Oxford cross-sectional study (Verkasalo et al., 2001), there were no differences seen in oestradiol or SHBG levels in 456 postmenopausal women consuming daily soy milk. Another arm of the EPIC study (Norfolk) found that urinary and serum levels of daidzein and genistein were negatively correlated with plasma oestradiol (Low et al., 2005), but that when women with certain genotypes were excluded these correlations were no longer significant. The author concluded that polymorphisms in genes encoding for enzymes involved in oestradiol metabolism (i.e, ESR1 PvuII CC genotype) may affect the concentration of plasma oestradiol, suggesting diet-gene interactions.

A study looking at correlations between levels of soy intake (assessed through FFQ) in 50 pre-menopausal Japanese women found negative correlations with oestradiol levels on day 11 and 22 of the cycle and no associations with SHBG (Nagata et al., 1997). Positive correlations were found between urinary phytoestrogens and plasma SHBG in other studies (Adlercreutz et al., 1987; Adlercreutz et al., 1992). More recent cross sectional studies of post-menopausal women (Low et al., 2005; Shoff et al., 1998; Thomas et al., 1999; Verkasalo et al., 2001) assessed phytoestrogen intakes through dietary questionnaires and found correlations between intake and SHBG or oestradiol were not significant.

Certainly in this study we did not see any significant change in hormone concentration in pre or post-menopausal women when on supplement.

However, the almost significant luteal increase in SHBG has been seen in other studies (Duncan et al., 2000; Wu et al., 2000). We saw no consistent association of lignan status with SHBG, in contrast with (Adlercreutz et al., 1987), who found that intake of total fibre correlated with enterolactone, enterodiol and plasma SHBG.

Further investigations are needed to determine whether isoflavones can cause hormonal changes and whether these are length of intervention and dose dependant. However data from this and other short term feeding studies suggest that Isoflavone effects on serum hormones and the menstrual cycle are unlikely to be the primary mechanisms by which isoflavones may prevent cancer in pre-menopausal women and certainly do not appear to have estrogenic-like effects in post-menopausal women.

Longer term supplementation than that carried out in this study, with thought given to the type of supplement (whole food or extracted supplement) and the subjects selected, may be required to fully elucidate this matter. This matter is of considerable importance in terms of the possible role of isoflavones in breast cancer prevention, where two early intervention studies focusing on the effects of phytoestrogens on the breast have both found evidence of a possible oestrogenic effect (McMichael-Phillips et al., 1998; Petrakis et al., 1996). It may be that the anti-cancer effects of isoflavones/soybeans are non-hormonal (Anderson, 1999).

5.4.4 IGF-1 and binding protein concentrations

Red clover isoflavone supplementation had no effect on IGF-1, IGFBP-1 or IGFBP-3 in post-menopausal women. In pre-menopausal women, isoflavone supplementation had no effect when multiple time points over the menstrual

cycle were considered, although there seemed to be a slight protective effect of isoflavone supplementation when the analysis was carried out simply before and after the supplementation period. This is likely to be due, however, to the large difference in IGF-1 levels at baseline and cannot be attributed to the isoflavone supplement. The effect of isoflavone supplementation on IGF status would therefore appear to be relatively weak. The clinical relevance of this small change may, in fact, be negligible. Our numbers were small, and the variance observed in IGF-1 and its binding proteins was high, so the area requires further study.

5.4.5 IGF and menstrual cycle

IGF-1 appears to be influenced by stages of the menstrual cycle. We observed a significantly higher concentration of IGF-1 and IGF-BP3 in pre-menopausal subjects on placebo in the luteal phase (days 21-23) compared to the menstrual phase (days 1-3). These changes over the menstrual cycle between the luteal and follicular phase are of a small magnitude (12.8% for placebo and 16.5% for supplement) and are in agreement with other studies (Caufriez et al., 1986; Juul et al., 1997; Lavigne et al., 2004). An early study in 5 premenopausal women (Caufriez et al., 1986) showed higher levels in the luteal (+4-+9) vs follicular (-3 - - 8) phases of the menstrual cycle and (Juul et al., 1997) also showed that, in ten healthy pre-menopausal women, there were significantly higher levels of IGF-1 in the luteal (0 - +10) vs follicular (-10 - 0) phase of the cycle ($P < 0.01$). There were no cyclic changes in BP-3, while BP-1 had a pre-ovulation peak and was negatively correlated with SHBG levels. A more recent study in 37 pre-menopausal women (Lavigne et al., 2004) also found that IGF-1 increased by 7.6% in the luteal phase (days 21-23) and 3.3% in the preovulatory phase (days

12-15), compared with the follicular phase (days 5-7) ($p=0.004$), and saw no change in BP-3 levels. These small changes may explain why several other studies have not seen any significant cyclic variation in IGF-1, BP-1 or BP-3 (Blake et al., 1997; Guidoux et al., 1986; Helle et al., 1998; Thierry van Dessel et al., 1996; Wang et al., 1995).

A study in 9 pre-menopausal women (Helle et al., 1998) found IGF-1 and BP-3 levels to be highest in the follicular phase in 8 out of 9 women studied.

However, the differences between the follicular and luteal phase were minimal with the major difference in plasma IGF-1 due to lower levels in the menstrual phase. As discussed previously (Duncan et al., 1999a), a feeding study in 14 pre-menopausal women (Wangen et al., 2000) found that IGF-1 was lower in the menstrual phase compared with the mid-follicular and mid-luteal phases ($p=0.03$ and $p=0.05$ respectively) whilst on the control ISP powder.

In this study, there were some consistent associations observed between IGF and binding proteins and hormones at certain stages of the menstrual cycle, but numbers were small, and the study not originally designed to examine this. In other studies, results for this have been conflicting. Positive correlations between oestradiol and IGF-1 have been seen (Helle et al., 1998; Ho et al., 1987; Massa et al., 1993) however others have found no correlations (Juul et al., 1997; Thierry van Dessel et al., 1996). A study by (Kelly et al., 1993) found that levels of serum oestrogen in post-menopausal women receiving exogenous oestrogens was actually negatively correlated with IGF-1 levels.

Certainly the positive correlations found in this study between BP-1 and SHBG levels throughout the menstrual cycle on both placebo and supplement reflects findings that both are regulated by insulin under various physiological conditions

(Holly, 1991) and has been seen in studies of post-menopausal breast cancer patients (Lonning et al., 1995).

These results therefore indicate that controlling closely for menstrual cycle phase may be necessary to observe IGF changes of small magnitude.

5.4.6 Isoflavones and IGF

To our knowledge this is the first published randomised controlled trial to examine the effects of isolated isoflavones on IGF-1 status in humans (Campbell et al., 2004). Since conducting this study between 1998-1999, other intervention studies have also published their findings on the effects of soy supplementation on IGF status (**Table 1.3.5.1**), including one recent study that has also looked at isolated isoflavone supplementation in men.

Of the twelve studies published to date, half have seen a significant increase in serum IGF-1 concentrations while the others have seen no effect (**Table 1.3.5.1**). The discrepancy in results may be due to differences in endogenous estrogen levels between subjects studied i.e the study of males versus females, and pre or post-menopausal women or may be explained by the increased kilocalorie/ protein intake (Clemmons & Underwood, 1991) from the actual soy supplement used which may be masking the IGF lowering effects of isoflavones alone (Voskuil et al., 2005).

In animal studies, soy protein resulted in reduced IGF-1 in both normal and polycystic kidney disease affected male animals (Aukema & Housini, 2001), while Isoflavones reduced IGF-1 and inhibited growth of prostate cancer in mice (Zhou et al., 1999). It should be noted however that rodents convert most of their daidzein to equol (Setchell et al., 2002).

In our one week study using a soy, rye and linseed bar (see chapter 4) (Woodside et al., 2006), both IGF-1 and IGF-BP3 were significantly elevated after intervention.

Two studies were conducted using soy protein with a milk protein control. In men, serum IGF-1 was higher in those supplemented with soy protein than those consuming milk protein (Khalil et al., 2002). The same supplements given to post-menopausal women (Arjmandi et al., 2003) resulted in significant increases in IGF-1 on both soy protein ($p<0.0001$) and the milk protein control ($p=0.0045$). However like the men, the greatest increases in IGF-1 (97%) was seen in women on the Soy protein especially those who were not on HRT.

These two studies would suggest that protein itself does increase IGF-1 but that soy protein had an effect on IGF-1 independent of its protein content. Calcium was not involved as both supplements had similar calcium content.

By contrast, in a three month study looking at markers of bone turnover (Wangen et al., 2000), in pre-menopausal women, IGF-1 and IGF-BP3 were increased by a low isoflavone diet, while in post-menopausal women, there were trends towards decreased IGF-1 and IGF-BP3 concentrations with increasing isoflavone concentration. The authors concluded that, although soy isoflavones do affect markers of bone turnover, the changes observed were of small magnitude and were not likely to be clinically relevant. In a study looking at soy protein supplementation with or without 83 mg isoflavones (Adams et al., 2003), it was found that changes in serum IGF-1 and BP3 were similar in both groups and there appeared to be no effect of soy isoflavones on IGF-1 when compared to the protein supplement alone.

Similarly, a one year soy protein supplementation in post-menopausal women ($n=62$) (Arjmandi et al., 2005), found that both soy and dietary protein

supplements significantly increased serum IGF-1, however only those consuming soy protein had a significant increase in IGF-BP3.

In a 2 year soya intervention (~30mg isoflavones) in pre-menopausal women (Maskarinec et al., 2005b), there was also no significant change in serum IGF-1 or BP-3 throughout. They did see a positive correlation between total isoflavone excretion and IGF-1 ($p=0.004$). The author concluded that adding soya foods to the diet of pre-menopausal women does not appear to lower serum levels of IGF-1 and BP-3, if anything the greater protein intake from soy may lead to a small increase in IGF-1 serum levels.

There have recently been a few studies that have examined the effects of soy supplementation in combination with low fat/high fibre diets. A study in 17 prostate cancer patients (Spentzos et al., 2003) found that a low fat diet with the subsequent addition of a soy protein supplement (containing 114mg isoflavones), significantly increased serum IGF-1 levels from baseline ($p=0.02$), while a study by (Gann et al., 2005) found that in 153 pre-menopausal women randomised to the consumption of a low fat high fibre diet followed by soy protein with or without isoflavones vs the usual diet, there was no difference in the LFHF diet groups vs the usual diet. There were also no differences in IGF levels between those on soy with or without isoflavones. However, there was a small but significant decrease in BP-3 in both groups post soy protein supplementation regardless of isoflavone presence. In a recent study in post-menopausal women (Kaaks et al., 2003) involving five month dietary intervention with foods rich in dietary fibre/phytoestrogens (~45mg isoflavones) and a reduction in total fat and refined carbohydrate intake, there were also no significant changes in serum IGF-1 levels. However the intervention group did have increased levels of IGF-BP1, BP-2 and SHBG.

In all of the studies mentioned above it is difficult to disentangle the effects of isoflavones from soy protein (Voskuil et al., 2005) especially in those studies which have looked at soy supplementation as part of an overall healthy dietary intervention (Gann et al., 2005; Kaaks et al., 2003; Spentzos et al., 2003). Only our study (Campbell et al., 2004) and to our knowledge one subsequent study published recently (Vrieling et al., 2007), has looked at the effects of isoflavone supplementation on its own. The study by (Vrieling et al., 2007) also used a supplement derived from red clover (by the same manufacturer) and found that in the 37 men, with a family history of colorectal cancer or a personal history or adenomas, 84mg/d Isoflavone supplementation did not significantly affect serum IGF-1, BP-3 or BP-1 serum concentrations. However, they did find that the change in serum IGF-1 concentrations post supplement were negatively correlated to serum equol concentrations ($r=-0.49$, $p=0.002$). This requires further investigation in larger studies and as our numbers were small we cannot corroborate this finding in the 4 subjects who were classified as equol producers.

The results from our study and the intervention studies above are inconclusive but would point to there being little clinical effect of soy and isoflavones on serum levels of IGF-1. The results possibly show opposing effects of Isoflavones and soy protein, with IGF-1 decreased by isoflavones but increased by soy protein and possibly protein in general. Certainly feeding studies (Clemmons et al., 1985; Smith et al., 1995) have shown that a diet restricted of certain essential amino acids reduces IGF-1 levels possibly by decreasing production of GH which in turn reduces hepatic production of IGF-1 (Ketelslegers et al., 1995) and increases BP-1 (Jousse et al., 1998). The status of endogenous oestrogen concentrations may also contribute to the differences

seen with IGF-1 association between men and women, pre and post-menopausal, HRT vs none, etc.

However further larger and long-term intervention studies using isolated isoflavone supplementation should determine whether the isoflavone content of soy is beneficial to IGF status.

Since this study was conducted there have also been a number of cross-sectional studies published that have looked at links between dietary intake and IGF-1 concentrations (**Table 1.3.4.1**). Some of the studies have been reviewed (Voskuil et al., 2005) and a few of them have also examined associations with intake of Soy/phytoestrogens (Nagata et al., 2003; Probst-Hensch et al., 2003; Vrieling et al., 2004). No correlations were seen between soy product intake and IGF-1 or BP-3 levels in 261 Japanese pre-menopausal women (Nagata et al., 2003) or the 386 pre and post-menopausal Dutch women (Vrieling et al., 2004) assessed using food frequency questionnaires. However, there was a positive association found between soy intake and IGF-1 in 321 Chinese men studied (Probst-Hensch et al., 2003) and a positive association with plant lignans and BP1 concentrations in 162 post-menopausal Dutch women (Vrieling et al., 2004). It would appear then that the findings from the cross-sectional studies are in agreement with results from our study and the other intervention studies, i.e. there appears to be very little effect of soy/isoflavones on IGF-1 concentrations and that, rather than reducing levels of IGF-1, consumption of soy may in fact be increasing levels particularly in men and post-menopausal women who are not on HRT (Arjmandi et al., 2003; Khalil et al., 2002).

Other nutritional factors that have been associated with IGF-1 levels are energy and protein intake, with malnutrition and fasting states resulting in significant reductions of serum IGF-1 levels (Thissen et al., 1994). The physiological

mechanisms are thought to be linked to low levels of insulin, resulting in consistently high levels of GH, which leads to hepatic resistance and binding of GH, with decreased production of hepatic IGF-1 (Thissen et al., 1994). However, these correlations only appear to apply with extreme dietary restrictions as only a few cross-sectional studies, done in well-nourished individuals, have shown positive associations with energy intake (Heald et al., 2003) where subjects with BMI < 25 kg/m² are analysed separately (Giovannucci et al., 2003; Holmes et al., 2002). We did not see a positive association between IGF-1 and total energy intake but we did find a number of positive correlations with anthropometric data (**Table 5.A.18**) including correlations between IGF-1 and waist, weight and hip measurements. We also did not detect any correlations with protein intake and IGF-1 in pre-menopausal women. Ten of the seventeen cross-sectional studies done to date (**Table 1.3.4.1**) did not see any significant correlations between IGF-1 and protein intake. Seven saw positive associations with protein intake (Allen et al., 2002; Devine et al., 1998; Holmes et al., 2002), (Giovannucci et al., 2003; Heald et al., 2003; Larsson et al., 2005; Norat et al., 2007) and only one study (Maskarinec et al., 2005a) saw a suggestive negative correlation with IGF-1 and protein in the form of fish intake.

This study showed some associations between serum levels of triglycerides, total cholesterol and IGF-1 concentrations in pre-menopausal women, which is consistent with findings by (Probst-Hensch et al., 2003) which also found strong positive associations between IGF-1 concentrations and serum levels of cholesterol, LDL and triglycerides. Other cross-sectional studies have examined fat intake through dietary questionnaires and have found significant positive correlations with IGF-1 concentrations (Baibas et al., 2003; Gunnell et al., 2003;

Heald et al., 2003; Kaklamani et al., 1999), with one study showing a negative association between intake of fat from meat and IGF-1 (DeLellis et al., 2004).

Interestingly we found that, in pre-menopausal women, BP-1 levels were consistently positively associated with HDL levels and as there have been no other studies to examine this link it would be worth further investigation.

Studies linking the IGF system with the development of cardiovascular disease (Bayes-Genis et al., 2000), have been examined fairly recently, finding associations between insulin like growth factors and its binding proteins with serum lipid levels, (Fischer et al., 2004; Harrela et al., 2002; Janssen et al., 1998; Juul et al., 2002; Laughlin et al., 2004).

The positive associations seen in our pre-menopausal women between BP-1 and HDL, agree with findings by (Janssen et al., 1998) and (Laughlin et al., 2004), while the positive associations for IGF-1 with triglycerides and cholesterol (although not consistent at all timepoints) appear to agree with findings by (Laughlin et al., 2004) but not (Harrela et al., 2002; Juul et al., 2002).

The links are not entirely clear, however, with (Juul et al., 2002) showing that low IGF-1 levels puts patients at significantly higher risk of developing ischaemic heart disease (IHD), while recent findings by (Fischer et al., 2004) suggest a positive association with IGF-1, BP-3 levels and CVD. A review by (Juul, 2003) suggesting that the controversy around clinical studies of IGF-1 being either cardioprotective or a risk factor for IHD may reflect that IGF-1 was studied at different time points in relation to the ischaemic heart event.

Although none of the cross-sectional studies done to date have looked at associations with serum levels of antioxidants, the negative correlations we found between IGF-1 and its binding proteins and serum levels of carotenoids are similar to findings by the recent large EPIC cross-sectional study examining

diets of women in Europe (Norat et al., 2007) which saw negative correlations with dietary intake of β -carotene, while an earlier study of 112 Greek men (Mucci et al., 2001) showed that greater consumption of cooked tomatoes (the largest source of lycopene) was associated with lower concentrations of IGF-1 ($p=0.041$). However, positive correlations have also been observed in cross-sectional studies (Holmes et al., 2002; Maskarinec et al., 2005a).

There also appears to be a strong suggestion that milk and/or dairy consumption is associated with higher levels of IGF-1, with many of the cross-sectional studies showing positive correlations with IGF-1 concentrations and milk/dairy foods (Gunnell et al., 2003; Ma et al., 2001; Norat et al., 2007) and/or calcium and vitamin D intake (Giovannucci et al., 2003; Gunnell et al., 2003; Holmes et al., 2002; Nagata et al., 2003; Norat et al., 2007; Probst-Hensch et al., 2003). While consumption of $\frac{3}{4}$ pint of soya milk daily in vegan women significantly increased IGF-1 concentrations (Allen et al., 2002), these increases in IGF-1 could be attributable to increased levels of protein (including essential amino acids and possibly ingestion of bovine IGF-1 contained in milk (Giovannucci, 2003). Other nutrients that have been shown to be positively associated with IGF-1 include intake of vitamin B6 and B12 (Norat et al., 2007). (Maskarinec et al., 2005a), found that women in the top quartiles of dietary fiber intake had 5.7% higher IGF-1 levels than women in the bottom quartiles ($p=0.004$).

We did not see any associations with IGF levels and alcohol consumption unlike other studies (Probst-Hensch et al., 2003; Vrieling et al., 2004), nor did we find any associations between Zinc intake and IGF-1 (Devine et al., 1998; Giovannucci et al., 2003; Holmes et al., 2002; Larsson et al., 2005).

Due to the small numbers in our study it is difficult to draw conclusions from the correlations found between serum levels of lipids and antioxidants, dietary intake and IGF-1, BP3 and BP1 levels. Further larger studies and intervention studies should be done to determine if there is a causal effect and what the relationships are with these dietary components, including analysis of serum concentrations of these nutrients which may give a more accurate assessment of dietary intake.

5.4.7 Isoflavones and lipids

We have shown an effect of isoflavone supplementation within one month on HDL-cholesterol in post-menopausal subjects. Several studies have provided evidence that phytoestrogens in general can modulate plasma lipid and lipoprotein concentrations, but the data are not entirely consistent (Anderson et al., 1995). A recent overview concluded that soy could improve blood lipid parameters in both normocholesterolaemic and hypercholesterolaemic subjects, although the use of soy alone may not allow patients with hyperlipidaemia to achieve target lipid parameters (Costa & Summa, 2000). Soy, with which most of the studies have been carried out, has a variety of properties unrelated to isoflavone content that may contribute to its lipid-modulating properties (Potter, 1995). Our post-menopausal subjects had normal lipid levels, but displayed an increase in HDL-cholesterol during the isoflavone supplementation phase. This may be pertinent in light of the fact that, a short-term study using the same purified red clover isoflavone pills found no effect on plasma lipids and lipoprotein concentrations (Nestel et al., 1999) and another found only small changes in levels of triglycerides (Schult et al., 2004). However other longer term studies using this supplement have demonstrated an effect of isoflavones

on HDL₃ (Samman et al., 1999) in pre-menopausal women and total HDL (Clifton-Bligh et al., 2001; Knight et al., 1999) in pre-menopausal and post-menopausal women. A recent study in post-menopausal women with a different isoflavone supplement not derived from red clover (containing 47mg as aglycone form) found that HDL cholesterol significantly increased from baseline in the combination of walking and supplement but not on supplement alone (Wu et al., 2006). The apparent difference in these findings compared to those obtained with soy-derived isoflavones (Dewell et al., 2002; Gardner et al., 2001; Somekawa et al., 2001) could be accounted for by the high levels of biochanin and fomonomentin in red clover isoflavone extracts.

5.4.8 Isoflavones and antioxidants

Isoflavones can act as antioxidants (Bingham et al., 1998), both *in vitro* and test systems. (Kapiotis et al., 1997) observed that LDL oxidation products, assayed as thiobarbituric acid-reactive substances, were strongly inhibited by genistein, somewhat inhibited by daidzein, but not affected by genistin (the glycosylated form of genistein) or control. Furthermore, both genistein and daidzein were found to protect against cytotoxic effects of oxidised LDL as assessed by cellular morphologic features and lactate dehydrogenase release by cultured endothelial cells. Genistein has also been shown to inhibit hydrogen peroxide production and increase the activity of antioxidant enzymes, such as catalase, superoxide dismutase, glutathione peroxidase and glutathione reductase (Lissin & Cooke, 2000). A protective/antioxidant effect has been shown by phytoestrogen supplementation on the susceptibility of LDL to oxidation in another study (Tikkanen et al., 1998). Six healthy volunteers received 19 mg isoflavones/day for 2 weeks. Lag phases of LDL oxidation curves were reduced

post-supplement compared with baseline values. Another study, however, failed to show an inhibition of lipid peroxidation after 8 weeks of isoflavonoid supplementation in subjects with high-normal blood pressure (Hodgson et al., 1999), measuring urinary F-2-isoprostanes. Another study in 24 subjects showed that consumption of soy containing burgers containing 21mg Daidzein and 35mg genistein isoflavones, reduced lipid peroxidation in plasma in the form of plasma F(2)isoprostane and increased the resistance of LDL to oxidation (Wiseman et al., 2000). We found little effect of isoflavone supplementation on markers of antioxidant status. Our subjects were healthy and had normal antioxidant levels; it may be that isoflavones will exert marked antioxidant effects *in vivo* only in those with an already compromised antioxidant status.

5.4.9 Diet

Background diet did not seem to affect the findings of this study, with the magnitude of any differences observed being unlikely to affect our conclusions. Overall analysis of dietary intake showed that mean intakes were comparable to national averages in the UK (Swan, 2004). The percentage of calories from protein, carbohydrate, fat and alcohol were very similar between FFQs and FDs. In terms of biomarkers, overall serum levels of carotenoids appeared to correlate strongly with both FFQ and FDs. Serum lipid levels of LDL, cholesterol and MDA showed some correlation with FFQ estimated intakes of fats but not at all with intakes estimated from FDs. The FFQ estimated intake of calories, cholesterol, % fat and fibre also showed some interesting correlations with anthropometric data.

Thus, it would appear that both methods of dietary assessment showed a reasonable level of agreement. However the FFQ appeared to perform better when compared with biomarkers and anthropometric data. Therefore the FFQ may well give a better indication of long-term dietary habits.

5.5 Conclusion

Red clover-derived isoflavone supplementation over one month has no effect on IGF status in post-menopausal women, and at most has a small effect on IGF-1 in pre-menopausal women. Red clover isoflavones seem to increase HDL-cholesterol in post-menopausal women, but have little or no effect on other lipids or antioxidant status. It must be remembered, in the interpretation of these data, that the study was small and had low power to detect a difference between placebo and supplement, particularly when multiple comparisons were being carried out. Further work in larger numbers will determine whether these dietary compounds may play a role in endocrine-responsive cancer and Cardiovascular disease prevention.

CHAPTER SIX – THE EFFECT OF ISOFLAVONES ON DNA SYNTHESIS IN MCF-7 AND MDA-MB-231 HUMAN BREAST CANCER CELLS

6.1 Introduction

In vitro, isoflavones inhibit the growth of both oestrogen dependent and independent breast cancer cells by a number of different mechanisms (Dampier et al., 2001; Magee & Rowland, 2004; Peterson & Barnes, 1991; Sarkar & Li, 2003).

Isoflavones bind to mammalian oestrogen receptors with between 1×10^{-2} and 1×10^{-4} the activity of 17β -oestradiol and are conventionally considered weak oestrogenic compounds (Markiewicz et al., 1993). However the discovery of the oestrogen receptor β subtype (ER β) (Kuiper et al., 1996; Kuiper & Gustafsson, 1997) has added a new dimension to the understanding of oestrogen and anti-oestrogenic actions of these compounds. The relative binding affinities and transcription activities of isoflavones compared to oestrogen, in both α and β subtypes have been investigated (Kuiper et al., 1997; Kuiper et al., 1998). Isoflavones, and in particular genistein and equol, have 30-100 fold higher binding affinities and transcription activities for ER β compared with ER α (An et al., 2001; Barkhem et al., 1998; Kostelac et al., 2003) (McCarty, 2006).

Normal and malignant breast tissue express both receptors (Kuiper et al., 1997). Studies suggest that ER β is the principle oestrogen receptor in normal breast tissue with decreasing expression in poorly differentiated cancerous and metastatic tissue (Palmieri et al., 2002), with others finding that ER β is upregulated in breast tumours that develop antioestrogen resistance (Speirs &

Kerin, 2000). Although the precise role of ER β activation is unclear (Park et al., 2003) there is evidence that it opposes the proliferative effects of ER α *in vitro* (Caetano et al., 2006; Paruthiyil et al., 2004; Strom et al., 2004).

Expression of the oestrogen receptors in breast cancer cell lines, shows that the ER positive MCF-7 cells strongly express ER α , while both MCF-7 and the ER negative MDA-MB-231 cells express low levels of ER β (Dampier et al., 2001; Fuqua et al., 1999; Tong et al., 2002; Vladusic et al., 2000).

Genistein and daidzein bind to ER α with 4% and 0.1% affinity of oestradiol respectively (Kuiper et al., 1998). Equol has similar binding affinity to genistein but a much stronger ability to induce transcription (Morito et al., 2001). In ER α positive MCF-7 cells, isoflavone concentrations $\leq 1\mu\text{M}$ increase cell proliferation, with genistein shown to be a strong oestrogenic agonist for both ER isoforms (Maggiolini et al., 2001). However, higher concentrations ($\geq 10\mu\text{M}$) have been shown to inhibit hormone dependent growth by oestrogen antagonistic activity, and these biphasic effects on cell growth are now well documented (Fioravanti et al., 1998; Hsieh et al., 1998; Hsu et al., 1999; Le Bail et al., 1998; Power & Thompson, 2003; Wang & Kurzer, 1997; Wang et al., 1996; Zava & Duwe, 1997). Research by (Hsieh et al., 1998) found that concentrations as low as 10nM of genistein could have proliferative effects in MCF-7 cells, with 100nM achieving proliferative effects similar to 1nM of oestradiol. Others have also found that genistein and equol induced oestrogen regulated pS2 expression at concentrations of 1nM to 1 μM , with their growth stimulation linked to the binding affinity for ER α (Zava & Duwe, 1997). They also found that genistein's ability to override the growth inhibitory effects of tamoxifen helped demonstrate that the inhibitory mechanisms of isoflavones may not be the same as classical oestrogen agonists like tamoxifen. (Le Bail et al., 1998) found that, in the

presence of 0.1nM oestradiol, the proliferative effects of isoflavones at lower concentrations were no longer present while their inhibitory effects at higher concentrations (~50 μ M) were unaffected by oestradiol.

Oestradiol and IGF-1 appear to work together in the proliferation of breast cancer cells with increasing support for cross talking between the ER and IGF-1R signalling pathway (Kato et al., 2000; Laban et al., 2003). Oestradiol has been shown to stimulate expression of IGF-1, the IGF-1R, IRS-1 and down regulate expression of the IGF binding proteins (Ruan et al., 1995; Stewart et al., 1990), (Huynh et al., 1996b; Lee et al., 1999). The cytotoxic effects of tamoxifen are partially mediated through IGF-1 receptor (IGF-1R) signalling (Guvakova & Surmacz, 1997), while genistein has recently been shown to enhance the IGF-1 signalling pathway in proliferation of MCF-7 cells at low concentrations (Chen & Wong, 2004).

In ER α negative cells, isoflavones do not appear to cause biphasic effects on cell proliferation. A study comparing the effects of genistein in different breast cancer cell types, found that growth promotion at low concentrations of 1-10 μ M were only apparent in MCF-7 cells, and that different mechanisms were found to promote growth depending on breast cell type (Dampier et al., 2001). Whether these antiproliferative effects are mediated through binding to ER β has yet to be fully examined. However research by (Caetano et al., 2006) has recently found that in MDA-231 cells, genistein increases expression of all genes involved in cell cycle control, apoptosis and DNA repair including BRAC1, p53, p21, Rad51, BARD, Bax GADD45 and RF1, suggesting it may decrease the inhibition exerted by oestrogens on the transcription of these genes by antagonising ER β .

A recent study using cDNA microarray technology has shown that 50 μ M Genistein can also upregulate genes such as HSP, involved in cell salvage response, and down regulate SRF transcription factor, involved in cell growth /differentiation through oestrogen receptor independent pathways (Chen et al., 2003). Other oestrogen independent mechanisms have been reviewed (Middleton et al., 2000; Sarkar & Li, 2003), and include: inhibition of AKt signalling and activation of nuclear factor KB DNA binding activity needed for cell survival, inhibition of antiproliferative enzymes such as topoisomerase I and II and 5 α reductase, inhibition of protein tyrosine kinase mediated signalling pathway, inhibition of angiogenesis and metastasis, stimulation of immunity and antioxidant activity (Akiyama et al., 1987; Akiyama & Ogawara, 1991; Chen et al., 2003; Kim et al., 1998; Ruiz-Larrea et al., 1997; Shao et al., 1998a; Shao et al., 1998b; Sierens et al., 2001; Su et al., 2005; Zhang et al., 1999).

However, caution is warranted in the interpretation of some of these effects, as they have been observed at higher concentrations than may be found *in vivo*. Reports on plasma concentrations of isoflavones do vary, with levels reported to reach no higher than 18.5 μ M (Barnes & Peterson, 1995). Soy rich Japanese diets can result in total concentrations above 5 μ M (Morton et al., 1994) with mean concentrations of genistein in Japanese men (493nM) compared with men from UK (33nM). It was found that 58% of Japanese men and 38% of women had serum equol concentrations above 20nM compared with none and 2.2% of comparable samples in the UK population (Morton et al., 2002). Total levels of isoflavones following moderate consumption of soy are reported to be ~2 μ M (Franke et al., 1998) with studies following consumption of genistein rich foods, finding that plasma concentrations are generally <1 μ M of genistein (Barnes et al., 1996), (Adlercreutz et al., 1995a; Gooderham et al., 1996;

Wiseman et al., 2004; Xu et al., 1994). Recent bioavailability studies looking at the American caucasian female population have found peak plasma genistein concentrations of 0.5-2.2 μM after 4-8 hours of ingestion (Setchell et al., 2003), with the bioavailability from aglycone forms higher than glycosides for daidzein with no difference seen for genistein (Zubik & Meydani, 2003). Studies suggest that as little as ~0.5-2% of genistein in serum is actually available as the unbound unconjugated form of genistein (Barnes, 1995; Nagel et al., 1998) which may be equivalent to ~10nM genistein concentrations *in vitro* (McCarty, 2006).

However, cellular concentrations may in fact be higher than this, with oestrogen concentrations found to be up to 40 fold higher in breast ductal fluid than in serum (Ernster et al., 1987). A study by (Hargreaves et al., 1999) also found significantly higher isoflavone levels in nipple aspirate fluid than in serum, in both control and soy supplemented pre-menopausal women, suggesting accumulation of phytoestrogens within breast ducts. Similar concentrated effects have also been seen in prostatic fluid (Morton et al., 1997). However, (Pumford et al., 2002) found that breast tissue levels of phytoestrogens are comparable to serum levels, and increase with isoflavone supplementation.

In vitro studies in breast cancer cells can improve our understanding of the possible therapeutic benefits that isoflavone compounds may have. Thus, the present study was designed to investigate the effects of isoflavones on DNA proliferation in the oestrogen dependent, non-invasive MCF-7 breast carcinoma cell line and the oestrogen-independent, invasive MDA-MB-231 breast carcinoma cell line. Interactions between the isoflavones, antiestrogen Tamoxifen and IGF-1 were also examined to investigate possible mechanisms

of action. To examine the mechanisms by which isoflavones inhibit cell growth, concentrations of 10 μ M (~IC₅₀) of isoflavones were combined with:

- 1nM Oestradiol to imitate levels seen in plasma for premenopausal women (Pike et al., 1993),
- 10nM IGF-1 which gave near maximum promotion of DNA synthesis in both cell lines
- 1 μ M tamoxifen to imitate levels in women receiving tamoxifen for prevention/treatment of breast cancer (Lien et al., 1989) and can inhibit ER and IGF-1 mediated proliferation of MCF-7 cells (Jordan, 1994) (Wosikowski et al., 1993).

6.2 Materials and methods

6.2.1 Cell Culture

MCF-7 and MDA MB -231 cells (kindly donated by Ludwig cancer research group) were maintained in Phenol Red IMEM (GIBCO) supplemented with 5% FCS (GIBCO), 1% L-Glutamine (sigma) and 1% Fungizone (GIBCO) in 5% CO₂ at 37°C. Conditions involved growing cells in phenol red free IMEM supplemented with 5%FCS for 72 hours prior to plating. Cells were then seeded in 96 well plates (NUNC) at 1×10^4 cells per well in 5% DCC-FCS (Dextran charcoal coated foetal calf serum) phenol red free IMEM and incubated for 24 hours. Medium was changed to serum free IMEM for another 24 hours before the compounds were added. Stock solutions of tamoxifen, genistein (4,5,7-trihydroxyisoflavone), daidzein (4,7-dihydroxyisoflavone), equol and 17 β -oestradiol (Sigma) in ethanol were used to produce the required concentrations of compounds in culture medium. Control conditions also contained the appropriate amount of ethanol (0.1%).

6.2.2 BrdU assay

The principle of the assay is to determine cell proliferation by measuring 5-bromo-2-deoxyuridine (BrdU) incorporation during DNA synthesis in proliferating cells. This assay indicates growth rate as opposed to cumulative cell growth. Conventional methods use thymidine, however this technique is based on the incorporation of the pyrimidine analogue 5-bromo-2'-deoxyuridine (BrdU). The BrdU assay was performed using a colorimetric ELISA kit (Amersham-Pharmacia-Biotech, 1998). Results were from five replicate wells with experiments performed two to three times.

See Chapter two for further detail on Methods.

6.3 Results

Because we have not directly compared DNA synthesis with other assays, such as cell number or protein content, we cannot assume that these results are directly comparable. However, previous studies have compared assays and found DNA synthesis to parallel other assays and directly correlate with the [³H]thymidine incorporation assay (Amersham-Pharmacia-Biotech, 1998).

The work presented here does appear to correlate quite closely with results from studies that have used thymidine to analyse DNA synthesis in these cell types, and could therefore be considered as a useful and safer alternative.

6.3.1 Effects of E₂ and IGF-1 on DNA synthesis

DNA synthesis was significantly stimulated with E₂ after 24 hour incubations in MCF-7 cells (**Table 6.3.1.1** and **Fig 6.3.1.1**). Induction was 16%, 173%, 187% and 159% of the 0% control at 0.01nM, 0.1nM, 1nM and 10nM, respectively. (P<0.01). In MDA-MB-231 cells E₂ did not stimulate DNA synthesis but appeared to inhibit cells to between 87 and 89% of the control at higher concentrations (P<0.05) (**Table 6.3.1.2** and **Fig 6.3.1.2**).

IGF-1 significantly increased DNA synthesis in MCF-7 cells to 123%, 149%, 198% and 214% of the control at 0.1nM, 1nM, 10nM and 100nM, respectively (P<0.01) (**Table 6.3.1.1** and **Fig 6.3.1.1**). In MDA-231 cells IGF-1 did not stimulate DNA synthesis but inhibited to 92 and 89% of the control at higher concentrations (P<0.05) (**Table 6.3.1.2** and **Fig 6.3.1.2**).

6.3.2 Effects of Isoflavones and Tamoxifen on DNA synthesis

The isoflavones genistein, daidzein and equol all stimulated DNA synthesis to a maximum of 207%, 198% and 209% at 1μM respectively. Both genistein and

equol caused significant stimulation between concentration of 0.01 μ M-10 μ M, ($P < 0.01$), while daidzein significantly increased stimulation between 0.1 μ M-100 μ M, ($P < 0.01$). Biphasic effects, meant that at higher concentrations, genistein and equol were able to inhibit DNA synthesis, with IC_{50} values of 93 and 55 μ M respectively. Tamoxifen did not have biphasic effects and inhibited DNA synthesis from 0.1 μ M ($P < 0.05$) through to 100 μ M ($P < 0.01$) with an IC_{50} value of 1.17 μ M (**Table 6.3.1.1** and **Fig 6.3.2.1**).

In MDA-MB-231 cells the isoflavones did not stimulate DNA synthesis at any concentration but significantly inhibited induction as concentrations increased. Genistein inhibited DNA synthesis from concentrations of 0.1 μ M ($P < 0.01$) upwards with an IC_{50} value of 14 μ M. Daidzein and equol inhibited DNA synthesis at concentrations greater than 1 μ M ($P < 0.01$) with an IC_{50} value of 15 μ M for equol. Tamoxifen inhibited synthesis from concentrations of 0.1 μ M ($P < 0.01$) onwards with an IC_{50} value of 1.35 μ M (**Table 6.3.1.2** and **Fig 6.3.2.2**).

Table 6.3.1.1: Dose response of MCF-7 cells incubated with Isoflavones, Tamoxifen, IGF-1 and oestradiol for 24 hours.

	0.01uM	0.1uM	1.0uM	10uM	100uM	IC ₅₀
Daidzein	96.4(15.6)	153.5(34.6)*	198.0(50.5)*	161.2(60.2)*	133.8(41.2)*	>100µM
Genistein	163.8(29.7)*	197.7(25.3)*	207.1(43)*	156.3(39.2)*	41.0(27.4)*	93µM
Equol	122.4(14.6)*	193.8(43.2)*	209.1(54.2)*	141.5(45.4)*	14.7(1.9)*	54.8µM
Tamoxifen	92.5(9.4)	91.5(8.0)*	59.7(11.6)*	7.45(0.7)*	9.11(0.8)*	1.17µM
	1pM	0.01nM	0.1nM	1nM	10nM	100nM
IGF-1		102.9(14.4)	123.3(26.0)*	149.2(18.0)*	197.6(45.2)*	214.3(46.5)*
Oestradiol	97.4(17.7)	160.9(22.9)*	172.9(25.9)*	187.4(45.8)*	159.3(39.2)*	

Levels given as percentage of the control. Mean (SD) *P<0.01 Mann Whitney U test.

Table 6.3.1.2: Dose response of MDA-231 cells incubated with Isoflavones, Tamoxifen, IGF-1 and oestradiol for 24 hours.

	0.01µM	0.1µM	1.0µM	10µM	100µM	IC ₅₀
Daidzein	100.4(9.3)	104.5(11.2)	98.3(12.1)	80.4(8.1)*	54.0(7.9)*	113µM
Genistein	96.4(10.3)	89.5(5.7)*	85.6(7.8)*	59.7(10.5)*	19.5(5.4)*	14.1µM
Equol	94.0(7.0)	100.8(7.7)	97.4(14.9)	65.2(11.0)*	9.0(1.7)*	14.55µM
Tamoxifen	96.1(5.6)	86.9(6.8)*	74.5(12.4)*	4.7(0.3)*	5.1(0.5)*	1.35µM
	1pM	0.01nM	0.1nM	1nM	10nM	100nM
IGF-1		101.3(6.2)	106.7(12.3)	101.1(10.5)	91.7(9.2)*	89.3(5.8)*
Oestradiol	97.4(13.0)	91.5(13.5)*	88.7(8.4)*	92.6(9.4)*	86.7(15.0)*	

Levels given as percentage of the control. Mean (SD) * P<0.01 Mann Whitney U test.




Figure 6.3.1.1: Dose response of MCF-7 cells incubated with IGF-1 and oestradiol for 24 hours before labelling with BrDU. Mean (SE)




Figure 6.3.1.2: Dose response of MDA-MB-231 cells incubated with IGF-1 and oestradiol for 24 hours before labelling with BrDU. Mean (SE)




Figure 6.3.2.1: Dose response of MCF-7 cells incubated with isoflavones and Tamoxifen for 24 hours before labelling with BrDU. Mean (SE)




Figure 6.3.2.2: Dose response of MDA-MB-231 cells incubated with isoflavones and Tamoxifen for 24 hours before labelling with BrDU. Mean (SE)

6.3.3 Interactions between isoflavones and Tamoxifen with Oestradiol

In MCF-7 cells maximal stimulation of DNA synthesis with 1nM E₂ alone was 187% of the control. When incubated in combination with 10μM genistein, daidzein and equol, synthesis induced by 1nM E₂ was decreased by 51%, 41% and 43% respectively (P<0.01).

When incubated in combination with 1μM tamoxifen, the decrease of E₂ induced synthesis was 28% (P<0.01) (**Table 6.3.3.1**).

In MDA-231 cells the inhibition caused by isoflavones alone was counteracted by the addition of 1nM Estradiol such that the inhibition of DNA synthesis was reduced by 10%, 7% and 11% for genistein, daidzein and equol respectively (p<0.01).

This was not the case with tamoxifen as there was very little difference seen between inhibition with or without E₂ (**Table 6.3.3.2**).

6.3.4 Interactions between Isoflavones with Tamoxifen

In MCF-7 cells, 1μM tamoxifen inhibited DNA synthesis by 60% (P<0.01).

However, when added to 10μM genistein and daidzein, tamoxifen increased DNA synthesis by a further 22% and 15% (P<0.05). This increase was not seen when tamoxifen was added to equol (**Table 6.3.3.1**).

In MDA-231 cells, 1μM tamoxifen inhibited DNA synthesis to 74.5% of the control. Unlike the MCF-7 cells, the addition of genistein and equol reduced this inhibition further by 11% (P<0.01) and 22% (P<0.01) (**Table 6.3.3.2**).

6.3.5 Interactions between Isoflavones and Tamoxifen with IGF-1

Stimulation of DNA synthesis with 10nM IGF-1 was 198% of the control (P<0.01). When incubated with isoflavones, induction was decreased. With

equol, this was a significant reduction of 46.2% ($P < 0.01$). Tamoxifen also reduced IGF-1 induced synthesis but this was not significant. No significant increase in DNA synthesis was seen with the addition of E_2 to IGF-1 (**Table 6.3.3.1**).

In MDA-231 cells the addition of 10nM IGF-1 reduced the ability of genistein, daidzein and equol to inhibit DNA synthesis by 11%, 20% and 17%, respectively ($P < 0.05$). However, this was not the case with tamoxifen as there was no difference seen in the amount of DNA inhibition between tamoxifen on its own or combined with IGF-1. There was a small yet significant increase in DNA synthesis with the addition of E_2 to IGF-1 ($P < 0.01$) (**Table 6.3.3.2**).

Table 6.3.3.1 Response of MCF-7 cells incubated for 24 hours with different combinations of Isoflavones, oestradiol, Tamoxifen and IGF-1.

		Oestradiol(1nM)	Tamoxifen(1μM)	IGF-1(10nM)
Genistein(10μM)	156.3(39.2)	136.9(7.9)	178.6(17.2)	168.7(9.2)
Daidzein(10μM)	161.2(60.2)	146.5(12.9)	176.2(17.6)	181.3(12.4)
Equol(10μM)	141.5(45.4)	144.6(13.4)	143.2(15.2)	151.4(22.3)
Tamoxifen(1μM)	59.7(11.6)	159.0(49.8)	-	169.6(41.0)
Oestradiol(1nM)	187.4(45.8)	-	159.0(49.8)	190.2(9.6)
IGF-1(10nM)	197.6(45.2)	190.2(9.6)	169.6(41.0)	-

Levels given as a percentage of the control. Mean (SD)

Table 6.3.3.2: Response of MDA231 cells incubated for 48 hours with different combinations of Isoflavones, oestradiol, Tamoxifen and IGF-1.

		Oestradiol(1nM)	Tamoxifen(1μM)	IGF-1(10nM)
Genistein(10μM)	59.7(8.1)	69.9(4.3)	63.4(6.0)	70.2(12.8)
Daidzein(10μM)	80.4(10.5)	87.0(6.8)	91.4(6.6)	100.2(18.6)
Equol(10μM)	65.2(11.0)	76.4(4.0)	52.9(5.1)	81.9(14.2)
Tamoxifen(1μM)	74.5(12.4)	68.2(12.9)	-	71.1(9.1)
Oestradiol(1nM)	92.6(9.4)	-	68.2(12.9)	102.1(11.2)
IGF-1(10nM)	91.7(9.2)	102.1(11.2)	71.1(9.1)	-

Levels given as a percentage of the control. Mean (SD)

6.4 Discussion

6.4.1 Isoflavones

DNA synthesis was significantly increased by E_2 in MCF-7 ($ER\alpha+/ER\beta+$) cells but not MDA-MB-231 ($ER\alpha-/ER\beta+$) cells. These results confirm the role of $ER\alpha$ in facilitating the dose dependent proliferation effects of E_2 in MCF-7 cells (Dampier et al., 2001). The isoflavones genistein, daidzein and its metabolite equol also significantly stimulated DNA synthesis in a dose dependant manner in MCF-7 cells. Maximal stimulation for all three isoflavones tested was reached at 1.0 μM , with an increase of 207%, 198% and 209% for genistein, daidzein and equol respectively. However, genistein and equol concentrations $>10\mu M$ inhibited DNA synthesis with IC_{50} values of $93\mu M$ and $54.8\mu M$ respectively. This biphasic response of MCF-7 cells to increasing concentrations of isoflavones has also been reported in other studies looking at DNA synthesis. Incubation for 24 hours with 1 μM genistein and 10 μM daidzein in the presence of 5% DCC-fCS, maximized DNA synthesis to 215% and 210% of the control while, inhibition occurred between 50-100 μM with an IC_{50} value for genistein of 41 μM (Liu & Wang, 1997; Wang & Kurzer, 1998). (Hoffman, 1995) found genistein IC_{50} values of 12.6 μM in the presence of no growth factors and an IC_{50} of 5.9 μM in the presence of 10ng/ml IGF-1. Studies using BRDU incorporation to assess DNA synthesis have also shown biphasic effects in MCF-7 cells. (Hsu et al., 1999) found that in MCF-7 cells, biochanin A (the precursor to genistein) concentrations of less than 10 $\mu g/ml$ increased DNA synthesis by up to 280% of the control while greater than 30 $\mu g/ml$ significantly reduced DNA synthesis with an IC_{50} value of 40 $\mu g/ml$.

The biphasic effects on DNA synthesis caused by increasing concentrations of isoflavones in MCF-7 cells agrees with results from other studies done in varying assays including:

- direct cell counting with a Coulter Counter (Dampier et al., 2001; Schmitt et al., 2001)
- cell doubling time after 7 days incubation (Matsumura et al., 2005)
- cell number-colorimetric MTT assay (Le Bail et al., 1998)
- protein content-sulphorhodamine colorimetric assay (Sathyamoorthy & Wang, 1997; Sathyamoorthy et al., 1994; Wang et al., 1996)
- DNA content- diphenylamine assay (Fioravanti et al., 1998)
- DNA content- Krishans propidium iodide stain (Zava & Duwe, 1997)
- DNA content- Hoeshst reagent (Hsieh et al., 1998).

Studies examining MCF-7 cell number, found genistein IC₅₀ values of 36 μ M (Monti & Sinha, 1994), 39 μ M (Peterson & Barnes, 1991) and 24-38 μ M (Matsukawa et al., 1993). Our comparatively high IC₅₀ values could well be lowered if concentrations between 10 and 100 μ M had been examined, as inhibition appears to increase steeply in the cells between these concentrations. A recent paper examining DNA synthesis using the BRDU assay and expression of ER α /ER β in MCF-7 and T47D cells (Power & Thompson, 2003) found biphasic effects in both cell lines, with maximal proliferation of ~250% control at 10 μ M genistein which then decreased at 100 μ M in MCF-7 cells. Western blot analysis showed that 1 μ M genistein caused a down-regulation of ER α while having no effect on ER α in T47D cells and an upregulation of ER β in both cell types. This pattern of differential regulation of the ERs between the two cell lines was also seen with oestradiol stimulation at 1nM. This suggests that proliferation of DNA synthesis by genistein at this concentration is using similar

mechanisms to that of oestradiol, while the inhibitory action at higher concentrations appears to be only partially mediated through ER dependent pathways (Chen et al., 2003).

In the ER α negative MDA-MB 231 cells, isoflavones did not cause proliferation of DNA synthesis at lower concentrations but showed significant inhibition at higher concentrations. The IC₅₀ values of 14 μ M, 113 μ M and 15 μ M for genistein, daidzein and equol respectively are consistent with findings reported by other studies in other assay types.

The IC₅₀ for DNA synthesis, as determined through thymidine incorporation, for genistein and daidzein, were 27 μ M and 81 μ M respectively (Liu & Wang, 1997). Assays of cell number in MDA-231 cells found IC₅₀ of 15 μ M for genistein (Monti & Sinha, 1994) and in MDA-468 cells 24 μ M and 134 μ M for genistein and daidzein, respectively (Peterson & Barnes, 1991). Genistein concentrations >1 μ M markedly inhibited MDA-468 cell growth measured as DNA content (Zava & Duwe, 1997) and genistein concentrations of 0.01-10 μ M were inhibitory of DNA content in MDA-231 cells (Wang et al., 1996). (Fioravanti et al., 1998) also found that > 1 μ M genistein inhibited cell growth as measured through DNA content with 5 μ M giving 20% inhibition and significant counteraction of IGF-1 stimulation ($p < 0.01$). A 10 μ M concentration of genistein significantly inhibited cell number in MDA-MB 231 cells (Dampier et al., 2001). (Tanos et al., 2002) also showed inhibition of MDA-MB 231 cells between the genistein concentrations of 0.4-4 μ M with 54.6% of cell growth at 4 μ M using methylene blue staining method. (Shao et al., 1998a) found a genistein IC₅₀ for cell growth of 20 μ g/ml (74 μ M) for both MCF-7 and MDA-231 cells, while (Schmitt et al., 2001) showed that concentrations of Daidzein and Equol >10 μ M inhibited cell proliferation in this cell line.

6.4.2 Interactions between Isoflavones and oestradiol

In the ER α positive MCF-7 cells, the three isoflavones studied at 10 μ M concentrations all significantly inhibited 1nM oestradiol induced synthesis.

These results agree with other studies which have shown that, like the biphasic effects seen when isoflavones are given on their own, low concentrations enhance oestradiol proliferation and high concentrations ($\geq 10\mu$ M) inhibit oestradiol induced proliferation (Harris et al., 2005; Le Bail et al., 1998; Power & Thompson, 2003; Wang & Kurzer, 1998). (Kuiper et al., 1998) investigated the induction of gene expression by a combination of oestradiol with genistein and found that low concentrations of 10^{-7} M and 10^{-6} M had additive effects on the activation of transcription in both ER α and ER β .

At higher concentrations (Zava & Duwe, 1997) found that the pronounced growth-inhibitory actions of genistein over the 10-20 μ M concentration were not modified by 0.3nM E₂. (So et al., 1997) showed that inhibition in cell growth caused by 18.5 μ M genistein in MCF-7 cells was restored by 100nM oestradiol, while (Schmitt et al., 2001) also found that 1-10 μ M of daidzein and equol in combination with 100pM oestradiol reduced cell proliferation compared to the treatment with oestradiol alone. (Matsumura et al., 2005) also showed an inhibition of oestradiol stimulated growth after 7 days when combined with genistein and equol concentrations of 10^{-5} M.

(Power & Thompson, 2003) found that when genistein was combined with 1nM oestradiol and compared to cells treated with oestradiol alone it significantly decreased cell proliferation at concentrations $\geq 10\mu$ M in MCF-7 and $\geq 1\mu$ M in T47D cells. They found that ER β was upregulated when 1 μ M genistein or 1nM oestradiol was given alone but not when given in combination.

These results could be explained by a competitive reaction of the test compounds with oestradiol for the receptor subtypes, as genistein competes strongly with oestradiol, with (Kuiper et al., 1998) showing that at concentrations of 1 μ M, genistein has a greater oestrogenic potency (in the form of trans activation activity for ER α and ER β).

(Zava & Duwe, 1997) suggests that genistein, while mediating its actions via ER as an estrogen agonist, apparently also cross talks with other ER-independent cellular mechanisms at higher concentrations to inhibit cell proliferation induced by genistein through ER pathways. However, whether this activity may potentially be due to extragenomic actions of these ligands as concluded by (Harris et al., 2005) such as antioxidant activity G2M arrest of the cell cycle and/or possibly the result of ER β exerting an inhibitory action on ER α mediated gene expression, still remains to be elucidated (Matthews & Gustafsson, 2003). A recent study by (Chen et al., 2003) showed that genistein at high concentrations down regulates the SRF response element, which is involved in both ER and IGF-R mediated pathways (Duan et al., 2001). However this occurred in the presence of a pure antiestrogen ICI 162, 780 which suggests this is not purely an ER dependent mechanism.

In MDA-231 cells however, the addition of 1nM oestradiol appeared to rescue the cells slightly from the inhibitory effects of the isoflavones with an increase in synthesis of 10%, 7.9% and 11% of genistein, daidzein and equol respectively. The reason for this may be due to competitive binding for ER β , however the combinatory effects of isoflavones with oestradiol have not been investigated by others in ER negative cell lines and thus requires further investigation.

6.4.3 Interactions between oestradiol and isoflavones with Tamoxifen

In MCF-7 cells, tamoxifen inhibited DNA synthesis and this inhibition was significant from 0.1 μ M onwards with an IC₅₀ of 1.2 μ M. These results agree with other studies in this cell line (Faye et al., 1987; Jordan et al., 1985; Power & Thompson, 2003; Tanos et al., 2002; Taylor et al., 1984).

Concentrations of 1 μ M tamoxifen also inhibited 1 nM oestradiol induced synthesis by 28% ($P < 0.01$). This is consistent with results by (Wang & Kurzer, 1998) who showed a decrease in DNA synthesis of 23% at these concentrations. Other studies have also shown inhibitory effects of tamoxifen on oestradiol induced proliferation at varying concentrations (Power & Thompson, 2003; Schmitt et al., 2001). The combination of 1 μ M tamoxifen and 1 nM oestradiol induced the same pattern of ER α / β regulation as tamoxifen alone in MCF-7 cells (Power & Thompson, 2003). Tamoxifen has been shown to counteract oestradiol induced synthesis in part by binding and causing transcriptional activation of ER β (Paech et al., 1997).

Conversely, in this study the antiestrogenic actions of 1 μ M tamoxifen which inhibited DNA synthesis to 60% when given alone or by 28% when given with oestradiol, was completely counteracted when combined with 10 μ M genistein and daidzein such that DNA synthesis actually increased by 22% and 15% respectively ($P < 0.05$) compared with isoflavones given alone.

This is consistent with findings by (Wang & Kurzer, 1998) who showed that tamoxifen at concentrations of 0.1-1 μ M enhanced the oestrogenic effects of low concentrations (0.1-10 μ M) of genistein.

In contrast, others have found that although genistein may override the growth inhibitory actions of tamoxifen at concentrations of 10 nM-1 μ M, tamoxifen still reduces genistein induced protein content when compared with genistein given

alone (Zava & Duwe, 1997). While at 0.01 μ M tamoxifen can further inhibit cell proliferation of 0-37 μ M genistein concentrations (Tanos et al., 2002) and decrease expression of genistein stimulated pS2 mRNA expression (Wang et al., 1996).

It is interesting to note that the increase we saw in DNA synthesis with genistein and daidzein was not seen when tamoxifen was combined with 10 μ M equol and the reasons for this requires further investigation. Studies by (Sathyamoorthy & Wang, 1997) have found equol to be 100 fold more potent than daidzein in stimulating pS2 expression in MCF-7 cells. They found that 1 μ M tamoxifen was able to completely inhibit pS2 expression by 0.1 μ M daidzein but could only partially inhibit equol stimulation (55%) at this concentration. Equol was far more effective at inhibiting the binding of oestradiol to ER with 0.05 μ M resulting in 50% inhibition compared to 20% for Daidzein at the same concentration. A study by (Magee et al., 2006) has also found that R and S-equol enantiomers have different biological properties with the Racemic equol responsible for antioxidant effects preventing DNA damage in MCF-7 cells.

In the ER α negative MDA 231 cells, the opposite was true. The addition of genistein and equol caused a further inhibition of 11% and 22% respectively than antiproliferative effects seen with 1 μ M tamoxifen alone (74.5%) or with 1nM Estradiol alone (68.2%). This synergistic inhibition has also been reported by (Shen et al., 1999) who found that 5 μ M tamoxifen and genistein decreased cell inhibition to 44% of the control in MDA-435 cells and agrees with results by (Schmitt et al., 2001; Tanos et al., 2002). An explanation for the observed synergistic anti-proliferative action may be the result of tamoxifen and genistein attacking different targets in signal transduction at different phases in the cell cycle (G1 and G2M respectively)(Shen et al., 1999). Why this synergism was

not seen in MCF-7 cells is a matter for discussion, however, it would seem that the presence of ER α complicates the non-estrogenic actions of these compounds. We should note that we did not see this synergism in inhibition with daidzein and this may be explained by some reports which have shown that daidzein does not bind with any great affinity to ER β compared with genistein and equol (Kuiper et al., 1998; Matsumura et al., 2005).

6.4.4 Interactions between isoflavones and Tamoxifen with IGF-1

Early studies showed that genistein inhibited tyrosine kinase activity of growth factor receptors EGF, PDGF, insulin and IGF-1 (Akiyama et al., 1987; Akiyama & Ogawara, 1991; Kawase et al., 1995). Studies have also shown that antiestrogens can inhibit the IGF-1 induced transcriptional activation of the ER (Clayton et al., 1997; Lee et al., 1997a; Newton et al., 1994; Westley & May, 1994) with increasing substantiation of interactions between the ER and IGF-1R pathways in ER α positive breast cancer cells (Cascio et al., 2007; Kato et al., 2000; Martin & Stoica, 2002; Mendez et al., 2003; Surmacz & Bartucci, 2004). To investigate these effects, DNA synthesis was measured in both MCF-7 cells and MDA-231 cells after incubation with isoflavones in the presence of IGF-1. The stimulation of IGF-1 in MCF-7 cells was reduced by the addition of all three isoflavones and tamoxifen. However, although reductions of 29%, 16% and 28% were seen for genistein, daidzein and tamoxifen respectively, only the equol reduction of 46.2% ($P < 0.01$) was significant.

These results compare with two previous studies examining genistein inhibition of IGF-1 or insulin stimulated growth. (Hoffman, 1995) found an IC₅₀ value of 5.9 μ M genistein for inhibition of 10 ng/ml IGF-1 stimulated DNA synthesis of MCF-7 cells. (Fioravanti et al., 1998) found that 5 μ M genistein significantly

counteracted 10ng/ml IGF-1 stimulation of DNA content ($p < 0.001$). (Pagliacci et al., 1994) also found that 100 μ M genistein inhibited insulin induced tyrosine phosphorylation and a strong increase in the percentage of S phase cells. (Wang & Kurzer, 1998) found that 0.1 μ M and 1.0 μ M genistein significantly increased insulin induced DNA synthesis, while 10 μ M genistein inhibited DNA synthesis induced by 1-2 μ g/ml insulin and $\geq 25\mu$ M genistein inhibited DNA synthesis induced by 0.5-2 μ g/ml insulin ($p < 0.01$). (Panno et al., 1996) found that $>37\mu$ M Genistein inhibited insulin induced growth in MCF-7 cells. However, to our knowledge there are no other studies that have examined the effects of daidzein and equol on IGF-1 induced DNA synthesis in MCF-7 cells.

The tamoxifen reductions in IGF-1 induced DNA synthesis also agree with other studies, which have shown that 1 μ M tamoxifen can significantly inhibit IGF-1 induced proliferation in ER α positive cell lines by inhibiting IGF-1 mediated transcriptional activation of the endogenous ER (Lee et al., 1997a), down regulate IGF-1 induced tyrosine phosphorylation of IGF-1R and inhibit IRS-1 (Guvakova & Surmacz, 1997). Antiestrogens such as tamoxifen, have also been shown to down regulate the expression of IGF binding proteins (Winston et al., 1994). Certainly the interaction between oestrogens and the IGF-1 signaling pathway could explain some of the isoflavone biphasic effects seen in ER positive cell lines and possibly the synergistic effects seen with combinations of tamoxifen and isoflavones.

Oestradiol has been shown to upregulate IGF-1 receptor mRNA expression (Stewart et al., 1990), enhance IGF-1 signalling through IRS-1 expression (Lee et al., 1999; Surmacz & Bartucci, 2004), activate MAP Kinase pathway (Migliaccio et al., 1996) and prevent apoptosis through a synergistic increase in Akt protein activity (Ahmad et al., 1999), with studies by (Martin & Stoica, 2002),

showing that IGF-1 regulates both the expression and the activity of ER α in part through the phosphatidylinositol-3-kinase/Akt pathway. More recent studies by (Cascio et al., 2007) show that both oestradiol and IGF-1 differentially regulate ER transcription at ERE and AP-1 sites. A recent study by (Chen & Wong, 2004) has looked directly at the effects of genistein at low concentrations (1 μ M) on the IGF-1 signalling pathway. They found that 1 μ M genistein like 10nM oestradiol increases protein, mRNA expression and enhances tyrosine phosphorylation of the IGF-1R and downstream signalling protein IRS-1. These effects were completely counteracted by the antiestrogens ICI 182,780 and tamoxifen, suggesting that ER and more specifically the AF-2 domain of the ER is most likely involved in mediating these effects. The author concluded that the IGF signalling pathway is at least partly responsible for the growth promoting effects of genistein in MCF-7 cells at low concentrations. However, further investigation should be done to determine if higher doses of genistein cause inhibition of the IGF-1 signalling pathway or mediate their antiproliferative actions through other mechanisms. It would also be valuable to conduct these experiments with other isoflavones and antiestrogens in combination and in cell lines that do not express ER α , such as MDA-231 cells, to understand more fully the mechanisms involved.

In MDA-231 cells the addition of IGF-1 actually reduced the ability of genistein, daidzein and equol to inhibit DNA synthesis alone by 11%, 20% and 17% respectively ($P < 0.05$). Genistein inhibited IGF-1 stimulated growth by 22% compared with IGF-1 alone, which is similar to the findings of (Fioravanti et al., 1998) who found that 5 μ M genistein was able to inhibit IGF-1 stimulated DNA content by 20%. However, we did not see this with tamoxifen, which showed no

difference in the amount of DNA inhibition with or without IGF-1 at the concentrations used.

The synergistic increase in DNA synthesis when oestradiol and IGF-1 were combined ($P < 0.01$) is in line with studies that have looked at these effects in ER α positive cell lines (Stewart et al., 1990) but have not been investigated with the MDA-231 ER α negative cell line.

These results suggest that the isoflavones are working differently to tamoxifen in that part of the isoflavone inhibitory action (possibly through ER- β) is counteracted by the addition of IGF-1 and stimulation of the IGF-1R mediated pathway. To our knowledge there are as yet no published studies that have examined the interactions of daidzein and equol with IGF-1 in MCF-7 or MDA-231 ER α negative cell lines.

6.5 Conclusion

In conclusion, the most plausible explanation for the biphasic effects of these isoflavone compounds in MCF-7 cells is that, while mediating their actions via ER as oestrogen agonists, they can also cross talk with the IGF-1 signalling pathway using this and other ER independent cellular mechanisms at higher concentrations to inhibit cell proliferation. Certainly the presence of ER α is not required for the isoflavones to inhibit tumor cell growth as seen by the results in the MDA-231 cells. The isoflavones themselves also appear to have different qualities with the metabolite equol showing greater inhibition than the parent compound daidzein in both cell lines.

Isoflavones do appear to be inhibitory in concentrations $\geq 10\mu\text{M}$ and the synergism of tamoxifen especially with equol in the ER α negative MDA-231 cells is certainly worth further investigation. The clinical concern is whether

physiological concentrations could be sustainable at these levels and the evidence for this is conflicting. Reports also indicate that not everyone produces equol as a metabolite and thus it may need to be provided in supplement form. Finally, to our knowledge, this is the first report to assess the effect of isoflavones daidzein and equol on IGF-1 induced DNA synthesis in MCF-7 and MDA-231 cells and although the results do not provide direct biochemical evidence for antagonism of the IGF-1R signalling pathway they do show that biological effects of IGF-1 on MCF-7 and MDA-231 cells are counteracted by isoflavones, especially equol.

CHAPTER SEVEN – CONCLUSION

The potential role for isoflavones in the prevention of breast cancer is yet to be fully established, whilst the mechanisms involved also remain to be further elucidated.

Certainly the intervention studies conducted here and others published to date show that these compounds have only moderate effects on hormonal status and little impact in lowering endogenous levels of IGF-1, unlike pharmacological compounds such as tamoxifen. Further well-designed and controlled intervention studies using isolated isoflavones are required to determine any differences when compared with soy protein, which appears to increase circulating levels of IGF-1 and its binding proteins.

The heterogeneity of response to isoflavone exposure, which depends on gut microflora activity, necessitates an increase in the numbers of subjects required for an adequately powered study (Bingham et al., 1998) as increasing evidence suggests that equol producers may have greater clinical benefits. Differences in lipid and antioxidant status in equol excretors versus non-excretors in our one month supplementation study would appear to support this. The effect of habitual dietary intake on the proportion of metabolites excreted remains to be determined. Further studies are also needed to determine the amounts of equol ingested directly from meat and dairy products as opposed to created through a conducive microflora environment. It may also be of interest to study supplementation with isolated equol supplements to reduce study complications.

Certainly other dietary components may well be more effective in altering IGF-1 status and this requires further investigation. In particular it may be more effective to intervene with more global dietary changes such as a low fat high fibre diet as seen with recent large long term intervention studies, rather than supplementing with isolated compounds.

With isoflavone effects on endogenous levels of IGF apparently minimal it may be that their benefits are more localised to the breast. *In vitro* results from our studies in breast cancer cell lines and others suggest that isoflavones, especially equol, may be useful in establishing the pharmacological qualities of compounds needed to overcome resistance to antiestrogens which may be occurring through the IGF-1R and ER cross-talking mechanisms. Whether or not the isoflavones' greater affinity for ER β may be conferring an ability to act synergistically with tamoxifen remains to be examined. Further investigation should also be done to examine the isoflavones' ability to increase signal transduction of the IGF-1R pathway, which occurs at low concentrations and could well explain the biphasic effects seen with these compounds.

Certainly the ability of isoflavones to increase proliferation in ER α positive breast cancer cell lines at lower concentrations should not be ignored, with current opinion that breast cancer survivors and those at risk should avoid excess consumption of these "weak estrogenic compounds" until further research has been carried out. Further studies should be done to fully establish concentrations at the cellular level in populations who already consume soy isoflavones, while future intervention studies in subpopulations at higher risk of breast cancer, and in those with breast cancer, may give us the answers required.

Emerging animal study evidence suggests that short-term neonatal and pre-pubertal exposure to dietary isoflavones decreases carcinogen-induced breast cancer by increasing the proportion of differentiated cells in the mammary gland (Lamartiniere et al., 1995; Miller et al., 1994). This would support a concept that the protective effect of the traditional Asian diet occurs early in life, (Colditz & Frazier, 1995) and may explain why epidemiological studies and intervention studies, focussing on adult intake, have to date been unimpressive. Future prospective studies examining outcomes of adults who received soy based formula as infants may be of interest although a recent study comparing infants and children on cows' milk vs soya formula suggested no lasting effect of early life isoflavone exposure on isoflavone metabolism (Hoey et al., 2004). Finally, further epidemiological evidence coming from large prospective studies should point to nutritional links with breast cancer but may not be able to disentangle the beneficial effects of higher fibre/lignans/flaxseed or low fat diets that go hand in hand with soy consumption and an overall healthier diet pattern. Long term larger intervention studies with isolated isoflavones including equol, in populations at higher risk of developing breast cancer, will hopefully add to the evidence base. Certainly IGF-1 and its signalling pathway appear to be prominent targets for future treatments given they have such an important role in protecting tumour cells from apoptosis. Lowering physiological levels and expression of the IGF-1R and its ligand IGF-1 also appear to be important components in the prevention of metastasis, with antiestrogenic treatment influencing levels *in vivo*. Just how effective natural antiestrogenic compounds like isoflavones are in lowering IGF-1 production, influencing the signalling pathway and ultimately breast cancer risk remains to be determined and certainly requires further research.

APPENDIX

Table 5.A.1: Correlations of equol excretion with serum lipids, serum antioxidants, dietary intake (pre and post-menopausal women) and anthropometric measurements.

	Equol (ug/day) P(1-3)	Equol (ug/day) P(26-28)	Equol (ug/day) S(1-3)	Equol (ug/day) S(26-28)
LDL P(1-3)				R=0.487 P=0.034 N=19
LDL S(26-28)				R=0.571 P=0.011 N=19
MDA P(1-3)				R=0.676 P=0.001 N=19
CHOL P(1-3)				R=0.579 P=0.009 N=19
CHOL S(26-28)				R=0.528 P=0.020 N=19
TRIG P(1-3)				R=0.685 P=0.001 N=19
TRIG S(1-3)		R=0.486 P=0.030 N=20		R=0.577 P=0.010 N=19
RET P(1-3)				R=0.487 P=0.034 N=19
g-TOC P(1-3)				R=0.873 P=0.000 N=19
g-TOC P(26-28)		R=0.469 P=0.037 N=20		
a-TOC P(1-3)				R=0.630 P=0.004 N=19
a-TOC S(1-3)		R=0.475 P=0.035 N=20		
LYCO P(1-3)			R=0.593 P=0.007 N=19	
Iodine PFFQ				R=0.590 P=0.016 N=16
%Alcohol PFFQ	R=0.823 P=0.000 N=17			
Carbo SFFQ			R=0.498 P=0.036 N=18	
Energy Kcal SFFQ			R=0.548 P=0.019 N=18	

Starch SFFQ			R=0.639 P=0.004 N=18	
Sodium SFFQ			R=0.500 P=0.035 N=18	
Iron SFFQ			R=0.548 P=0.019 N=18	
%Protein SFFQ		R=0.555 P=0.017 N=18		
%alcohol SFFQ	R=0.815 P=0.000 N=17			
Water PFD			R=-0.583 P=0.014 N=17	
Fat PFD	R=0.552 P=0.022 N=17			
Energy Kcal PFD	R=0.508 P=0.037 N=17			
PUFA PFD	R=0.528 P=0.029 N=17			R=0.551 P=0.022 N=17
Chol PFD	R=0.615 P=0.009 N=17			
Starch PFD			R=0.518 P=0.033 N=17	
Selenium PFD		R=0.482 P=0.050 N=17	R=0.576 P=0.016 N=17	
Vit D PFD				R=0.516 P=0.034 N=17
Niacin PFD	R=0.497 P=0.042 N=17			
%alcohol PFD	R=0.508 P=0.037 N=17			
Vit D SFD				R=0.577 P=0.019 N=16
% alcohol SFD	R=0.748 P=0.001 N=17			
Weight				R=0.503 P=0.028 N=19
Waist				R=0.588 P=0.013 N=17
BMI				R=0.517 P=0.023 N=19

Data compared using pearson correlation and presented (r, P)

Table 5.A.2: Correlations of Enterolactone excretion with serum lipids, serum antioxidants and dietary intake (pre and postmenopausal women).

	Enterolactone P(1-3)	Enterolactone P(26-28)	Enterolactone S(1-3)	Enterolactone S(26-28)
b-Cryp P(26-28)			R=0.506 P=0.019 N=21	R=0.554 P=0.009 N=21
a-Caro P(1-3)			R=0.531 P=0.013 N=21	
a-Caro P(26-28)				R=0.580 P=0.006 N=21
a-Caro S(1-3)	R=0.573 P=0.007 N=21	R=0.530 P=0.014 N=21		R=0.644 P=0.002 N=21
a-Caro S(26-28)		R=0.453 P=0.039 N=21	R=0.553 P=0.009 N=21	R=0.586 P=0.005 N=21
b-Caro P(1-3)			R=0.440 P=0.046 N=21	
b-Caro P(26-28)				R=0.466 P=0.033 N=21
b-Caro S(1-3)	R=0.501 P=0.021 N=21	R=0.467 P=0.033 N=21		R=0.620 P=0.003 N=21
b-Caro S(26-28)			R=0.484 P=0.026 N=21	R=0.441 P=0.045 N=21
Manganese PFD			R=0.487 P=0.035 N=19	
Carotene PFD			R=0.459 P=0.048 N=19	
Zinc SFD			R=0.546 P=0.024 N=17	
Retinol SFD	R=0.579 P=0.009 N=19	R=0.511 P=0.030 N=18		R=0.517 P=0.028 N=18
Thiamin SFD			R=0.504 P=0.039 N=17	
Riboflavin SFD			R=0.518 P=0.033 N=17	

Data compared using pearson correlation and presented (r, P)

Table 5.A.3: Correlations of Enterodiol excretion with serum lipids, serum antioxidants and dietary intake (pre and postmenopausal women)

	Enterodiol P(1-3)	Enterodiol P(26-28)	Enterodiol S(1-3)	Enterodiol S(26-28)
MDA P(1-3)			R=0.437 P=0.048 N=21	
Chol S(26-28)		R=0.436 P=0.048 N=21		
Vit C P(1-3)			R=0.483 P=0.027 N=21	
g-TOC P(1-3)	R=0.581 P=0.006 N=21	R=0.555 P=0.009 N=21		R=0.706 P=0.000 N=21
a-TOC P(1-3)		R=0.450 P=0.041 N=21		
Zeax P(1-3)	R=0.457 P=0.037 N=21	R=0.515 P=0.017 N=21		
Iodine PFFQ				R=0.611 P=0.007 N=18
Protein PFD			R=0.552 P=0.014 N=19	
Carbohydrate PFD		R=0.502 P=0.034 N=18		
Energy Kcal PFD			R=0.482 P=0.037 N=19	
Nitrogen PFD			R=0.553 P=0.014 N=19	
PUFA PFD	R=0.471 P=0.042 N=19			
Chol PFD			R=0.501 P=0.029 N=19	
Sugar PFD		R=0.482 P=0.043 N=18		
Fibre1 PFD		R=0.578 P=0.012 N=18		
Fibre2 PFD		R=0.594 P=0.009 N=18		
Potassium PFD		R=0.692 P=0.001 N=18		
Magnesium PFD		R=0.605 P=0.008 N=18		
Phosphorous PFD		R=0.515 P=0.029 N=18		
Iron PFD		R=0.473		

		P=0.041 N=19		
Carotene PFD		R=0.547 P=0.019 N=18		
Vitamin D PFD	R=0.659 P=0.002 N=19	R=0.646 P=0.004 N=18		
Vitamin E PFD	R=0.610 P=0.006 N=19			
Niacin PFD			R=0.533 P=0.019 N=19	
Tryptophan PFD			R=0.543 P=0.016 N=19	
Vit B6 PFD		R=0.648 P=0.004 N=18		
Vit B12 PFD		R=0.682 P=0.002 N=18		
Pantothenic PFD		R=0.469 P=0.050 N=18	R=0.634 P=0.004 N=19	
Biotin PFD		R=0.508 P=0.031 N=18		
Retinol SFD	R=0.538 P=0.017 N=19			
Vit D SFD		R=0.644 P=0.004 N=18		
%Fat SFD	R=0.511 P=0.025 N=19			
%Alcohol SFD			R=0.519 P=0.033 N=17	

Data compared using pearson correlation and presented (r, P)

Table 5.A.4: Correlations of Genistein excretion with serum lipids, serum antioxidants and dietary intake (pre and post-menopausal women)

	Genistein P(1-3)	Genistein P(26-28)	Genistein S(1-3)	Genistein S(26-28)
HDL P(26-28)				R=-0.479 P=0.038 N=19
TRIG P(1-3)				R=0.525 P=0.021 N=19
TRIG P(26-28)				R=0.457 P=0.049 N=19
TRIG S(26-28)				R=0.532 P=0.019 N=19
Retin P(1-3)	R=0.486 P=0.025 N=21			
Vit C S(26-28)				R=-0.482 P=0.043 N=18
Lutein P(1-3)			R=-0.451 P=0.046 N=20	
Lutein P(26-28)	R=0.492 P=0.023 N=21			
Lutein S(1-3)	R=0.544 P=0.011 N=21			
Lutein S(26-28)	R=0.501 P=0.021 N=21			
Zeax P(26-28)			R=-0.463 P=0.040 N=20	
Zeax S(26-28)				R=-0.486 P=0.035 N=19
BCryp P(1-3)	R=0.464 P=0.034 N=21			
Bcryp P(26-28)	R=0.467 P=0.033 N=21			
a-Caro P(1-3)	R=0.439 P=0.046 N=21		R=-0.542 P=0.013 N=20	
b-Caro P(1-3)			R=-0.630 P=0.003 N=20	
b-Caro S(26-28)			R=-0.446 P=0.049 N=20	
Lyco P(1-3)			R=-0.658 P=0.002 N=20	
Carotene PFFQ	R=0.527 P=0.020 N=19			
%Alcohol PFFQ		R=0.687		

		P=0.002 N=18		
Water PFD				R=0.524 P=0.026 N=18
Iodine PFD				R=0.483 P=0.043 N=18
Niacin PFD		R=0.560 P=0.016 N=18		
Vitamin C PFD	R=0.517 P=0.023 N=19			
%Alcohol PFD		R=0.494 P=0.037 N=18		
Carotene SFD				R=-0.526 P=0.030 N=17
Thiamin SFD	R=0.525 P=0.021 N=19			
Riboflavin SFD	R=0.462 P=0.046 N=19			
%Protein SFD		R=-0.478 P=0.045 N=18	R=-0.501 P=0.041 N=17	
%Alcohol SFD		R=0.540 P=0.021 N=18		
Age				R=0.512 P=0.025 N=19
Weight				R=0.648 P=0.003 N=19
Waist				R=0.579 P=0.015 N=17
Hip				R=0.659 P=0.004 N=17
BMI				R=0.574 P=0.010 N=19

Data compared using pearson correlation and presented (r, P)

Table 5.A.5: Correlations of Daidzein excretion with serum lipids, serum antioxidants, dietary intake and anthropometric measurements (pre and postmenopausal women)

	Daidzein P(1-3)	Daidzein P(26-28)	Daidzein S(1-3)	Daidzein S(26-28)
RETIN P(26-28)		R=0.449 P=0.041 N=21		
G-Toc S(1-3)			R=0.498 P=0.022 N=21	
G-Toc S(26-28)			R=0.485 P=0.026 N=21	
A-Toc S(1-3)		R=0.464 P=0.034 N=21		
b-Cryp P(26-28)			R=0.445 P=0.043 N=21	
b-Cryp S(26-28)			R=0.539 P=0.012 N=21	
a-Caro P(26-28)		R=0.469 P=0.032 N=21		
Carbo PFD				R=0.710 P=0.001 N=19
Energy Kcal PFD				R=0.548 P=0.015 N=19
Starch PFD				R=0.589 P=0.008 N=19
Sugar PFD				R=0.54- P=0.017 N=19
Fibre 1 PFD		R=0.550 P=0.018 N=18	R=0.478 P=0.039 N=19	R=0.811 P=0.000 N=19
Fibre 2 PFD		R=0.627 P=0.005 N=18	R=0.512 P=0.025 N=19	R=0.689 P=0.001 N=19
Potassium PFD		R=0.595 P=0.009 N=18		R=0.574 P=0.010 N=19
Magnesium PFD		R=0.584 P=0.011 N=18		R=0.620 P=0.005 N=19
Phosphorous PFD				R=0.479 P=0.038 N=19
Iron PFD		R=0.527 P=0.025 N=18		R=0.565 P=0.012 N=19
Copper PFD		R=0.619 P=0.006 N=18		
Zinc PFD		R=0.558 P=0.016		R=0.493 P=0.032

		N=18		N=19
Manganese PFD	R=0.604 P=0.006 N=19	R=0.780 P=0.000 N=18	R=0.70 P=0.001 N=19	R=0.630 P=0.004 N=19
Riboflavin PFD		R=0.536 P=0.022 N=18		
Nicain PFD		R=0.522 P=0.026 N=18		
VitB6 PFD		R=0.610 P=0.007 N=18		R=0.560 P=0.013 N=19
Vit B12 PFD		R=0.607 P=0.001 N=18		
Folate PFD		R=0.562 P=0.015 N=18	R=0.528 P=0.020 N=19	
Pantothenic PFD		R=0.692 P=0.001 N=18	R=0.516 P=0.024 N=19	R=0.581 P=0.009 N=19
Biotin PFD		R=0.557 P=0.016 N=18	R=0.508 P=0.026 N=19	
Water SFD			R=0.543 P=0.024 N=17	
Protein SFD			R=0.577 P=0.015 N=17	
Fat SFD		R=0.577 P=0.012 N=18		
Cabo SFD		R=0.474 P=0.047 N=18		R=0.584 P=0.011 N=18
Energy Kcal SFD		R=0.549 P=0.018 N=18		R=0.512 P=0.030 N=18
Nitrogen SFD			R=0.577 P=0.015 N=17	
SFA SFD		R=0.468 P=0.050 N=18		
MUFA SFD		R=0.588 P=0.010 N=18		
Sugar SFD				R=0.543 P=0.020 N=18
Fibre 1 SFD		R=0.557 P=0.016 N=18		R=0.482 P=0.043 N=18
Fibre 2 SFD		R=0.523 P=0.026 N=18	R=0.513 P=0.035 N=17	R=0.552 P=0.018 N=18
Magnesium SFD		R=0.526 P=0.025 N=18	R=0.653 P=0.004 N=17	R=0.534 P=0.022 N=18
Phosphorous			R=0.606	R=0.479

SFD			P=0.010 N=17	P=0.044 N=18
Iron SFD		R=0.667 P=0.003 N=18	R=0.644 P=0.005 N=17	R=0.559 P=0.016 N=18
Copper SFD	R=0.544 P=0.016 N=19	R=0.616 P=0.007 N=18	R=0.574 P=0.016 N=17	
Zinc SFD		R=0.485 P=0.041 N=18	R=0.670 P=0.003 N=17	
Manganese SFD	R=0.605 P=0.006 N=19	R=0.683 P=0.002 N=18	R=0.745 P=0.001 N=17	
Selenium SFD		R=0.621 P=0.006 N=18	R=0.635 P=0.006 N=17	R=0.668 P=0.002 N=18
Retinol SFD		R=0.542 P=0.020 N=18		
Vit D SFD				R=0.493 P=0.038 N=18
Thiamin SFD		R=0.614 P=0.007 N=18	R=0.541 P=0.025 N=17	
Riboflavin SFD			R=0.698 P=0.002 N=17	
Niacin SFD			R=0.649 P=0.005 N=17	R=0.551 P=0.018 N=18
Tryptophan SFD			R=0.555 P=0.021 N=17	
Vit B6 SFD			R=0.541 P=0.025 N=17	R=0.471 P=0.048 N=18
Vit B12 SFD		R=0.632 P=0.005 N=18	R=0.667 P=0.003 N=17	R=0.778 P=0.000 N=18
Folate SFD			R=0.563 P=0.019 N=17	
Pantothenic SFD	R=0.461 P=0.047 N=19	R=0.533 P=0.023 N=18	R=0.611 P=0.009 N=17	
Biotin SFD	R=0.522 P=0.022 N=19			
Age			R=0.495 P=0.022 N=21	

Data compared using pearson correlation and presented (r, P)

Table 5.A.6: Correlations of serum concentrations of Isoflavones (ug/day) in Pre-menopausal women with Serum IGF-1, BP-3 and BP-1 throughout the menstrual cycle.

	Genistein Supp/post	Daidzein Supp/post	Equol Placeb/ba se	Equol Placeb/po st	Equol Supp/base	Equol Supp/post
IGF1 Placebo(day1-3)					R=0.638 P=0.014 N=14	
IGF1 Placebo(day 6-8)	R=0.736 P=0.015 N=10					
IGF1 Placebo(day12-15)				R=-0.741 P=0.002 N=15		
IGF1 Placebo(day21-23)						R=0.548 P=0.042 N=14
IGF1 Supp(day1-3)		R=0.619 P=0.018 N=14	R=0.541 P=0.046 N=14			
IGF1 Supp(day12-15)		R=0.600 P=0.030 N=13				
BP3 Placebo(day6-8)				R=0.631 P=0.016 N=14		
BP1 Placebo(day1-3)			R=-0.566 P=0.035 N=14			
BP1 Supp(day12-15)			R=-0.597 P=0.031 N=13			

Data compared using pearson correlation and presented (*r*, *P*)

Table 5.A.7: Correlations of serum concentrations of Enterolactone (ug/day) in pre-menopausal women with Serum IGF-1, BP-3 and BP-1 throughout the menstrual cycle.

	Enterolactone Placebo/Base	Enterolactone Placebo/post	Enterolactone Supp/post
BP-1 Placebo (day1-3)	R=0.590 P=0.026 N=14	R=0.505 P=0.055 N=15	R=0.655 P=0.011 N=14
BP-1 Placebo (day 6-8)	R=0.639 P=0.019 N=13	R=0.574 P=0.032 N=14	R=0.733 P=0.007 N=12
BP-1 Placebo (day 12-15)	R=0.528 P=0.052 N=14	R=0.454 P=0.089 N=15	R=0.716 P=0.004 N=14
BP-1 Placebo (day 21-23)	R=0.731 P=0.003 N=14	R=0.59 P=0.021 N=15	R=0.741 P=0.002 N=14
BP-1 Placebo (day 26-28)	R=0.664 P=0.010 N=14	R=0.525 N=0.045 N=15	R=0.709 P=0.005 N=14
BP-1 Supp (day 1-3)	R=0.686 P=0.007 N=14	R=0.604 P=0.017 N=15	R=0.811 P=0.000 N=14
BP-1 Supp (day 6-8)	R=0.652 P=0.011 N=14	R=0.550 P=0.034 N=15	R=0.775 P=0.001 N=14
BP-1 Supp (day 12-15)			R=0.673 P=0.012 N=13
BP-1 Supp (day 21-23)	R=0.684 P=0.007 N=14	R=0.578 P=0.024 N=15	R=0.782 P=0.001 N=14
BP-1 Supp (day 26-28)			R=0.710 P=0.004 N=14
IGF-1 Supp (day 26-28)			R=-0.670 P=0.009 N=14

Data compared using pearson correlation and presented (r, P)

Table 5.A.8: Correlations of serum concentrations of Isoflavones (ug/day) in post-menopausal women with Serum IGF-1, BP-3 and BP-1.

	Genistein Placebo base	Genistein Placebo post	Genistein Supp post	Daidzein Placebo base	Daidzein Placebo post	Daidzein Supp post	Equol Placebo Post	Equol Supp Base	Equol Supp Post	Enterelact Placebo base
IGF1 Placebo base						R=0.804 P=0.029 N=7				R=0.806 P=0.029 N=7
IGF1 Placebo Post									R=0.917 P=0.028 N=5	
IGF1 Supp Base									R=0.983 P=0.003 N=5	R=0.791 P=0.034 N=7
IGF1 Supp Post					R=0.967 P=0.002 N=6	R=0.914 P=0.004 N=7				R=0.313 R=0.034 N=7
BP3 Placebo Base									R=0.989 P=0.001 N=5	
BP3 Placebo Post							R=0.898 P=0.038 N=5		R=0.912 P=0.031 N=5	
BP3 Supp Base									R=0.938 P=0.018 N=5	
BP3 Supp Post		R=0.920 P=0.009 N=6							R=0.933 P=0.020 N=5	
BP1 Placebo Base								R=0.931 P=0.021 N=5		
BP1 Supp Base							R=-0.940 P=0.017 N=5			
BP1 Supp Post	R=-0.776 P=0.040 N=7		R=0.804 P=0.029 N=7	R=-0.881 P=0.009 N=7						

Data compared using pearson correlation and presented (r, P)

Table 5.A.9: Correlations of hormones with excretion of Isoflavones (ug/day) in post-menopausal women.

	Enterodiol P(1-3) Ug/day	Enterodiol P(26-28) Ug/day	Enterodiol S(1-3) Ug/day	Enterodiol S(26-28) Ug/day	Equol P(1-3) Ug/day	Equol S(1-3) Ug/day	Equol S(26-28) Ug/day	Daid S(26-28)	Genistein P(1-3) Ug/day	Genistein S(1-3) Ug/day	Enterolactone P(26-28) Ug/day	Enterolactone S(1-3) Ug/day
Progesterone P(1-3)											R=0.954 P=0.003 N=6	
Progesterone P(26-28)										R=0.919 P=0.003 N=7		
Progesterone S(26-28)	R=0.761 P=0.047 N=7						R=-0.970 P=0.006 N=5					
SHBG P(1-3)		R=0.834 P=0.039 N=6	R=0.875 P=0.010 N=7	R=0.836 P=0.019 N=7				R=0.906 P=0.005 N=7				
SHBG P(26-28)			R=0.809 P=0.027 N=7	R=0.837 P=0.019 N=7				R=0.852 P=0.015 N=7		R=-0.792 P=0.034 N=7		
SHBG S(1-3)				R=0.886 P=0.008 N=7						R=-0.841 P=0.018 N=7		
SHBG S(26-28)				R=0.817 P=0.025 N=7						R=-0.834 P=0.020 N=7		
Estrodiol S(1-3)						R=-0.890 P=0.043 N=5						
Estrodiol P(26-28)					R=0.939 P=0.018 N=5							
LH P(1-3)				R=0.713 P=0.042 N=7								
LH P(26-28)				R=0.787 P=0.036 N=7								
LH									R=0.797			R=0.760

S(1-3)									P=0.032 N=7			P=0.047 N=7
FSH P(1-3)				R=0.834 P=0.020 N=7						R=-0.797 P=0.032 N=7		
FSH S(1-3)										R=-0.839 P=0.018 N=7		
FSH S(26-28)				R=0.759 P=0.048 N=7						R=-0.763 P=0.046 N=7		

Data compared using pearson correlation and presented (r, P)

Table 5.A.10: Correlations of hormones with excretion of Isoflavones (ug/day) in pre-menopausal women.

	Enterodiol P(1-3) Ug/day	Enterodiol P(26-28) Ug/day	Equol P(1-3) Ug/day	Equol P(26-28) Ug/day	Equol S(1-3) Ug/day	Daid P(1-3) Ug/day	Daid S(1-3) Ug/day	Daid S(26-28) Ug/day	Genistein P(1-3) Ug/day	Enterolactone P(1-3) Ug/day	Enterolactone P(26-28) Ug/day	Enterolactone S(1-3) Ug/day	Enterolactone S(26-28) Ug/day
Progesterone S(1-3)										R=0.653 P=0.011 N=14	R=0.617 P=0.014 N=		R=0.760 P=0.002 N=14
Progesterone S(12-15)													
Progesterone S(21-23)					R=0.620 P=0.018 N=14					R=0.588 P=0.034 N=13	R=0.589 P=0.027 N=14		R=0.669 P=0.012 N=13
Progesterone S(26-28)									R=-0.708 P=0.007 N=13			R=-0.598 P=0.031 N=13	
SHBG P(1-3)			r=-0.626 P=0.017 N=14										
SHBG S(1-3)			R=-0.535 P=0.049 N=14										
SHBG P(6-8)			R=-0.580 P=0.038 N=14							R=0.645 P=0.017 N=13	R=0.539 P=0.047 N=14		R=0.696 P=0.008 N=13
SHBG S(6-8)								R=-0.583 P=0.029 N=14					
SHBG P(12-15)			R=-0.654 P=0.011 N=14					R=-0.664 P=0.013 N=13					
SHBG S(12-15)								R=-0.599 P=0.030 N=14					
SHBG P(21-23)			R=-0.563 P=0.036 N=14										
SHBG S(21-23)			R=-0.578 P=0.03 N=14					R=-0.579 P=0.030 N=14					
SHBG P(26-28)			R=-0.613 P=0.020										R=0.583 P=0.029

			N=14									N=14
SHBG S(26-28)			R=-0.554 P=0.049 N=13					R=-0.534 P=0.049 N=14				
Estrodiol P(1-3)		R=0.564 P=0.028 N=15										
Estrodiol S(1-3)												
Estrodiol P(6-8)	R=0.796 P=0.001 N=13	R=0.539 P=0.047 N=14							R=0.950 P=0.000 N=13	R=0.907 P=0.000 N=14		R=0.959 P=0.000 N=12
Estrodiol S(6-8)		R=0.568 P=0.027 N=15										
Estrodiol P(26-28)		R=-0.619 P=0.014 N=15										
LH P(1-3)							R=0.564 P=0.036 N=14					
LH P(6-8)						R=0.628 P=0.022 N=13	R=0.571 P=0.041 N=13		R=0.604 P=0.029 N=13			
LH S(6-8)						R=0.825 P=0.000 N=14	R=0.870 P=0.009 N=14		R=0.641 P=0.014 N=14			R=0.553 P=0.040 N=14
LH P(12-15)							R=0.535 P=0.049 N=14					
LH S(12-15)									R=0.661 P=0.014 N=13			
LH P(21-23)						R=0.812 P=0.000 N=14	R=0.555 P=0.039 N=14		R=0.702 P=0.005 N=14			R=0.534 P=0.049 N=14
LH S(21-23)												
LH P(26-28)									R=0.670 P=0.009 N=14			
LH S(26-28)						R=0.573 P=0.041 N=13			R=0.732 P=0.004 N=13			R=0.648 P=0.017 N=13
FSH												

P(1-3)													
FSH S(1-3)						R=0.706 P=0.005 N=14	R=0.820 P=0.000 N=14						
FSH P(6-8)						R=0.771 P=0.002 N=13	R=0.851 P=0.000 N=13					R=0.577 P=0.039 N=13	
FSH S(6-8)						R=0.744 P=0.002 N=14	R=0.866 P=0.000 N=14					R=0.540 P=0.046 N=14	
FSH P(12-15)						R=0.849 P=0.000 N=14	R=0.880 P=0.000 N=14					R=0.623 P=0.017 N=14	
FSH S(12-15)						R=0.796 P=0.001 N=13	R=0.826 P=0.001 N=13		R=0.614 P=0.026 N=13			R=0.767 P=0.002 N=13	
FSH P(21-23)						R=0.871 P=0.000 N=14	R=0.829 P=0.000 N=14		R=0.562 P=0.036 N=14			R=0.645 P=0.013 N=14	
FSH S(21-23)							R=0.635 P=0.015 N=14						
FSH P(26-28)				R=0.690 P=0.004 N=15						R=0.671 P=0.009 N=14	R=0.688 P=0.005 N=14	R=0.677 P=0.008 N=14	R=0.576 P=0.031 N=14
FSH S(26-28)						R=0.765 P=0.002 N=13	R=0.709 P=0.007 N=13		R=0.713 P=0.006 N=13			R=0.696 P=0.008 N=13	

Data compared using pearson correlation and presented (r, P)

Table 5.A.11: Correlations of Estradiol and Progesterone with IGF-1, BP-3 and BP-1 in pre-menopausal women (n=16).

		Estradiol Placebo (6-8)	Estradiol Placebo (21-23)	Estradiol Supplement (12-14)	Estradiol Supplement (21-23)	Proges Supplement (1-3)	Proges Supplement (6-8)	Proges Supplement (12-14)
IGF-1(1-3)	placeb		-0.54, 0.031					
	supp				-0.64, 0.008			
IGF-1(6-8)	placeb							
	supp		-0.61, 0.012		-0.72, 0.002			
IGF-1(12-14)	placeb							
	supp				-0.76, 0.001			
IGF-1 (21-23)	placeb				-0.64, 0.008			
	supp				-0.65, 0.006			-0.55, 0.032
IGF-1 (26-28)	placeb							
	supp				-0.53, 0.04	-0.54, 0.038		-0.68, 0.008
BP-3 (1-3)	Placeb							
	supp				-0.51, 0.045			
BP-3 (6-8)	placeb			-0.60, 0.030				
	supp				-0.56, 0.023			
BP-3 (12-14)	Place							
	supp				-0.53, 0.041		-0.58, 0.025	
BP-3 (21-23)	Placeb							
	supp							
BP-3 (26-28)	Placeb							
	supp							
BP-1 (1-3)	placebo	0.61, 0.020				0.508, 0.045		0.69, 0.004
	supp	0.74, 0.003				0.79, 0.000		0.78, 0.001
BP-1 (6-8)	placebo	0.66, 0.010				0.63, 0.016		0.75, 0.003
	supp	0.68, 0.007			0.56, 0.023	0.65, 0.006		0.64, 0.01
BP-1 (12-14)	placebo	0.56, 0.04			0.53, 0.034	0.67, 0.006		0.71, 0.003
	supp				0.505, 0.055	0.59, 0.022		0.53, 0.043
BP-1 (21-23)	placebo	0.73, 0.003				0.55, 0.026		0.63, 0.009
	supp	0.69, 0.006			0.511, 0.043	0.69, 0.003		0.66, 0.008
BP-1 (26-28)	Placebo	0.69, 0.006				0.61, 0.012		0.62, 0.013
	supp				0.63, 0.012	0.49, 0.063		0.63, 0.016

Data compared using pearson correlation and presented (r, P).

Table 5.A.12: Correlations of SHBG with IGF-1 and BP-1 in pre-menopausal women (n=16).

		SHBG (Placebo phase)					SHBG (Supplement phase)				
		1-3	6-8	12-14	21-23	26-28	1-3	6-8	12-14	21-23	26-28
IGF-1(1-3)	placebo		-0.54, 0.036				-0.48, 0.061		-0.54, 0.036		-0.54, 0.039
	supp			-0.53, 0.044							
IGF-1(6-8)	placebo										
	supp		-0.55, 0.034	-0.56, 0.029					-0.54, 0.034		
IGF-1 (12-14)	placebo										
	supp			-0.57, 0.032							
	supp			-0.62, 0.014							
BP-1 (1-3)	placebo	0.55, 0.027	0.74, 0.002	0.73, 0.002	0.6, 0.014	0.61, 0.011	0.61, 0.012		0.58, 0.022	0.58, 0.017	0.60, 0.019
	supp	0.59, 0.016	0.75, 0.001	0.77, 0.001	0.63, 0.009	0.66, 0.005	0.60, 0.015		0.59, 0.019	0.56, 0.023	0.62, 0.013
BP-1 (6-8)	placebo	0.62, 0.019	0.76, 0.001	0.74, 0.002	0.63, 0.016	0.65, 0.011	0.60, 0.023		0.63, 0.022	0.57, 0.034	0.61, 0.028
	supp	0.63, 0.009	0.78, 0.001	0.75, 0.001	0.67, 0.004	0.66, 0.005	0.66, 0.005	0.51, 0.042	0.61, 0.017	0.61, 0.011	0.66, 0.008
BP-1 (12-14)	placebo	0.55, 0.027	0.62, 0.014	0.62, 0.014	0.56, 0.023	0.58, 0.019	0.49, 0.053				
	supp	0.50, 0.055	0.54, 0.050	0.57, 0.032	0.52, 0.048	0.57, 0.028	0.51, 0.051				
BP-1 (21-23)	placebo		0.68, 0.005	0.66, 0.007		0.53, 0.036	0.54, 0.031		0.53, 0.043	0.50, 0.051	0.57, 0.026
	supp		0.59, 0.021	0.65, 0.09		0.50, 0.049					0.48, 0.07
BP-1 (26-28)	placebo		0.62, 0.013	0.56, 0.037		0.50, 0.049					0.50, 0.06
	supp	0.55, 0.033	0.57, 0.032								

Data compared using pearson correlation and presented (r, P)

Table 5.A.13: Correlations of hormones with IGF-1, BP-1 and BP-3 in post - menopausal women (n=7).

	LH (P1)	LH (S28)	Prog (P28)	Prog (S28)	SHBG (P1)	SHBG (P28)
BP1 (P1)	0.78, 0.037					
BP1 (S1)			0.75, 0.050			
BP3 (S1)		-0.78, 0.039				
IGF-1 (P1)					0.81, 0.028	0.77, 0.043
IGF-1 (P28)			-0.85, 0.016	-0.93, 0.0023		
IGF-1 (S28)					0.79, 0.033	

Data compared using pearson correlation and presented (r, P)

Table 5.A.14: Correlations of serum concentrations of IGF-1 & BP-1 and Serum Lipids (no correlations seen with BP-3) in pre-menopausal women.

	HDL S(1-3)	HDL P(26- 28)	HDL S(26- 28)	Trig P(1-3)	Trig S(1-3)	Trig P(26- 28)	Trig S(26- 28)	Chol P(1-3)
BP1 S(1-3)	R=0.622 P=0.010 N=16	R=0.584 P=0.018 N=16	R=0.549 P=0.028 N=16					
BP1 S(6-8)	R=0.551 P=0.027 N=16				R=- 0.523 P=0.038 N=16			
BP1 P(12- 15)	R=0.555 P=0.026 N=16							
BP1 S(12- 15)	R=0.756 P=0.001 N=15	R=0.533 P=0.041 N=15	R=0.629 P=0.012 N=15					
BP1 P(21- 23)					R=- 0.538 P=0.032 N=16			
BP1 S(21- 23)	R=0.600 P=0.014 N=16							
BP1 S(26- 28)	R=0.635 P=0.011 N=15		R=0.522 P=0.046 N=15					
IGF-1 P(1-3)				R=0.748 P=0.001 N=16	R=0.676 P=0.004 N=16		R=0.699 P=0.003 N=16	R=0.504 P=0.046 N=16
IGF-1 S(1-3)				R=0.510 P=0.027 N=16	R=0.822 P=0.000 N=16	R=0.565 P=0.022 N=16	R=0.538 P=0.032 N=16	
IGF-1 P(6-8)					R=0.608 P=0.023 N=14			
IGF-1 S(6-8)					R=0.576 P=0.019 N=16			
IGF-1 S(12- 15)					R=0.671 P=0.006 N=15			
IGF-1 P(21- 23)	R=- 0.534 P=0.033 N=16	R=- 0.509 P=0.044 N=16		R=0.713 P=0.002 N=16	R=0.594 P=0.015 N=16	R=0.502 P=0.048 N=16	R=0.678 P=0.004 N=16	
IGF-1 S(21- 23)					R=0.604 P=0.013 N=16			
IGF-1 P(26- 28)				R=0.550 P=0.027 N=16			R=0.504 P=0.047 N=16	
IGF-1 S(26- 28)		R=- 0.530 P=0.042 N=15	R=- 0.528 P=0.043 N=15					

Data compared using pearson correlation and presented (*r*, *P*)

Table 5.A.15: Correlations of serum concentrations of IGF1 with Serum antioxidants in pre menopausal women.

	IGF-1 P(1-3)	IGF-1 S(1-3)	IGF-1 P(6-8)	IGF-1 S(6-8)	IGF-1 P(12-15)	IGF-1 S(12-15)	IGF-1 P(21-23)	IGF-1 P(26-28)
Retinol P(1-3)	R=0.503 P=0.047 N=16	R=0.586 P=0.017 N=16						
Retinol S(1-3)	R=-0.655 P=0.006 N=16			R=-0.509 P=0.044 N=16			R=-0.591 P=0.016 N=16	R=-0.509 P=0.04 N=16
Vit C S(1-3)			R=-0.567 P=0.035 N=14					
Vit C S(26-28)			R=-0.540 P=0.046 N=14					
G- Tocoph P(1-3)	R=0.600 P=0.014 N=16							
G- Tocoph P(6-8)	R=-0.501 P=0.046 N=14							
A-Tocoph P(26-28)							R=0.623 P=0.010 N=16	
Lutein P(1-3)	R=0.503 P=0.047 N=16					R=0.533 P=0.041 N=15		
Zeaxanthyn P(1-3)						R=0.535 P=0.040 N=15		
Bcryp S(1-3)	R=-0.519 P=0.040 N=16				R=-0.557 P=0.025 N=16			
Bcryp P(26-28)					R=-0.495 P=0.050 N=16			
Lycopene S(1-3)	R=-0.557 P=0.025 N=16		R=-0.636 P=0.015 N=14	R=-0.610 P=0.012 N=16				
Lycopene P(26-28)			R=-0.585 P=0.028 N=14					
MDA P(1-3)	R=0.615 P=0.011 N=16				R=0.500 P=0.048 N=16			
MDA S(1-3)	R=0.533 P=0.033 N=16							

Data compared using pearson correlation and presented (r, P)

Table 5.A.16: Correlations of serum concentrations of BP1 and Serum antioxidants in premenopausal women.

	BP1 S(1-3)	BP1 P(6-8)	BP1 S(6-8)	BP1 S(21-23)	BP1 P(26-28)
B-Carotene S(1-3)	R=0.580 P=0.018 N=16	R=0.635 P=0.015 N=14	R=0.655 P=0.006 N=16	R=0.570 P=0.021 N=16	
B-Carotene S(26-28)	R=0.575 P=0.020 N=16	R=0.540 P=0.046 N=14			
A-Carotene S(1-3)	R=0.606 P=0.013 N=16	R=0.587 P=0.027 N=14	R=0.656 P=0.006 N=16	R=0.631 P=0.009 N=16	
A-Carotene P(26-28)			R=0.519 P=0.039 N=16		
A-Carotene S(26-28)	R=0.661 P=0.005 N=16	R=0.569 P=0.034 N=14			R=0.585 P=0.017 N=16
Lycopene S(1-3)		R=0.652 P=0.012 N=14	R=0.561 P=0.024 N=16		
A-Tocoph P(26-28)					R=0.532 P=0.034 N=16

Data compared using pearson correlation and presented (r, P)

Table 5.A.17: Correlations of serum concentrations of BP-3 and Serum antioxidants in pre-menopausal women.

	BP3 P(1-3)	BP3 S(1-3)	BP3 P(6-8)	BP3 S(6-8)	BP3 P(12-14)	BP3 S(12-14)	BP3 P(21-23)	BP3 S(21-23)	BP3 P(26-28)	BP3 S(26-28)
Retinol S(1-3)		R=0.540 P=0.031 N=16						R=0.576 P=0.020 N=16	R=0.718 P=0.002 N=16	R=0.738 P=0.002 N=15
Retinol P(26-28)		R=0.635 P=0.008 N=16	R=0.626 P=0.017 N=14		R=0.499 P=0.049 N=16	R=0.586 P=0.022 N=15		R=0.589 P=0.016 N=16	R=0.638 P=0.008 N=16	R=0.697 P=0.004 N=15
Retinol S(26-28)										R=0.565 P=0.028 N=15
G-tocoph S(1-3)			R=0.589 P=0.027 N=14							
G-tocoph S(26-28)	R=0.552 P=0.027 N=16	R=0.520 P=0.039 N=16	R=0.633 P=0.015 N=14	R=0.50 P=0.049 N=16	R=0.584 P=0.017 N=16	R=0.573 P=0.026 N=15	R=0.519 P=0.039 N=16		R=0.574 P=0.020 N=16	R=0.536 P=0.040 N=15
a-tocoph S(1-3)									R=0.582 P=0.018 N=16	
Zeaxan P(26-28)							R=0.575 P=0.20 N=16			
B-cryp P(1-3)							R=0.598 P=0.014 N=16			
B-cryp P(26-28)			R=0.545 P=0.044 N=14		R=0.539 P=0.031 N=16	R=0.666 P=0.007 N=15	R=0.595 P=0.015 N=16	R=0.548 P=0.028 N=16	R=0.649 P=0.006 N=16	R=0.579 P=0.024 N=15
B-cryp S(1-3)			R=0.581 P=0.029 N=14			R=0.611 P=0.015 N=15			R=0.603 P=0.013 N=16	R=0.609 P=0.016 N=15
A-Caro P(26-28)									R=0.570 P=0.021 N=16	R=0.546 P=0.035 N=15
B-Caro P(26-28)									R=0.523 P=0.038	

								N=16		
Lyco P(26-28)							R=0.613 P=0.011 N=16			
MDA S(1-3)		R=-0.588 P=0.017 N=16		R=-0.519 P=0.039 N=16				R=-0.503 P=0.047 N=16		
MDA S(26-28)		R=-0.590 P=0.016 N=16	R=-0.702 P=0.005 N=14	R=-0.550 P=0.027 N=16	R=-0.641 P=0.007 N=16		R=-0.676 P=0.004 N=16		R=-0.527 P=0.036 N=16	R=-0.514 P=0.050 N=15

Data compared using pearson correlation and presented (r, P)

Table 5.A.18: Correlations of serum concentrations of IGF-1, & BP-1 with anthropometric data in premenopausal women.
N.B post-menopausal correlations in last two rows

	Waist	Weight	Height	BMI	Hip
IGF P(1-3)	R=0.669 P=0.009 N=14	R=0.561 P=0.024 N=16		R=0.627 P=0.009 N=16	R=0.712 P=0.004 N=14
IGF-1 S(1-3)	R=0.538 P=0.047 N=14	R=0.626 P=0.009 N=16		R=0.527 P=0.036 N=16	R=0.615 P=0.019 N=14
IGF-1 P(6-8)	R=0.587 P=0.027 N=14				R=0.575 P=0.040 N=13
IGF-1 S(6-8)	R=0.501 P=0.048 N=16				
IGF-1 P(21-23)	R=0.560 P=0.024 N=16				R=0.551 P=0.041 N=14
IGF-1 S(21-23)			R=0.513 P=0.042 N=16		
IGF-1 S(26-28)		R=0.528 P=0.043 N=15			
BP1 P(1-3)		R=-0.54 P=0.031 N=16			
BP1 S(1-3)		R=-0.499 P=0.049 N=16			
BP1 P(6-8)	R=-0.557 P=0.048 N=13	R=-0.59 P=0.026 N=14		R=-0.627 P=0.016 N=14	R=-0.615 P=0.025 N=13
BP1 S(6-8)		R=-0.625 P=0.010 N=16		R=-0.586 P=0.017 N=16	R=-0.551 P=0.041 N=14
BP1 P(12-15)		R=-0.534 P=0.033 N=16			
BP1 P(21-23)		R=-0.604 P=0.013 N=16		R=-0.537 P=0.032 N=16	R=-0.600 P=0.023 N=14
BP1 S(21-23)		R=-0.642 P=0.007 N=16		R=-0.512 P=0.043 N=16	R=-0.512 P=0.043 N=16
BP1 S(26-28)		R=-0.639 P=0.010 N=15			
Postmen IGF-1 P(1-3)	R=-0.766 P=0.046 N=7				
Postmen BP-1 P(26-28)			R=0.803 P=0.030 N=7		

Data compared using pearson correlation and presented (r, P)

Table 5.A.19:Correlations of BP-1 with serum lipids in post-menopausal women.
N.B no correlations seen with BP-3 or IGF-1.

	BP1 P(1-3)	BP1 P(26-28)	BP1 S(26-28)
LDL P(1-3)		R=0.830 P=0.021 N=7	R=0.802 P=0.030 N=7
LDL P(26-28)		R=0.964 P=0.000 N=7	
LDL S(1-3)	R=0.792 P=0.034 N=7		R=0.782 P=0.038 N=7
LDL S(26-28)		R=0.899 P=0.006 N=7	
MDA P(26-28)			R=0.769 P=0.043 N=7
Chol P(1-3)			R=0.783 P=0.037 N=7
Chol P(26-28)		R=0.925 P=0.003 N=7	
Chol S(1-3)	R=0.782 P=0.038 N=7		

Data compared using pearson correlation and presented (r , P).

Table 5.A.20: Correlations of IGF-1, BP-3 and BP-1 with serum antioxidants in postmenopausal women.

	Vit C P(1-3)	Vit C S(1-3)	Gtoc P(1-3)	Lutein P(26-28)	Zeax P(1-3)	Zeax P(26-28)
IGF P(1-3)	R=-0.862 P=0.013 N=7					
IGF P(26-28)	R=-0.776 P=0.040 N=7					
IGF S(26-28)	R=-0.760 P=0.048 N=7					
BP3 P(1-3)		R=-0.777 P=0.040 N=7				
BP3 P(26-28)			R=0.822 P=0.023 N=7			R=0.787 P=0.036 N=7
BP1 S(1-3)				R=-0.815 P=0.026 N=7	R=-0.775 P=0.041 N=7	R=-0.932 P=0.002 N=7
BP1 S(26-28)				R=-0.8000 P=0.031 N=7		

Data compared using pearson correlation.

Table 5.A.21: Correlations of Serum Lipid concentrations and FFQ/FD dietary intake in pre and postmenopausal women (n=23).

	Placebo					Supplement			
Serum Lipids	PUFA FFQ	Fat FFQ	MUFA FFQ	SFA FFQ	CHOL FFQ	PUFA FFQ	Fat FFQ	MUFA FFQ	%Fat
LDL Placebo/baseline	r= 0.461 P=0.041								
LDL Placebo/post	r=0.533 P=0.016								
LDL Supp/baseline	r=0.589 P=0.006	r=0.447 P=0.048							
Cholesterol Placebo/baseline	r=0.451 P=0.046								
Cholesterol Placebo/post	r=0.547 P=0.013	r=0.480 P=0.032	r=0.479 P=0.033						
Cholesterol Supp/baseline	r=0.585 P=0.007	r=0.506 P=0.023	r=0.455 P=0.044						
MDA Placebo/post		r=0.485 P=0.030	r=0.448 P=0.048	r=0.461 P=0.041	r=0.506 P=0.023				r=0.483 P=0.026
MDA Supp/baseline		r=0.460 P=0.041				r=0.522 P=0.015	r=0.515 P=0.017	r=0.437 P=0.048	

Data compared using pearson correlation and presented (r, P)

Table 5.A.22: Correlations of Serum concentrations of antioxidants with FFQ/FD intake in pre and postmenopausal women (n=23).
Serum vitamin C vs dietary vitamin C, Serum α and γ tocopherol vs dietary Vitamin E and serum α and β carotene vs dietary Carotene.

	Vit C			Vit D			Vit E			Carotene				Biotin		Retinol
Serum	P-ffq	P-fd	S-fd	S-ffq	P-fd	S-fd	S-ffq	P-fd	S-fd	P-ffq	S-ffq	P-fd	S-fd	P-ffq	S-ffq	P-ffq
Vit C P-base										R=0.476 P=0.034		R=0.494 P=0.027				
Vit C S-base	R=0.450 P=0.046															
Vit C S-post					R=0.476 P=0.034					R=0.524 P=0.021						
Retinol P-base			R=-0.46 P=0.045		R=0.530 P=0.016											
G-toc P-base					R=0.657 P=0.002	R=0.502 P=0.029		R=0.481 P=0.032								
G-toc P-post				R=0.624 P=0.002					R=0.472 P=0.041							
G-toc S-base																
G-toc S-post																R=0.468 P=0.037
a-toc P-base					R=0.631 P=0.003	R=0.634 P=0.004										
a-toc P-post									R=0.476 P=0.039							
a-toc S-base																
a-toc S-post															R=0.452 P=0.040	
Lutein P-post										R=0.716 P=0.000		R=0.613 P=0.004				
Lutein S-base										R=0.702 P=0.001		R=0.513 P=0.021				
Lutein S-post		R=0.458 P=0.042								R=0.507 P=0.022		R=0.69 P=0.001	R=0.54 P=0.017	R=0.522 P=0.018		
Zeax					R=0.445							R=0.485				

P-base					P=0.049						P=0.03					
Zeax P-post										R=0.662 P=0.001						
Zeax S-base										R=0.654 P=0.002						
Zeax S-post										R=0.506 P=0.023						
Bcryp P-post													R=0.475 P=0.034	R=0.451 P=0.040		
Bcryp S-base											R=0.488 P=0.029					
Acaro P-base											R=0.635 P=0.003					
Acaro P-post											R=0.504 P=0.024					
Acaro S-base									R=0.487 P=0.029		R=0.682 P=0.001					
Acaro S-post											R=0.692 P=0.001	R=0.550 P=0.015				
Bcaro P-base											R=0.610 P=0.004					
Bcaro P-post											R=0.547 P=0.013					
Bcaro S-base											R=0.711 P=0.000	R=0.457 P=0.049				
Bcaro S-post											R=0.642 P=0.002	R=0.522 P=0.022	R=0.447 P=0.048			
Lyco P-post										R=0.542 P=0.014						
Lyco S-base										R=0.629 P=0.003						
Lyco S-post				R=0.627 P=0.002			R=0.50 P=0.022			R=0.468 P=0.038						

Data compared using pearson correlation and presented (r, P)

Table 5.A.23: Correlations of IGF-1 concentrations with FFQ/FD intake in pre-menopausal women.

	IGF-1 P(1-3)	IGF-1 S(1-3)	IGF-1 P(6-8)	IGF-1 S(6-8)	IGF-1 S(12-14)	IGF-1 P(21-23)	IGF-1 S(21-23)	IGF-1 P(26-28)
Carotene S FFQ			R=0.657 P=0.015 n=13					
Vit B6 S FFQ		R=0.604 P=0.022 n=14					R=0.545 P=0.044 n=14	
Vit B12 S FFQ								R=0.537 P=0.048 n=14
Vit B12 S FD	R=0.694 P=0.012 n=12			R=0.606 P=0.037 N=12				
Biotin S FFQ		R=0.627 P=0.016 n=14		R=0.691 P=0.009 n=13				
Vit C S FFQ			R=0.610 P=0.027 n=13		R=0.594 P=0.025 n=14	R=0.557 P=0.038 n=14		
PUFA S FFQ			R=0.637 P=0.019 n=13					R=0.574 P=0.032 n=14
Fat P FD								R=-0.652 P=0.012 N=13
SFA P FD								R=-0.612 P=0.026 N=13
MUFA P FD								R=-0.646 P=0.017 N=13
Energy P FD								R=-0.581 P=0.037 N=13
Starch P FD								R=-0.567 P=0.043 N=13
Sodium P FD						R=-0.621 P=0.023 N=13		R=-0.729 P=0.005 N=13
Copper P FD							R=-0.566 P=0.044 N=13	R=-0.631 P=0.021 N=13
Chloride P FD								R=-0.645 P=0.017 N=13
Selenium P FD						R=-0.602 P=0.029 N=13		R=-0.628 P=0.022 N=13
Vit D P FD		R=0.625 P=0.022 N=13			R=0.605 P=0.037 N=12			
Vit D S FD		R=0.755 P=0.005 N=12				R=0.592 P=0.042 N=12		R=0.630 P=0.028 N=12
Vit E S FD			R=-0.671 P=0.024 N=11	R=-0.587 P=0.045 N=12				
Fibre (Englyst) S FD	R=-0.725 P=0.008 N=12							

Data compared using pearson correlation and presented (r, P)

Table 5.A.24: Correlations of BP-1 concentrations with FFQ/FD intake in pre-menopausal women.

	BP1 P(1-3)	BP1 S(1-3)	BP1 P(6-8)	BP1 S(6-8)	BP1 P(12-14)	BP1 S(12-14)	BP1 P(21-23)	BP1 S(21-23)	BP1 P(26-28)	BP1 S(26-28)
Carotene P FFQ								R=0.557 P=0.048 N=13		
%Carbo P FFQ						R=0.668 P=0.018 N=12				
Protein S FFQ							R=-0.558 P=0.027 N=14			
Magnesium S FFQ							R=-0.575 P=0.031 N=14			
Phosphorous S FFQ							R=-0.556 P=0.039 N=14			
Tryptophan S FFQ							R=-0.578 P=0.030 N=14			
Water P FD										R=-0.585 P=0.045 N=12
Manganese P FD						R=-0.588 P=0.044 N=12				
Selenium S FD							R=-0.634 P=0.027 N=12			
Iodine S FD	R=-0.597 P=0.041 N=12					R=-0.607 P=0.022 N=12			R=-0.651 P=0.022 N=12	
Vit E S FD	R=0.626 P=0.029 N=12	R=0.635 P=0.027 N=12	R=0.742 P=0.009 N=11	R=0.7221 P=0.008 N=12	R=0.745 P=0.005 N=12		R=0.649 P=0.002 N=12	R=0.608 P=0.036 N=12		
Vit B12 S FD						R=-0.655 P=0.021 N=12	R=-0.593 P=0.041 N=12			R=-0.664 P=0.026 N=11
Vit B6 P FFQ						R=0.578 P=0.049 N=12				

Data compared using pearson correlation and presented (r, P)

Table 5.A.25: Correlations of BP-3 concentrations with FFQ/FD intake in pre-menopausal women.

	BP3 P(1-3)	BP3 S(1-3)	BP3 P(6-8)	BP3 S(6-8)	BP3 P(12-14)	BP3 S(12-14)	BP3 P(21-23)	BP3 S(21-23)	BP3 P(26-28)	BP3 S(26-28)
Carbo P FFQ		R=-0.632 P=0.021 N=13	R=-0.577 P=0.049 N=12	R=-0.647 P=0.017 N=13	R=-0.580 P=0.038 N=13				R=-0.631 P=0.021 N=13	
Starch S FD			R=-0.705 P=0.015 N=11							
Sugar P FFQ		R=-0.565 P=0.044 N=13					R=-0.640 P=0.018 N=13			
Fibre Southgate P FFQ	R=-0.643 P=0.018 N=13	R=-0.658 P=0.015 N=13		R=-0.674 P=0.012 N=13	R=-0.621 P=0.023 N=13	R=-0.611 P=0.035 N=12		R=-0.577 P=0.039 N=13		
Fibre Southgate S FFQ				R=-0.546 P=0.043 N=14	R=-0.566 P=0.035 N=14		R=-0.553 P=0.040 N=14			
Fibre Englyst P FFQ	R=-0.587 P=0.035 N=13	R=-0.597 P=0.031 N=13		R=-0.560 P=0.047 N=13						
Fibre Englyst S FFQ				R=-0.541 P=0.046 N=14						
Fibre Southgate S FD	R=-0.632 P=0.028 N=12									
Calcium S FD	R=-0.725 P=0.008 N=12				R=-0.694 P=0.012 N=12		R=-0.688 P=0.013 N=12			
Magnesium P FFQ	R=-0.560 P=0.046 N=13							R=-0.567 P=0.043 N=13		
Magnesium S FFQ							R=-0.548 P=0.043 N=14			
Phosphorous S FD	R=-0.720 P=0.008 N=12						R=-0.684 P=0.014 N=12		R=-0.621 R=0.031 N=12	

Zinc S FD	R=-0.578 P=0.049 N=12									
Manganese S FFQ							R=-0.556 P=0.039 N=14			
Copper S FFQ							R=-0.545 P=0.044 N=14			
Vit D P FFQ									R=0.558 P=0.047 N=13	
% Carbo P FFQ			R=-0.581 P=0.048 N=12				R=-0.634 P=0.020 N=13		R=-0.651 P=0.016 N=13	
% Fat P FFQ						R=0.584 P=0.046 N=12			R=0.616 P=0.025 N=13	
% Protein S FFQ					R=0.536 P=0.048 N=14					
% Protein S FD										R=-0.636 P=0.035 N=11
Chol S FFQ										R=0.608 P=0.027 N=13
Chol S FD	R=-0.657 P=0.020 N=12					R=-0.600 P=0.039 N=12	R=-0.644 R=0.024 N=12		R=-0.692 P=0.013 N=12	R=-0.629 P=0.038 N=11
Vit E S FFQ			R=-0.632 P=0.021 N=13		R=-0.560 P=0.037 N=14		R=-0.583 P=0.029 N=14			
Vit C P FD					R=0.608 P=0.028 N=13			R=0.555 P=0.049 N=13		
Protein S FD	R=-0.694 P=0.012 N=12							R=-0.595 P=0.041 N=12		R=-0.666 P=0.025 N=11
Carbo					R=-0.625					

S FD					P=0.030 N=12					
Nitrogen S FD	R=-0.693 P=0.013 N=12							R=-0.585 P=0.046 N=12		R=-0.656 P=0.028 N=11
Chloride S FD	R=-0.584 P=0.046 N=12		R=-0.621 P=0.041 N=11		R=-0.579 P=0.048 N=12					
Selenium S FD	R=-0.616 P=0.033 N=12	R=-0.650 P=0.022 N=12			R=-0.613 P=0.034 N=12		R=-0.602 P=0.038 N=12	R=-0.722 P=0.008 N=12		R=-0.708 P=0.015 N=11
Tryptophan S FD	R=-0.674 P=0.016 N=12						R=-0.637 P=0.026 N=12	R=-0.595 P=0.041 N=12	R=-0.585 P=0.046 N=12	R=-0.622 P=0.041 N=11
Energy Kcal S FD	R=-0.688 P=0.013 N=12									
Sodium S FD	R=-0.602 P=0.038 N=12									
Fat S FD	R=-0.741 P=0.006 N=12									
SFA S FD	R=-0.762 P=0.004 N=12									
MUFA S FD	R=-0.672 P=0.017 N=12									

Data compared using pearson correlation and presented (r, P)

Table 5.A.26: Correlations of IGF-1 with FFQ/FD in post-menopausal women

	IGF-1 P(1-3)	IGF-1 S(1-3)	IGF-1 P(26-28)	IGF-1 S(26-28)
Water PFFQ	R=-0.759 P=0.048 N=7			R=-0.861 P=0.013 N=7
Water SFFQ				R=-0.828 P=0.021 N=7
Protein PFFQ			R=-0.916 P=0.004 N=7	
Protein SFFQ	R=-0.809 P=0.028 N=7		R=-0.953 P=0.001 N=7	R=-0.840 P=0.018 N=7
Fat PFFQ			R=-0.825 P=0.022 N=7	
Carbo PFFQ			R=-0.788 P=0.035 N=7	
Energy PFFQ			R=-0.961 P=0.001 N=7	
Energy SFFQ		R=-0.846 P=0.016 N=7		R=-0.863 P=0.012 N=7
Nitrogen PFFQ			R=-0.960 P=0.003 N=7	
Nitrogen SFFQ	R=-0.820 P=0.024 N=7		R=-0.961 P=0.001 N=7	R=-0.836 P=0.019 N=7
MUFA PFFQ			R=-0.932 P=0.002 N=7	
PUFA SFFQ			R=-0.810 P=0.027 N=7	
Starch PFFQ			R=-0.903 P=0.005 N=7	
Starch PFD				R=0.793 P=0.034 N=7
Fibre (south) PFD				R=0.768 P=0.044 N=7
Fibre (engly) PFD	R=0.813 P=0.026 N=7			
Sodium PFFQ			R=-0.865 P=0.012 N=7	
Potass PFFQ				R=-0.777 P=0.040 N=7
Potass SFFQ		R=-0.809 P=0.028 N=7		R=-0.784 P=0.037 N=7
Calcium				R=-0.818

SFFQ				P=0.025 N=7
Phosph PFFQ	R=-0.814 P=0.026 N=7		R=-0.979 P=0.000 N=7	R=-0.791 P=0.034 N=7
Phosph SFFQ	R=-0.785 P=0.037 N=7		R=-0.830 P=-0.021 N=7	R=-0.898 P=0.006 N=7
Iron PFFQ			R=-0.876 P=0.010 N=7	
Copper PFD				R=0.776 P=0.040 N=7
Zinc PFFQ	R=-0.792 P=0.034 N=7		R=-0.971 P=0.000 N=7	
Zinc SFFQ			R=-0.870 P=0.011 N=7	R=-0.871 P=0.011 N=7
Chloride PFFQ			R=-0.890 P=0.007 N=7	
Mangan PFFQ			R=-0.755 P=0.050 N=7	
Mangan PFD				R=0.795 P=0.033 N=7
Selen PFD				R=0.772 P=0.042 N=7
Iodine SFFQ				R=-0.827 P=0.022 N=7
Thiamin PFFQ			R=-0.875 P=0.010 N=7	
Riboflav PFFQ			R=-0.779 P=0.039 N=7	
Niacin PFFQ			R=-0.846 P=0.016 N=7	
Tryptop PFFQ			R=-0.901 P=0.006 N=7	
Tryptop SFFQ	R=-0.799 P=0.031 N=7		R=-0.949 P=0.001 N=7	R=-0.826 P=0.022 N=7
Vit B6 SFFQ	R=-0.876 P=0.010 N=7		R=-0.794 P=0.033 N=7	R=-0.885 P=0.008 N=7
Pantoth PFFQ			R=-0.874 P=0.010 N=7	
Pantoth SFFQ	R=-0.816 P=0.025 N=7		R=-0.945 P=0.001 N=7	R=-0.874 P=0.010 N=7

Data compared using pearson correlation and presented (r, P)

Table 5.A.27: Correlations of BP-3 and BP-1 with FFQ/FD in post-menopausal women

	BP3 P(1-3)	BP3 S(1-3)	BP3 P(26-28)	BP3 S(26-28)	BP1 P(1-3)	BP1 S(1-3)	BP1 P(26-28)	BP1 S(26-28)
Water SFD					R=-0.843 P=0.017 N=7			
Fat PFFQ						R=0.878 P=0.009 N=7		
Fat SFFQ						R=0.800 P=0.031 N=7		
Carbo PFFQ						R=0.760 P=0.048 N=7		
Energy PFFQ						R=0.869 P=0.011 N=7		
Energy SFFQ	R=-0.791 P=0.034 N=7		R=-0.807 P=0.028 N=7	R=-0.755 P=0.05 N=7		R=0.780 P=0.039 N=7		
Nitrogen SFD					R=-0.758 P=0.049 N=7			R=-0.758 P=0.048 N=7
SFA PFFQ			R=-0.775 P=0.041 N=7	R=-0.779 P=0.039 N=7				
MUFA PFFQ						R=0.803 P=0.030 N=7		
MUFA SFFQ						R=0.755 P=0.050 N=7		
PUFA SFFQ						R=0.804 P=0.029 N=7		
Chol PFD					R=-0.795 P=0.033 N=7			
Starch PFFQ						R=0.850 P=0.015 N=7		
Starch SFD					R=-0.785 P=0.037 N=7			
Sugar SFFQ		R=-0.804 P=0.029 N=7		R=-0.755 P=0.050 N=7				
Sugar SFD			R=-0.762 P=0.047 N=7					
Fibre (south) PFD							R=-0.764 P=0.046 N=7	
Fibre (south) SFD								R=-0.853 P=0.015 N=7
Sodium PFFQ						R=0.852 P=0.015 N=7		
Sodium PFD					R=-0.861 P=0.013 N=7			
Sodium SFD					R=-0.802 P=0.030 N=7			R=0.755 P=0.050 N=7
Potass SFFQ		R=-0.755 P=0.050 N=7						
Potass SFD								R=-0.831 P=0.020 N=7
Magnes								R=-0.792 P=0.034

SFD								N=7
Phos SFD								R=-0.771 P=0.042 N=7
Iron SFD								R=-0.850 P=0.015 N=7
Copper SFFQ						R=-0.833 P=0.020 N=7		R=-0.757 P=0.049 N=7
Copper SFD					R=-0.840 P=0.018 N=7			R=-0.811 P=0.027 N=7
Chloride PFFQ						R=0.855 P=0.014 N=7		
Chloride PFD					R=-0.855 P=0.014 N=7			
Chloride SFD					R=-0.772 P=0.042 N=7			
Selenium PFFQ						R=0.902 P=0.005 N=7		R=0.755 P=0.050 N=7
Iodine SFFQ						R=0.763 P=0.046 N=7		
Iodine SFD								R=-0.766 P=0.044 N=7
Carotene PFFQ					R=-0.774 P=0.041 N=7			R=-0.760 P=0.047 N=7
Carotene SFFQ					R=-0.826 P=0.022 N=7			
Carotene PFD								R=-0.777 P=0.040 N=7
Carotene SFD								R=-0.795 P=0.033 N=7
Vit E SFFQ							R=0.817 P=0.025 N=7	
Vit E SFD		R=0.767 P=0.044 N=7						
Riboflavin SFD								R=-0.791 P=0.034 N=7
Niacin SFD								R=-0.913 P=0.004 N=7
Folate PFD							R=-0.763 P=0.046 N=7	
Folate SFD								R=-0.874 P=0.010 N=7
Vit B6 PFFQ							R=0.836 P=0.019 N=7	
Biotin PFD								R=-0.938 P=0.002 N=7
Vit C PFD							R=-0.756 P=0.049 N=7	
%carbo SFD					R=0.803 P=0.03 N=7			
%Fat PFD					R=-0.804 P=0.029 N=7			

Data compared using pearson correlation and presented (r ,P

REFERENCES

- Adams, K.F., Newton, K.M., Chen, C., Emerson, S.S., Potter, J.D., White, E. & Lampe, J.W. (2003). Soy isoflavones do not modulate circulating insulin-like growth factor concentrations in an older population in an intervention trial. *J Nutr*, **133**, 1316-9.
- Adlercreutz, C.H., Goldin, B.R., Gorbach, S.L., Hockerstedt, K.A., Watanabe, S., Hamalainen, E.K., Markkanen, M.H., Makela, T.H., Wahala, K.T. & Adlercreutz, T. (1995a). Soybean phytoestrogen intake and cancer risk. *J Nutr*, **125**, 757S-770S.
- Adlercreutz, H. (1990). Western diet and Western diseases: some hormonal and biochemical mechanisms and associations. *Scand J Clin Lab Invest Suppl*, **201**, 3-23.
- Adlercreutz, H. (2002a). Phytoestrogens and breast cancer. *J Steroid Biochem Mol Biol*, **83**, 113-8.
- Adlercreutz, H. (2002b). Phyto-oestrogens and cancer. *Lancet Oncol*, **3**, 364-73.
- Adlercreutz, H., Fotsis, T., Bannwart, C., Wahala, K., Makela, T., Brunow, G. & Hase, T. (1986). Determination of urinary lignans and phytoestrogen metabolites, potential antiestrogens and anticarcinogens, in urine of women on various habitual diets. *J Steroid Biochem*, **25**, 791-7.
- Adlercreutz, H., Fotsis, T., Heikkinen, R., Dwyer, J.T., Woods, M., Goldin, B.R. & Gorbach, S.L. (1982). Excretion of the lignans enterolactone and enterodiol and of equol in omnivorous and vegetarian postmenopausal women and in women with breast cancer. *Lancet*, **2**, 1295-9.
- Adlercreutz, H., Fotsis, T., Kurzer, M.S., Wahala, K., Makela, T. & Hase, T. (1995b). Isotope dilution gas chromatographic-mass spectrometric method for the determination of unconjugated lignans and isoflavonoids in human feces, with preliminary results in omnivorous and vegetarian women. *Anal Biochem*, **225**, 101-8.
- Adlercreutz, H., Fotsis, T., Bannwart, C., Makela, T., Wahala, K., Brunow, G. & Hase, T. (1985). Assay of lignans and phytoestrogens in urine of women and in cow milk by GCMS (SIM). In *In Advances in Mass Spectrometry*, Todd, J.F.J. (ed) pp. 1293-1294. Wiley: New York.
- Adlercreutz, H., Hockerstedt, K., Bannwart, C., Bloigu, S., Hamalainen, E., Fotsis, T. & Ollus, A. (1987). Effect of dietary components, including lignans and phytoestrogens, on enterohepatic circulation and liver metabolism of estrogens and on sex hormone binding globulin (SHBG). *J Steroid Biochem*, **27**, 1135-44.
- Adlercreutz, H., Honjo, H., Higashi, A., Fotsis, T., Hamalainen, E., Hasegawa, T. & Okada, H. (1991). Urinary excretion of lignans and isoflavonoid phytoestrogens in Japanese men and women consuming a traditional Japanese diet. *Am J Clin Nutr*, **54**, 1093-100.
- Adlercreutz, H., Mousavi, Y., Clark, J., Hockerstedt, K., Hamalainen, E., Wahala, K., Makela, T. & Hase, T. (1992). Dietary phytoestrogens and cancer: in vitro and in vivo studies. *J Steroid Biochem Mol Biol*, **41**, 331-7.
- Ahmad, S., Singh, N. & Glazer, R.I. (1999). Role of AKT1 in 17beta-estradiol- and insulin-like growth factor I (IGF-I)-dependent proliferation and prevention of apoptosis in MCF-7 breast carcinoma cells. *Biochem Pharmacol*, **58**, 425-30.

- Akaza, H., Miyanaga, N., Takashima, N., Naito, S., Hirao, Y., Tsukamoto, T., Fujioka, T., Mori, M., Kim, W.J., Song, J.M. & Pantuck, A.J. (2004). Comparisons of percent equol producers between prostate cancer patients and controls: case-controlled studies of isoflavones in Japanese, Korean and American residents. *Jpn J Clin Oncol*, **34**, 86-9.
- Akiyama, T., Ishida, J., Nakagawa, S., Ogawara, H., Watanabe, S., Itoh, N., Shibuya, M. & Fukami, Y. (1987). Genistein, a specific inhibitor of tyrosine-specific protein kinases. *J Biol Chem*, **262**, 5592-5.
- Akiyama, T. & Ogawara, H. (1991). Use and specificity of genistein as inhibitor of protein-tyrosine kinases. *Methods Enzymol*, **201**, 362-70.
- Ali, O., Cohen, P. & Lee, K.W. (2003). Epidemiology and biology of insulin-like growth factor binding protein-3 (IGFBP-3) as an anti-cancer molecule. *Horm Metab Res*, **35**, 726-33.
- Allen, N.E., Appleby, P.N., Davey, G.K., Kaaks, R., Rinaldi, S. & Key, T.J. (2002). The associations of diet with serum insulin-like growth factor I and its main binding proteins in 292 women meat-eaters, vegetarians, and vegans. *Cancer Epidemiol Biomarkers Prev*, **11**, 1441-8.
- Allen, N.E., Roddam, A.W., Allen, D.S., Fentiman, I.S., Dos Santos Silva, I., Peto, J., Holly, J.M. & Key, T.J. (2005). A prospective study of serum insulin-like growth factor-I (IGF-I), IGF-II, IGF-binding protein-3 and breast cancer risk. *Br J Cancer*, **92**, 1283-7.
- Allred, C.D., Allred, K.F., Ju, Y.H., Clausen, L.M., Doerge, D.R., Schantz, S.L., Korol, D.L., Wallig, M.A. & Helferich, W.G. (2004a). Dietary genistein results in larger MNU-induced, estrogen-dependent mammary tumors following ovariectomy of Sprague-Dawley rats. *Carcinogenesis*, **25**, 211-8.
- Allred, C.D., Allred, K.F., Ju, Y.H., Goepfing, T.S., Doerge, D.R. & Helferich, W.G. (2004b). Soy processing influences growth of estrogen-dependent breast cancer tumors. *Carcinogenesis*, **25**, 1649-57.
- Allred, C.D., Allred, K.F., Ju, Y.H., Virant, S.M. & Helferich, W.G. (2001). Soy diets containing varying amounts of genistein stimulate growth of estrogen-dependent (MCF-7) tumors in a dose-dependent manner. *Cancer Res*, **61**, 5045-50.
- Al-Zahrani, A., Sandhu, M.S., Luben, R.N., Thompson, D., Baynes, C., Pooley, K.A., Luccarini, C., Munday, H., Perkins, B., Smith, P., Pharoah, P.D., Wareham, N.J., Easton, D.F., Ponder, B.A. & Dunning, A.M. (2006). IGF1 and IGFBP3 tagging polymorphisms are associated with circulating levels of IGF1, IGFBP3 and risk of breast cancer. *Hum Mol Genet*, **15**, 1-10.
- Amersham-Pharmacia-Biotech. (1998). Biotrak Cellular communication assays-Cell proliferation ELISA system vs 2. Amersham Pharmacia Biotech UK Limited: England.
- An, J., Tzagarakis-Foster, C., Scharschmidt, T.C., Lomri, N. & Leitman, D.C. (2001). Estrogen receptor beta-selective transcriptional activity and recruitment of coregulators by phytoestrogens. *J Biol Chem*, **276**, 17808-14.
- Anderson, J.A., Messina, M., Garner, S.C. (1999). Effects of phyto-oestrogens on tissues. *Nutr Res Rev*, **12**, 75-116.
- Anderson, J.W., Johnstone, B.M. & Cook-Newell, M.E. (1995). Meta-analysis of the effects of soy protein intake on serum lipids. *N Engl J Med*, **333**, 276-82.

- Andersson, B., Johannsson, G., Holm, G., Bengtsson, B.A., Sashegyi, A., Pavo, I., Mason, T. & Anderson, P.W. (2002). Raloxifene does not affect insulin sensitivity or glycemic control in postmenopausal women with type 2 diabetes mellitus: a randomized clinical trial. *J Clin Endocrinol Metab*, **87**, 122-8.
- Anonymous. (1983). A prospective multicentre trial of the ovulation method of natural family planning. III. Characteristics of the menstrual cycle and of the fertile phase. *Fertil Steril*, **40**, 773-8.
- Anonymous. (1988). Effects of adjuvant tamoxifen and of cytotoxic therapy on mortality in early breast cancer. An overview of 61 randomized trials among 28,896 women. Early Breast Cancer Trialists' Collaborative Group. *N Engl J Med*, **319**, 1681-92.
- Antignac, J.P., Cariou, R., Le Bizec, B., Cravedi, J.P. & Andre, F. (2003). Identification of phytoestrogens in bovine milk using liquid chromatography/electrospray tandem mass spectrometry. *Rapid Commun Mass Spectrom*, **17**, 1256-64.
- Arjmandi, B.H., Khalil, D.A., Smith, B.J., Lucas, E.A., Juma, S., Payton, M.E. & Wild, R.A. (2003). Soy protein has a greater effect on bone in postmenopausal women not on hormone replacement therapy, as evidenced by reducing bone resorption and urinary calcium excretion. *J Clin Endocrinol Metab*, **88**, 1048-54.
- Arjmandi, B.H., Lucas, E.A., Khalil, D.A., Devareddy, L., Smith, B.J., McDonald, J., Arquitt, A.B., Payton, M.E. & Mason, C. (2005). One year soy protein supplementation has positive effects on bone formation markers but not bone density in postmenopausal women. *Nutr J*, **4**, 8.
- Arteaga, C.L., Kitten, L.J., Coronado, E.B., Jacobs, S., Kull, F.C., Jr., Allred, D.C. & Osborne, C.K. (1989). Blockade of the type I somatomedin receptor inhibits growth of human breast cancer cells in athymic mice. *J Clin Invest*, **84**, 1418-23.
- Atkinson, C., Berman, S., Humbert, O. & Lampe, J.W. (2004). In vitro incubation of human feces with daidzein and antibiotics suggests interindividual differences in the bacteria responsible for equol production. *J Nutr*, **134**, 596-9.
- Aukema, H.M. & Housini, I. (2001). Dietary soy protein effects on disease and IGF-I in male and female Han:SPRD-cy rats. *Kidney Int*, **59**, 52-61.
- Baibas, N., Bamia, C., Vassilopoulou, E., Sdrolas, J., Trichopoulou, A. & Trichopoulos, D. (2003). Dietary and lifestyle factors in relation to plasma insulin-like growth factor I in a general population sample. *Eur J Cancer Prev*, **12**, 229-34.
- Baird, D.D., Umbach, D.M., Lansdell, L., Hughes, C.L., Setchell, K.D., Weinberg, C.R., Haney, A.F., Wilcox, A.J. & McLachlan, J.A. (1995). Dietary intervention study to assess estrogenicity of dietary soy among postmenopausal women. *J Clin Endocrinol Metab*, **80**, 1685-90.
- Bajetta, E., Ferrari, L., Celio, L., Mariani, L., Miceli, R., Di Leo, A., Zilembo, N., Buzzoni, R., Spagnoli, I., Martinetti, A., Bichisao, E. & Seregni, E. (1997). The aromatase inhibitor letrozole in advanced breast cancer: effects on serum insulin-like growth factor (IGF)-I and IGF-binding protein-3 levels. *J Steroid Biochem Mol Biol*, **63**, 261-7.
- Barkhem, T., Carlsson, B., Nilsson, Y., Enmark, E., Gustafsson, J. & Nilsson, S. (1998). Differential response of estrogen receptor alpha and estrogen receptor beta to partial estrogen agonists/antagonists. *Mol Pharmacol*, **54**, 105-12.

- Barnes, S. (1995). Effect of genistein on in vitro and in vivo models of cancer. *J Nutr*, **125**, 777S-783S.
- Barnes, S. (2004). Soy isoflavones—phytoestrogens and what else? *J Nutr*, **134**, 1225S-1228S.
- Barnes, S. & Peterson, T.G. (1995). Biochemical targets of the isoflavone genistein in tumor cell lines. *Proc Soc Exp Biol Med*, **208**, 103-8.
- Barnes, S., Sfakianos, J., Coward, L. & Kirk, M. (1996). Soy isoflavonoids and cancer prevention. Underlying biochemical and pharmacological issues. *Adv Exp Med Biol*, **401**, 87-100.
- Baum, M. (1998). Tamoxifen—the treatment of choice. Why look for alternatives? *Br J Cancer*, **78 Suppl 4**, 1-4.
- Baum, M., Budzar, A.U., Cuzick, J., Forbes, J., Houghton, J.H., Klijn, J.G. & Sahmoud, T. (2002). Anastrozole alone or in combination with tamoxifen versus tamoxifen alone for adjuvant treatment of postmenopausal women with early breast cancer: first results of the ATAC randomised trial. *Lancet*, **359**, 2131-9.
- Baxter, R.C., Binoux, M., Clemmons, D.R., Conover, C., Drop, S.L., Holly, J.M., Mohan, S., Oh, Y. & Rosenfeld, R.G. (1998). Recommendations for nomenclature of the insulin-like growth factor binding protein (IGFBP) superfamily. *Growth Horm IGF Res*, **8**, 273-4.
- Bayes-Genis, A., Conover, C.A. & Schwartz, R.S. (2000). The insulin-like growth factor axis: A review of atherosclerosis and restenosis. *Circ Res*, **86**, 125-30.
- Bernstein, L. & Ross, R.K. (1993). Endogenous hormones and breast cancer risk. *Epidemiol Rev*, **15**, 48-65.
- Berrino, F., Bellati, C., Secreto, G., Camerini, E., Pala, V., Panico, S., Allegro, G. & Kaaks, R. (2001). Reducing bioavailable sex hormones through a comprehensive change in diet: the diet and androgens (DIANA) randomized trial. *Cancer Epidemiol Biomarkers Prev*, **10**, 25-33.
- Bingham, S.A., Atkinson, C., Liggins, J., Bluck, L. & Coward, A. (1998). Phytoestrogens: where are we now? *Br J Nutr*, **79**, 393-406.
- Bingham, S.A., Luben, R., Welch, A., Wareham, N., Khaw, K.T. & Day, N. (2003). Are imprecise methods obscuring a relation between fat and breast cancer? *Lancet*, **362**, 212-4.
- Blake, E.J., Adel, T. & Santoro, N. (1997). Relationships between insulin-like growth hormone factor-I and estradiol in reproductive aging. *Fertil Steril*, **67**, 697-701.
- Bohlke, K., Cramer, D.W., Trichopoulos, D. & Mantzoros, C.S. (1998). Insulin-like growth factor-I in relation to premenopausal ductal carcinoma in situ of the breast. *Epidemiology*, **9**, 570-3.
- Bonanni, B., Johansson, H., Gandini, S., Guerrieri-Gonzaga, A., Torrisi, R., Sandri, M.T., Cazzaniga, M., Mora, S., Robertson, C., Lien, E.A. & Decensi, A. (2001). Effect of low dose tamoxifen on the insulin-like growth factor system in healthy women. *Breast Cancer Res Treat*, **69**, 21-7.
- Bonorden, M.J., Greany, K.A., Wangen, K.E., Phipps, W.R., Feirtag, J., Adlercreutz, H. & Kurzer, M.S. (2004). Consumption of *Lactobacillus acidophilus* and *Bifidobacterium longum* do not alter urinary equol excretion and plasma reproductive hormones in premenopausal women. *Eur J Clin Nutr*, **58**, 1635-42.
- Bosetti, C., Spertini, L., Parpinel, M., Gnagnarella, P., Lagioui, P., Negri, E., Franceschi, S., Montella, M., Peterson, J., Dwyer, J., Giacosa, A. & La

- Vecchia, C. (2005). Flavonoids and breast cancer risk in Italy. *Cancer Epidemiol Biomarkers Prev*, **14**, 805-8.
- Bowey, E., Adlercreutz, H. & Rowland, I. (2003). Metabolism of isoflavones and lignans by the gut microflora: a study in germ-free and human flora associated rats. *Food Chem Toxicol*, **41**, 631-6.
- Brismar, K., Fernqvist-Forbes, E., Wahren, J. & Hall, K. (1994). Effect of insulin on the hepatic production of insulin-like growth factor-binding protein-1 (IGFBP-1), IGFBP-3, and IGF-I in insulin-dependent diabetes. *J Clin Endocrinol Metab*, **79**, 872-8.
- Brooks, J.D., Ward, W.E., Lewis, J.E., Hilditch, J., Nickell, L., Wong, E. & Thompson, L.U. (2004). Supplementation with flaxseed alters estrogen metabolism in postmenopausal women to a greater extent than does supplementation with an equal amount of soy. *Am J Clin Nutr*, **79**, 318-25.
- Bruning, P.F., Van Doorn, J., Bonfrer, J.M., Van Noord, P.A., Korse, C.M., Linders, T.C. & Hart, A.A. (1995). Insulin-like growth-factor-binding protein 3 is decreased in early-stage operable pre-menopausal breast cancer. *Int J Cancer*, **62**, 266-70.
- Brzezinski, A., Adlercreutz, H., Shaoul, R., Rosler, A., Shmueli, A., Tanos, V. & Schenker, J. G. (1997). Short-term effects of phytoestrogen-rich diet on postmenopausal women. *Menopause*, **4**, 89-94.
- Brzezinski, A. & Debi, A. (1999). Phytoestrogens: the "natural" selective estrogen receptor modulators? *Eur J Obstet Gynecol Reprod Biol*, **85**, 47-51.
- Buell, P. (1973). Changing incidence of breast cancer in Japanese American women. *J. Natl. Cancer Inst.*, **51**, 1479-83.
- Cabanes, A., Wang, M., Olivo, S., DeAssis, S., Gustafsson, J.A., Khan, G. & Hilakivi-Clarke, L. (2004). Prepubertal estradiol and genistein exposures up-regulate BRCA1 mRNA and reduce mammary tumorigenesis. *Carcinogenesis*, **25**, 741-8.
- Cade, J.E., Burley, V.J. & Greenwood, D.C. (2007). Dietary fibre and risk of breast cancer in the UK Women's Cohort Study. *Int J Epidemiol*.
- Caetano, B., Le Corre, L., Chalabi, N., Delort, L., Bignon, Y.J. & Bernard-Gallon, D.J. (2006). Soya phytonutrients act on a panel of genes implicated with BRCA1 and BRCA2 oncosuppressors in human breast cell lines. *Br J Nutr*, **95**, 406-13.
- Campbell, M.J., Woodside, J.V., Honour, J.W., Morton, M.S. & Leathem, A.J. (2004). Effect of red clover-derived isoflavone supplementation on insulin-like growth factor, lipid and antioxidant status in healthy female volunteers: a pilot study. *Eur J Clin Nutr*, **58**, 173-9.
- Campbell, M.J., Woodside, J.V., Secker-Walker, J., Titcomb, A. & Leathem, A.J. (2001). IGF status is altered by tamoxifen in patients with breast cancer. *Mol Pathol*, **54**, 307-10.
- Canzian, F., McKay, J.D., Cleveland, R.J., Dossus, L., Biessy, C., Boillot, C., Rinaldi, S., Llewellyn, M., Chajes, V., Clavel-Chapelon, F., Tehard, B., Chang-Claude, J., Linseisen, J., Lahmann, P.H., Pischon, T., Trichopoulos, D., Trichopoulou, A., Zilis, D., Palli, D., Tumino, R., Vineis, P., Berrino, F., Bueno-de-Mesquita, H.B., van Gils, C.H., Peeters, P.H., Pera, G., Barricarte, A., Chirlaque, M.D., Quiros, J.R., Larranaga, N., Martinez-Garcia, C., Allen, N.E., Key, T.J., Bingham, S.A., Khaw, K.T., Slimani, N., Norat, T., Riboli, E. & Kaaks, R. (2005). Genetic variation in the growth hormone synthesis pathway in relation to circulating insulin-

- like growth factor-I, insulin-like growth factor binding protein-3, and breast cancer risk: results from the European prospective investigation into cancer and nutrition study. *Cancer Epidemiol Biomarkers Prev*, **14**, 2316-25.
- Canzian, F., McKay, J.D., Cleveland, R.J., Dossus, L., Biessy, C., Rinaldi, S., Landi, S., Boillot, C., Monnier, S., Chajes, V., Clavel-Chapelon, F., Tehard, B., Chang-Claude, J., Linseisen, J., Lahmann, P.H., Pischon, T., Trichopoulos, D., Trichopoulou, A., Zilis, D., Palli, D., Tumino, R., Vineis, P., Berrino, F., Bueno-de-Mesquita, H.B., van Gils, C.H., Peeters, P.H., Pera, G., Ardanaz, E., Chirlaque, M.D., Quiros, J.R., Larranaga, N., Martinez-Garcia, C., Allen, N.E., Key, T.J., Bingham, S.A., Khaw, K.T., Slimani, N., Norat, T., Riboli, E. & Kaaks, R. (2006). Polymorphisms of genes coding for insulin-like growth factor 1 and its major binding proteins, circulating levels of IGF-I and IGFBP-3 and breast cancer risk: results from the EPIC study. *Br J Cancer*, **94**, 299-307.
- Carmichael, P.L., Pole, J.C. & Neven, P. (2000). Modulation of endometrial transforming growth factor beta (TGFbeta) by tamoxifen. *Eur J Cancer*, **36 Suppl 4**, S42-3.
- Cascio, S., Bartella, V., Garofalo, C., Russo, A., Giordano, A. & Surmacz, E. (2007). Insulin-like growth factor 1 differentially regulates estrogen receptor-dependent transcription at estrogen response element and AP-1 sites in breast cancer cells. *J Biol Chem*, **282**, 3498-506.
- Cassidy, A. (2006). Factors affecting the bioavailability of soy isoflavones in humans. *J AOAC Int*, **89**, 1182-8.
- Cassidy, A., Bingham, S. & Setchell, K. (1995). Biological effects of isoflavones in young women: importance of the chemical composition of soyabean products. *Br J Nutr*, **74**, 587-601.
- Cassidy, A., Bingham, S. & Setchell, K.D. (1994). Biological effects of a diet of soy protein rich in isoflavones on the menstrual cycle of premenopausal women. *Am J Clin Nutr*, **60**, 333-40.
- Cassidy, A. & Hooper, L. (2006). Phytoestrogens and cardiovascular disease. *J Br Menopause Soc*, **12**, 49-56.
- Catignani, G.L. & Bieri, J.G. (1983). Simultaneous determination of retinol and alpha-tocopherol in serum or plasma by liquid chromatography. *Clin Chem*, **29**, 708-12.
- Caufriez, A., Golstein, J., Tadjerouni, A., Bosson, D., Cantraine, F., Robyn, C. & Copinschi, G. (1986). Modulation of immunoreactive somatomedin-C levels by sex steroids. *Acta Endocrinol (Copenh)*, **112**, 284-9.
- Cauley, J.A., Norton, L., Lippman, M.E., Eckert, S., Krueger, K.A., Purdie, D.W., Farrerons, J., Karasik, A., Mellstrom, D., Ng, K.W., Stepan, J.J., Powles, T.J., Morrow, M., Costa, A., Silfen, S.L., Walls, E.L., Schmitt, H., Muchmore, D.B., Jordan, V.C. & Ste-Marie, L.G. (2001). Continued breast cancer risk reduction in postmenopausal women treated with raloxifene: 4-year results from the MORE trial. Multiple outcomes of raloxifene evaluation. *Breast Cancer Res Treat*, **65**, 125-34.
- Chae, Y.H., Coffing, S.L., Cook, V.M., Ho, D.K., Cassady, J.M. & Baird, W.M. (1991). Effects of biochanin A on metabolism, DNA binding and mutagenicity of benzo[a]pyrene in mammalian cell cultures. *Carcinogenesis*, **12**, 2001-6.
- Chen, W.F., Huang, M.H., Tzang, C.H., Yang, M. & Wong, M.S. (2003). Inhibitory actions of genistein in human breast cancer (MCF-7) cells. *Biochim Biophys Acta*, **1638**, 187-96.

- Chen, W.F. & Wong, M.S. (2004). Genistein enhances insulin-like growth factor signaling pathway in human breast cancer (MCF-7) cells. *J Clin Endocrinol Metab*, **89**, 2351-9.
- Chen, Y.M., Ho, S.C., Lam, S.S., Ho, S.S. & Woo, J.L. (2004). Beneficial effect of soy isoflavones on bone mineral content was modified by years since menopause, body weight, and calcium intake: a double-blind, randomized, controlled trial. *Menopause*, **11**, 246-54.
- Chong, Y.M., Colston, K., Jiang, W.G., Sharma, A.K. & Mokbel, K. (2006a). The relationship between the insulin-like growth factor-1 system and the oestrogen metabolising enzymes in breast cancer tissue and its adjacent non-cancerous tissue. *Breast Cancer Res Treat*, **99**, 275-88.
- Chong, Y.M., Williams, S.L., Elkak, A., Sharma, A.K. & Mokbel, K. (2006b). Insulin-like growth factor 1 (IGF-1) and its receptor mRNA levels in breast cancer and adjacent non-neoplastic tissue. *Anticancer Res*, **26**, 167-73.
- Clarke, D.B., Barnes, K.A. & Lloyd, A.S. (2004). Determination of unusual soya and non-soya phytoestrogen sources in beer, fish products and other foods. *Food Addit Contam*, **21**, 949-62.
- Clayton, S.J., May, F.E. & Westley, B.R. (1997). Insulin-like growth factors control the regulation of oestrogen and progesterone receptor expression by oestrogens. *Mol Cell Endocrinol*, **128**, 57-68.
- Clemmons, D.R., Seek, M.M. & Underwood, L.E. (1985). Supplemental essential amino acids augment the somatomedin-C/insulin-like growth factor I response to refeeding after fasting. *Metabolism*, **34**, 391-5.
- Clemmons, D.R. & Underwood, L.E. (1991). Nutritional regulation of IGF-I and IGF binding proteins. *Annu Rev Nutr*, **11**, 393-412.
- Clemons, M. & Goss, P. (2001). Estrogen and the risk of breast cancer. *N Engl J Med*, **344**, 276-85.
- Clifton-Bligh, P.B., Baber, R.J., Fulcher, G.R., Nery, M.L. & Moreton, T. (2001). The effect of isoflavones extracted from red clover (Rimostil) on lipid and bone metabolism. *Menopause*, **8**, 259-65.
- Cohen, L.A., Zhao, Z., Pittman, B. & Scimeca, J.A. (2000). Effect of intact and isoflavone-depleted soy protein on NMU-induced rat mammary tumorigenesis. *Carcinogenesis*, **21**, 929-35.
- Colditz, G.A. & Frazier, A.L. (1995). Models of breast cancer show that risk is set by events of early life: prevention efforts must shift focus. *Cancer Epidemiol Biomarkers Prev*, **4**, 567-71.
- Colletti, R.B., Roberts, J.D., Devlin, J.T. & Copeland, K.C. (1989). Effect of tamoxifen on plasma insulin-like growth factor I in patients with breast cancer. *Cancer Res*, **49**, 1882-4.
- Constantinou, A., Kiguchi, K. & Huberman, E. (1990). Induction of differentiation and DNA strand breakage in human HL-60 and K-562 leukemia cells by genistein. *Cancer Res*, **50**, 2618-24.
- Constantinou, A.I., Lantvit, D., Hawthorne, M., Xu, X., van Breemen, R.B. & Pezzuto, J.M. (2001). Chemopreventive effects of soy protein and purified soy isoflavones on DMBA-induced mammary tumors in female Sprague-Dawley rats. *Nutr Cancer*, **41**, 75-81.
- Cooke, G.M. (2006). A review of the animal models used to investigate the health benefits of soy isoflavones. *J AOAC Int*, **89**, 1215-27.
- Costa, R.L. & Summa, M.A. (2000). Soy protein in the management of hyperlipidemia. *Ann Pharmacother*, **34**, 931-5.

- Cotroneo, M.S., Wang, J., Fritz, W.A., Eltoum, I.E. & Lamartiniere, C.A. (2002). Genistein action in the prepubertal mammary gland in a chemoprevention model. *Carcinogenesis*, **23**, 1467-74.
- Coward, L., Barnes, N.C., Setchell, K.D.R. & Barnes, S. (1993). Genistein, daidzein, and their .beta.-glycoside conjugates: antitumor isoflavones in soybean foods from American and Asian diets. *J. Agric. Food Chem.*, **41**, 1961-1967.
- Craft, N.E. (1992). Carotenoid reversed-phase high-performance liquid chromatography methods: reference compendium. *Methods Enzymol*, **213**, 185-205.
- Crawley, H., Great Britain. Ministry of Agriculture, F. & Food. (1988). *Food portion sizes*.
- Crouse, J.R., 3rd, Morgan, T., Terry, J.G., Ellis, J., Vitols, M. & Burke, G.L. (1999). A randomized trial comparing the effect of casein with that of soy protein containing varying amounts of isoflavones on plasma concentrations of lipids and lipoproteins. *Arch Intern Med*, **159**, 2070-6.
- Cuzick, J., Forbes, J., Edwards, R., Baum, M., Cawthorn, S., Coates, A., Hamed, A., Howell, A. & Powles, T. (2002). First results from the International Breast Cancer Intervention Study (IBIS-I): a randomised prevention trial. *Lancet*, **360**, 817-24.
- Dai, Q., Shu, X.O., Jin, F., Potter, J.D., Kushi, L.H., Teas, J., Gao, Y.T. & Zheng, W. (2001). Population-based case-control study of soyfood intake and breast cancer risk in Shanghai. *Br J Cancer*, **85**, 372-8.
- Dampier, K., Hudson, E.A., Howells, L.M., Manson, M.M., Walker, R.A. & Gescher, A. (2001). Differences between human breast cell lines in susceptibility towards growth inhibition by genistein. *Br J Cancer*, **85**, 618-24.
- Daughaday, W.H., Hall, K., Raben, M.S., Salmon, W.D., Jr., van den Brande, J.L. & van Wyk, J.J. (1972). Somatomedin: proposed designation for sulphation factor. *Nature*, **235**, 107.
- Davis, J.N., Kucuk, O. & Sarkar, F.H. (1999). Genistein inhibits NF-kappa B activation in prostate cancer cells. *Nutr Cancer*, **35**, 167-74.
- Day, J.K., Besch-Williford, C., McMann, T.R., Hufford, M.G., Lubahn, D.B. & MacDonald, R.S. (2001). Dietary genistein increased DMBA-induced mammary adenocarcinoma in wild-type, but not ER alpha KO, mice. *Nutr Cancer*, **39**, 226-32.
- Decensi, A., Bonanni, B., Guerrieri-Gonzaga, A., Gandini, S., Robertson, C., Johansson, H., Travaglini, R., Sandri, M.T., Tessadrelli, A., Farante, G., Salinaro, F., Bettega, D., Barreca, A., Boyle, P., Costa, A. & Veronesi, U. (1998). Biologic activity of tamoxifen at low doses in healthy women. *J Natl Cancer Inst*, **90**, 1461-7.
- Decensi, A., Gandini, S., Guerrieri-Gonzaga, A., Johansson, H., Manetti, L., Bonanni, B., Sandri, M.T., Barreca, A., Costa, A., Robertson, C. & Lien, E.A. (1999). Effect of blood tamoxifen concentrations on surrogate biomarkers in a trial of dose reduction in healthy women. *J Clin Oncol*, **17**, 2633-8.
- Decensi, A., Gandini, S., Serrano, D., Cazzaniga, M., Pizzamiglio, M., Maffini, F., Pelosi, G., Daldoss, C., Omodei, U., Johansson, H., Macis, D., Lazzeroni, M., Penotti, M., Sironi, L., Moroni, S., Bianco, V., Rondanina, G., Gjerde, J., Guerrieri-Gonzaga, A. & Bonanni, B. (2007). Randomized Dose-Ranging Trial of Tamoxifen at Low Doses in Hormone Replacement Therapy Users. *J Clin Oncol*.

- Decensi, A., Johansson, H., Miceli, R., Mariani, L., Camerini, T., Cavadini, E., Di Mauro, M.G., Barreca, A., Gonzaga, A.G., Diani, S., Sandri, M.T., De Palo, G. & Formelli, F. (2001). Long-term effects of fenretinide, a retinoic acid derivative, on the insulin-like growth factor system in women with early breast cancer. *Cancer Epidemiol Biomarkers Prev*, **10**, 1047-53.
- Decensi, A., Robertson, C., Viale, G., Pigatto, F., Johansson, H., Kisanga, E.R., Veronesi, P., Torrissi, R., Cazzaniga, M., Mora, S., Sandri, M.T., Pelosi, G., Luini, A., Goldhirsch, A., Lien, E.A. & Veronesi, U. (2003). A randomized trial of low-dose tamoxifen on breast cancer proliferation and blood estrogenic biomarkers. *J Natl Cancer Inst*, **95**, 779-90.
- Decroos, K., Vanhemmens, S., Cattoir, S., Boon, N. & Verstraete, W. (2005). Isolation and characterisation of an equol-producing mixed microbial culture from a human faecal sample and its activity under gastrointestinal conditions. *Arch Microbiol*, **183**, 45-55.
- Del Giudice, M.E., Fantus, I.G., Ezzat, S., McKeown-Eyssen, G., Page, D. & Goodwin, P.J. (1998). Insulin and related factors in premenopausal breast cancer risk. *Breast Cancer Res Treat*, **47**, 111-20.
- DeLellis, K., Rinaldi, S., Kaaks, R.J., Kolonel, L.N., Henderson, B. & Le Marchand, L. (2004). Dietary and lifestyle correlates of plasma insulin-like growth factor-I (IGF-I) and IGF binding protein-3 (IGFBP-3): the multiethnic cohort. *Cancer Epidemiol Biomarkers Prev*, **13**, 1444-51.
- den Tonkelaar, I., Keinan-Boker, L., Veer, P.V., Arts, C.J., Adlercreutz, H., Thijssen, J.H. & Peeters, P.H. (2001). Urinary phytoestrogens and postmenopausal breast cancer risk. *Cancer Epidemiol Biomarkers Prev*, **10**, 223-8.
- D'Ercole, A.J., Applewhite, G.T. & Underwood, L.E. (1980). Evidence that somatomedin is synthesized by multiple tissues in the fetus. *Dev Biol*, **75**, 315-28.
- Devine, A., Rosen, C., Mohan, S., Baylink, D. & Prince, R.L. (1998). Effects of zinc and other nutritional factors on insulin-like growth factor I and insulin-like growth factor binding proteins in postmenopausal women. *Am J Clin Nutr*, **68**, 200-6.
- Dewell, A., Hollenbeck, C.B. & Bruce, B. (2002). The effects of soy-derived phytoestrogens on serum lipids and lipoproteins in moderately hypercholesterolemic postmenopausal women. *J Clin Endocrinol Metab*, **87**, 118-21.
- Diamandi, A., Mistry, J., Krishna, R.G. & Khosravi, J. (2000). Immunoassay of insulin-like growth factor-binding protein-3 (IGFBP-3): new means to quantifying IGFBP-3 proteolysis. *J Clin Endocrinol Metab*, **85**, 2327-33.
- Duan, R., Xie, W., Burghardt, R.C. & Safe, S. (2001). Estrogen receptor-mediated activation of the serum response element in MCF-7 cells through MAPK-dependent phosphorylation of Elk-1. *J Biol Chem*, **276**, 11590-8.
- Duncan, A.M., Merz, B.E., Xu, X., Nagel, T.C., Phipps, W.R. & Kurzer, M.S. (1999a). Soy isoflavones exert modest hormonal effects in premenopausal women. *J Clin Endocrinol Metab*, **84**, 192-7.
- Duncan, A.M., Merz-Demlow, B.E., Xu, X., Phipps, W.R. & Kurzer, M.S. (2000). Premenopausal equol excretors show plasma hormone profiles associated with lowered risk of breast cancer. *Cancer Epidemiol Biomarkers Prev*, **9**, 581-6.

- Duncan, A.M., Underhill, K.E., Xu, X., Lavalleur, J., Phipps, W.R. & Kurzer, M.S. (1999b). Modest hormonal effects of soy isoflavones in postmenopausal women. *J Clin Endocrinol Metab*, **84**, 3479-84.
- Dushek, E.J., de Valk-de Roo, G.W., Gooren, L.J. & Netelenbos, C. (2004). Effects of conjugated equine estrogen vs. raloxifene on serum insulin-like growth factor-I and insulin-like growth factor binding protein-3: a 2-year, double-blind, placebo-controlled study. *Fertil Steril*, **82**, 384-90.
- Ellis, M.J., Coop, A., Singh, B., Mauriac, L., Llombert-Cussac, A., Janicke, F., Miller, W.R., Evans, D.B., Dugan, M., Brady, C., Quebe-Fehling, E. & Borgs, M. (2001). Letrozole is more effective neoadjuvant endocrine therapy than tamoxifen for ErbB-1- and/or ErbB-2-positive, estrogen receptor-positive primary breast cancer: evidence from a phase III randomized trial. *J Clin Oncol*, **19**, 3808-16.
- Ellis, M.J., Singer, C., Hornby, A., Rasmussen, A. & Cullen, K.J. (1994). Insulin-like growth factor mediated stromal-epithelial interactions in human breast cancer. *Breast Cancer Res Treat*, **31**, 249-61.
- Eng-Wong, J., Hursting, S.D., Venzon, D., Perkins, S.N. & Zujewski, J.A. (2003). Effect of raloxifene on insulin-like growth factor-I, insulin-like growth factor binding protein-3, and leptin in premenopausal women at high risk for developing breast cancer. *Cancer Epidemiol Biomarkers Prev*, **12**, 1468-73.
- Ernster, V.L., Wrensch, M.R., Petrakis, N.L., King, E.B., Miike, R., Murai, J., Goodson, W.H., 3rd & Siiteri, P.K. (1987). Benign and malignant breast disease: initial study results of serum and breast fluid analyses of endogenous estrogens. *J Natl Cancer Inst*, **79**, 949-60.
- Faye, J.C., Fargin, A., Valette, A. & Bayard, F. (1987). Antiestrogens, different sites of action than the estrogen receptor? *Horm Res*, **28**, 202-11.
- Fioravanti, L., Cappelletti, V., Miodini, P., Ronchi, E., Brivio, M. & Di Fronzo, G. (1998). Genistein in the control of breast cancer cell growth: insights into the mechanism of action in vitro. *Cancer Lett*, **130**, 143-52.
- Fischer, F., Schulte, H., Mohan, S., Tataru, M.C., Kohler, E., Assmann, G. & von Eckardstein, A. (2004). Associations of insulin-like growth factors, insulin-like growth factor binding proteins and acid-labile subunit with coronary heart disease. *Clin Endocrinol (Oxf)*, **61**, 595-602.
- Fisher, B., Costantino, J.P., Wickerham, D.L., Redmond, C.K., Kavanah, M., Cronin, W.M., Vogel, V., Robidoux, A., Dimitrov, N., Atkins, J., Daly, M., Wieand, S., Tan-Chiu, E., Ford, L. & Wolmark, N. (1998). Tamoxifen for prevention of breast cancer: report of the National Surgical Adjuvant Breast and Bowel Project P-1 Study. *J Natl Cancer Inst*, **90**, 1371-88.
- Fisher, B., Powles, T.J. & Pritchard, K.J. (2000). Tamoxifen for the prevention of breast cancer. *Eur J Cancer*, **36**, 142-50.
- Fletcher, O., Gibson, L., Johnson, N., Altmann, D.R., Holly, J.M., Ashworth, A., Peto, J. & Silva Idos, S. (2005). Polymorphisms and circulating levels in the insulin-like growth factor system and risk of breast cancer: a systematic review. *Cancer Epidemiol Biomarkers Prev*, **14**, 2-19.
- Fletcher, R.J. (2003). Food sources of phyto-oestrogens and their precursors in Europe. *Br J Nutr*, **89 Suppl 1**, S39-43.
- Foth, D. & Nawroth, F. (2003). Effect of soy supplementation on endogenous hormones in postmenopausal women. *Gynecol Obstet Invest*, **55**, 135-8.
- Fotsis, T., Pepper, M., Adlercreutz, H., Hase, T., Montesano, R. & Schweigerer, L. (1995). Genistein, a dietary ingested isoflavonoid, inhibits cell proliferation and in vitro angiogenesis. *J Nutr*, **125**, 790S-797S.

- Francis, C., Millington, A. & Bailey, E. (1967). The distribution of oestrogenic isoflavones in the genus *Trifolium*. *Aust. Vet J*, **18**, 47-54.
- Franke, A.A., Custer, L.J. & Tanaka, Y. (1998). Isoflavones in human breast milk and other biological fluids. *Am J Clin Nutr*, **68**, 1466S-1473S.
- Frost, V.J., Helle, S.I., Lonning, P.E., van der Stappen, J.W. & Holly, J.M. (1996). Effects of treatment with megestrol acetate, aminoglutethimide, or formestane on insulin-like growth factor (IGF) I and II, IGF-binding proteins (IGFBPs), and IGFBP-3 protease status in patients with advanced breast cancer. *J Clin Endocrinol Metab*, **81**, 2216-21.
- Fuqua, S.A., Schiff, R., Parra, I., Friedrichs, W.E., Su, J.L., McKee, D.D., Slentz-Kesler, K., Moore, L.B., Willson, T.M. & Moore, J.T. (1999). Expression of wild-type estrogen receptor beta and variant isoforms in human breast cancer. *Cancer Res*, **59**, 5425-8.
- Gagliardi, A. & Collins, D.C. (1993). Inhibition of angiogenesis by antiestrogens. *Cancer Res*, **53**, 533-5.
- Gallo, D., Ferlini, C., Fabrizi, M., Prislei, S. & Scambia, G. (2006). Lack of stimulatory activity of a phytoestrogen-containing soy extract on the growth of breast cancer tumors in mice. *Carcinogenesis*, **27**, 1404-9.
- Gann, P.H., Kazer, R., Chatterton, R., Gapstur, S., Thedford, K., Helenowski, I., Giovanazzi, S. & Van Horn, L. (2005). Sequential, randomized trial of a low-fat, high-fiber diet and soy supplementation: effects on circulating IGF-I and its binding proteins in premenopausal women. *Int J Cancer*, **116**, 297-303.
- Gardner, C.D., Newell, K.A., Cherin, R. & Haskell, W.L. (2001). The effect of soy protein with or without isoflavones relative to milk protein on plasma lipids in hypercholesterolemic postmenopausal women. *Am J Clin Nutr*, **73**, 728-35.
- Gasco, M., Argusti, A., Bonanni, B. & Decensi, A. (2005). SERMs in chemoprevention of breast cancer. *Eur J Cancer*, **41**, 1980-9.
- Giovannucci, E. (1999). Insulin-like growth factor-I and binding protein-3 and risk of cancer. *Horm Res*, **51 Suppl 3**, 34-41.
- Giovannucci, E. (2003). Nutrition, insulin, insulin-like growth factors and cancer. *Horm Metab Res*, **35**, 694-704.
- Giovannucci, E., Pollak, M., Liu, Y., Platz, E.A., Majeed, N., Rimm, E.B. & Willett, W.C. (2003). Nutritional predictors of insulin-like growth factor I and their relationships to cancer in men. *Cancer Epidemiol Biomarkers Prev*, **12**, 84-9.
- Gooch, J.L., Van Den Berg, C.L. & Yee, D. (1999). Insulin-like growth factor (IGF)-I rescues breast cancer cells from chemotherapy-induced cell death--proliferative and anti-apoptotic effects. *Breast Cancer Res Treat*, **56**, 1-10.
- Gooderham, M.H., Adlercreutz, H., Ojala, S.T., Wahala, K. & Holub, B.J. (1996). A soy protein isolate rich in genistein and daidzein and its effects on plasma isoflavone concentrations, platelet aggregation, blood lipids and fatty acid composition of plasma phospholipid in normal men. *J Nutr*, **126**, 2000-6.
- Grace, P.B., Taylor, J.I., Low, Y.L., Luben, R.N., Mulligan, A.A., Botting, N.P., Dowsett, M., Welch, A.A., Khaw, K.T., Wareham, N.J., Day, N.E. & Bingham, S.A. (2004). Phytoestrogen concentrations in serum and spot urine as biomarkers for dietary phytoestrogen intake and their relation to breast cancer risk in European prospective investigation of cancer and nutrition-norfolk. *Cancer Epidemiol Biomarkers Prev*, **13**, 698-708.

- Gu, L., House, S.E., Prior, R.L., Fang, N., Ronis, M.J., Clarkson, T.B., Wilson, M.E. & Badger, T.M. (2006). Metabolic phenotype of isoflavones differ among female rats, pigs, monkeys, and women. *J Nutr*, **136**, 1215-21.
- Guerrieri-Gonzaga, A., Robertson, C., Bonanni, B., Serrano, D., Cazzaniga, M., Mora, S., Gulisano, M., Johansson, H., Formelli, F., Intra, M., Latronico, A., Franchi, D., Pelosi, G., Johnson, K. & Decensi, A. (2006). Preliminary results on safety and activity of a randomized, double-blind, 2 x 2 trial of low-dose tamoxifen and fenretinide for breast cancer prevention in premenopausal women. *J Clin Oncol*, **24**, 129-35.
- Guidoux, S., Garnier, P.E. & Schimpff, R.M. (1986). Effect of the variations of female sex hormones during the menstrual cycle upon serum somatomedin and growth-promoting activity. *Horm Res*, **23**, 31-7.
- Gunnell, D., Oliver, S.E., Peters, T.J., Donovan, J.L., Persad, R., Maynard, M., Gillatt, D., Pearce, A., Hamdy, F.C., Neal, D.E. & Holly, J.M. (2003). Are diet-prostate cancer associations mediated by the IGF axis? A cross-sectional analysis of diet, IGF-I and IGFBP-3 in healthy middle-aged men. *Br J Cancer*, **88**, 1682-6.
- Guvakova, M.A. & Surmacz, E. (1997). Tamoxifen interferes with the insulin-like growth factor I receptor (IGF-IR) signaling pathway in breast cancer cells. *Cancer Res*, **57**, 2606-10.
- Hankinson, S.E., Willett, W.C., Colditz, G.A., Hunter, D.J., Michaud, D.S., Deroo, B., Rosner, B., Speizer, F.E. & Pollak, M. (1998). Circulating concentrations of insulin-like growth factor-I and risk of breast cancer. *Lancet*, **351**, 1393-6.
- Hargreaves, D.F., Potten, C.S., Harding, C., Shaw, L.E., Morton, M.S., Roberts, S.A., Howell, A. & Bundred, N.J. (1999). Two-week dietary soy supplementation has an estrogenic effect on normal premenopausal breast. *J Clin Endocrinol Metab*, **84**, 4017-24.
- Harper-Wynne, C.L., Sacks, N.P., Shenton, K., MacNeill, F.A., Sauven, P., Laidlaw, I.J., Rayter, Z., Miall, S., Howes, A., Salter, J., Hills, M.J., Lowe, F.M., A'Hern, R., Nasiri, N., Doody, D., Iqbal, J. & Dowsett, M. (2002). Comparison of the systemic and intratumoral effects of tamoxifen and the aromatase inhibitor vorozole in postmenopausal patients with primary breast cancer. *J Clin Oncol*, **20**, 1026-35.
- Harrela, M., Qiao, Q., Koistinen, R., Tuomilehto, J., Nissinen, A., Seppala, M. & Leinonen, P. (2002). High serum insulin-like growth factor binding protein-1 is associated with increased cardiovascular mortality in elderly men. *Horm Metab Res*, **34**, 144-9.
- Harris, D.M., Besselink, E., Henning, S.M., Go, V.L. & Heber, D. (2005). Phytoestrogens induce differential estrogen receptor alpha- or Beta-mediated responses in transfected breast cancer cells. *Exp Biol Med (Maywood)*, **230**, 558-68.
- Hashim, Y.Z., Eng, M., Gill, C.I., McGlynn, H. & Rowland, I.R. (2005). Components of olive oil and chemoprevention of colorectal cancer. *Nutr Rev*, **63**, 374-86.
- Heald, A.H., Cade, J.E., Cruickshank, J.K., Anderson, S., White, A. & Gibson, J.M. (2003). The influence of dietary intake on the insulin-like growth factor (IGF) system across three ethnic groups: a population-based study. *Public Health Nutr*, **6**, 175-80.
- Hedlund, T.E., Maroni, P.D., Ferucci, P.G., Dayton, R., Barnes, S., Jones, K., Moore, R., Ogden, L.G., Wahala, K., Sackett, H.M. & Gray, K.J. (2005).

- Long-term dietary habits affect soy isoflavone metabolism and accumulation in prostatic fluid in caucasian men. *J Nutr*, **135**, 1400-6.
- Helle, S.I., Anker, G.B., Meadows, K.A., Holly, J.M. & Lonning, P.E. (1998). Alterations in the insulin-like growth factor system during the menstrual cycle in normal women. *Maturitas*, **28**, 259-65.
- Helle, S.I., Holly, J.M., Tally, M., Hall, K., Vander Stappen, J. & Lonning, P.E. (1996). Influence of treatment with tamoxifen and change in tumor burden on the IGF-system in breast cancer patients. *Int J Cancer*, **69**, 335-9.
- Hilakivi-Clarke, L., Onojafe, I., Raygada, M., Cho, E., Skaar, T., Russo, I. & Clarke, R. (1999). Prepubertal exposure to zearalenone or genistein reduces mammary tumorigenesis. *Br J Cancer*, **80**, 1682-8.
- Hirayama, T. (1992). Life-style and cancer: from epidemiological evidence to public behavior change to mortality reduction of target cancers. *J Natl Cancer Inst Monogr*, 65-74.
- Hirohata, T., Shigematsu, T., Nomura, A.M., Nomura, Y., Horie, A. & Hirohata, I. (1985). Occurrence of breast cancer in relation to diet and reproductive history: a case-control study in Fukuoka, Japan. *Natl Cancer Inst Monogr*, **69**, 187-90.
- Hirose, K., Imaeda, N., Tokudome, Y., Goto, C., Wakai, K., Matsuo, K., Ito, H., Toyama, T., Iwata, H., Tokudome, S. & Tajima, K. (2005). Soybean products and reduction of breast cancer risk: a case-control study in Japan. *Br J Cancer*, **93**, 15-22.
- Hirose, K., Tajima, K., Hamajima, N., Inoue, M., Takezaki, T., Kuroishi, T., Yoshida, M. & Tokudome, S. (1995). A large-scale, hospital-based case-control study of risk factors of breast cancer according to menopausal status. *Jpn J Cancer Res*, **86**, 146-54.
- Ho, K.Y., Evans, W.S., Blizzard, R.M., Veldhuis, J.D., Merriam, G.R., Samojlik, E., Furlanetto, R., Rogol, A.D., Kaiser, D.L. & Thorner, M.O. (1987). Effects of sex and age on the 24-hour profile of growth hormone secretion in man: importance of endogenous estradiol concentrations. *J Clin Endocrinol Metab*, **64**, 51-8.
- Hodgson, J.M., Puddey, I.B., Croft, K.D., Mori, T.A., Rivera, J. & Beilin, L.J. (1999). Isoflavonoids do not inhibit in vivo lipid peroxidation in subjects with high-normal blood pressure. *Atherosclerosis*, **145**, 167-72.
- Hoey, L., Rowland, I.R., Lloyd, A.S., Clarke, D.B. & Wiseman, H. (2004). Influence of soya-based infant formula consumption on isoflavone and gut microflora metabolite concentrations in urine and on faecal microflora composition and metabolic activity in infants and children. *Br J Nutr*, **91**, 607-16.
- Hoffman, R. (1995). Potent inhibition of breast cancer cell lines by the isoflavonoid kievitone: comparison with genistein. *Biochem Biophys Res Commun*, **211**, 600-6.
- Holly, J.M. (1991). The physiological role of IGFBP-1. *Acta Endocrinol (Copenh)*, **124 Suppl 2**, 55-62.
- Holmes, M.D., Pollak, M.N., Willett, W.C. & Hankinson, S.E. (2002). Dietary correlates of plasma insulin-like growth factor I and insulin-like growth factor binding protein 3 concentrations. *Cancer Epidemiol Biomarkers Prev*, **11**, 852-61.
- Horn-Ross, P.L., Hoggatt, K.J., West, D.W., Krone, M.R., Stewart, S.L., Anton, H., Bernstei, C.L., Deapen, D., Peel, D., Pinder, R., Reynolds, P., Ross, R.K., Wright, W. & Ziogas, A. (2002). Recent diet and breast cancer risk:

- the California Teachers Study (USA). *Cancer Causes Control*, **13**, 407-15.
- Howell, A., Cuzick, J., Baum, M., Buzdar, A., Dowsett, M., Forbes, J.F., Hocht-Boes, G., Houghton, J., Locker, G.Y. & Tobias, J.S. (2005). Results of the ATAC (Arimidex, Tamoxifen, Alone or in Combination) trial after completion of 5 years' adjuvant treatment for breast cancer. *Lancet*, **365**, 60-2.
- Howes, J.B., Sullivan, D., Lai, N., Nestel, P., Pomeroy, S., West, L., Eden, J.A. & Howes, L.G. (2000). The effects of dietary supplementation with isoflavones from red clover on the lipoprotein profiles of post menopausal women with mild to moderate hypercholesterolaemia. *Atherosclerosis*, **152**, 143-7.
- Hsieh, C.Y., Santell, R.C., Haslam, S.Z. & Helferich, W.G. (1998). Estrogenic effects of genistein on the growth of estrogen receptor-positive human breast cancer (MCF-7) cells in vitro and in vivo. *Cancer Res*, **58**, 3833-8.
- Hsu, J.T., Hung, H.C., Chen, C.J., Hsu, W.L. & Ying, C. (1999). Effects of the dietary phytoestrogen biochanin A on cell growth in the mammary carcinoma cell line MCF-7. *J Nutr Biochem*, **10**, 510-7.
- Hunter, D.J. & Willett, W.C. (1996). Nutrition and breast cancer. *Cancer Causes Control*, **7**, 56-68.
- Hutchins, A.M., Lampe, J.W., Martini, M.C., Campbell, D.R. & Slavin, J.L. (1995). Vegetables, fruits, and legumes: effect on urinary isoflavonoid phytoestrogen and lignan excretion. *J Am Diet Assoc*, **95**, 769-74.
- Huynh, H., Nickerson, T., Pollak, M. & Yang, X. (1996a). Regulation of insulin-like growth factor I receptor expression by the pure antiestrogen ICI 182780. *Clin Cancer Res*, **2**, 2037-42.
- Huynh, H., Yang, X. & Pollak, M. (1996b). Estradiol and antiestrogens regulate a growth inhibitory insulin-like growth factor binding protein 3 autocrine loop in human breast cancer cells. *J Biol Chem*, **271**, 1016-21.
- Huynh, H.T., Tetenes, E., Wallace, L. & Pollak, M. (1993). In vivo inhibition of insulin-like growth factor I gene expression by tamoxifen. *Cancer Res*, **53**, 1727-30.
- Ingram, D., Sanders, K., Kolybaba, M. & Lopez, D. (1997). Case-control study of phyto-oestrogens and breast cancer. *Lancet*, **350**, 990-4.
- Janssen, J.A., Stolk, R.P., Pols, H.A., Grobbee, D.E. & Lamberts, S.W. (1998). Serum total IGF-I, free IGF-I, and IGFB-1 levels in an elderly population: relation to cardiovascular risk factors and disease. *Arterioscler Thromb Vasc Biol*, **18**, 277-82.
- Jenkins, D.J., Kendall, C.W., Vidgen, E., Vuksan, V., Jackson, C.J., Augustin, L.S., Lee, B., Garsetti, M., Agarwal, S., Rao, A.V., Cagampang, G.B. & Fulgoni, V., 3rd. (2000). Effect of soy-based breakfast cereal on blood lipids and oxidized low-density lipoprotein. *Metabolism*, **49**, 1496-500.
- Jing, Y., Nakaya, K. & Han, R. (1993). Differentiation of promyelocytic leukemia cells HL-60 induced by daidzein in vitro and in vivo. *Anticancer Res*, **13**, 1049-54.
- Johansson, H., Baglietto, L., Guerrieri-Gonzaga, A., Bonanni, B., Mariette, F., Macis, D., Serrano, D., Sandri, M.T. & Decensi, A. (2004). Factors associated with circulating levels of insulin-like growth factor-I and insulin-like growth factor binding protein-3 in 740 women at risk for breast cancer. *Breast Cancer Res Treat*, **88**, 63-73.
- Jordan, V.C. (1994). Molecular mechanisms of antiestrogen action in breast cancer. *Breast Cancer Res Treat*, **31**, 41-52.

- Jordan, V.C., Mittal, S., Gosden, B., Koch, R. & Lieberman, M.E. (1985). Structure-activity relationships of estrogens. *Environ Health Perspect*, **61**, 97-110.
- Jousse, C., Bruhat, A., Ferrara, M. & Fafournoux, P. (1998). Physiological concentration of amino acids regulates insulin-like-growth-factor-binding protein 1 expression. *Biochem J*, **334** (Pt 1), 147-53.
- Ju, Y.H., Allred, K.F., Allred, C.D. & Helferich, W.G. (2006a). Genistein stimulates growth of human breast cancer cells in a novel, postmenopausal animal model, with low plasma estradiol concentrations. *Carcinogenesis*, **27**, 1292-9.
- Ju, Y.H., Fultz, J., Allred, K.F., Doerge, D.R. & Helferich, W.G. (2006b). Effects of dietary daidzein and its metabolite, equol, at physiological concentrations on the growth of estrogen-dependent human breast cancer (MCF-7) tumors implanted in ovariectomized athymic mice. *Carcinogenesis*, **27**, 856-63.
- Juul, A. (2003). Serum levels of insulin-like growth factor I and its binding proteins in health and disease. *Growth Horm IGF Res*, **13**, 113-70.
- Juul, A., Scheike, T., Davidsen, M., Gyllenborg, J. & Jorgensen, T. (2002). Low serum insulin-like growth factor I is associated with increased risk of ischemic heart disease: a population-based case-control study. *Circulation*, **106**, 939-44.
- Juul, A., Scheike, T., Pedersen, A.T., Main, K.M., Andersson, A.M., Pedersen, L.M. & Skakkebaek, N.E. (1997). Changes in serum concentrations of growth hormone, insulin, insulin-like growth factor and insulin-like growth factor-binding proteins 1 and 3 and urinary growth hormone excretion during the menstrual cycle. *Hum Reprod*, **12**, 2123-8.
- Kaaks, R., Bellati, C., Venturelli, E., Rinaldi, S., Secreto, G., Biessy, C., Pala, V., Sieri, S. & Berrino, F. (2003). Effects of dietary intervention on IGF-I and IGF-binding proteins, and related alterations in sex steroid metabolism: the Diet and Androgens (DIANA) Randomised Trial. *Eur J Clin Nutr*, **57**, 1079-88.
- Kaaks, R., Berrino, F., Key, T., Rinaldi, S., Dossus, L., Biessy, C., Secreto, G., Amiano, P., Bingham, S., Boeing, H., Bueno de Mesquita, H.B., Chang-Claude, J., Clavel-Chapelon, F., Fournier, A., van Gils, C.H., Gonzalez, C.A., Gurree, A.B., Critselis, E., Khaw, K.T., Krogh, V., Lahmann, P.H., Nagel, G., Olsen, A., Onland-Moret, N.C., Overvad, K., Palli, D., Panico, S., Peeters, P., Quiros, J.R., Roddam, A., Thiebaut, A., Tjonneland, A., Chirlaque, M.D., Trichopoulou, A., Trichopoulos, D., Tumino, R., Vineis, P., Norat, T., Ferrari, P., Slimani, N. & Riboli, E. (2005). Serum sex steroids in premenopausal women and breast cancer risk within the European Prospective Investigation into Cancer and Nutrition (EPIC). *J Natl Cancer Inst*, **97**, 755-65.
- Kaaks, R. & Lukanova, A. (2001). Energy balance and cancer: the role of insulin and insulin-like growth factor-I. *Proc Nutr Soc*, **60**, 91-106.
- Kaaks, R., Lundin, E., Rinaldi, S., Manjer, J., Biessy, C., Soderberg, S., Lenner, P., Janzon, L., Riboli, E., Berglund, G. & Hallmans, G. (2002). Prospective study of IGF-I, IGF-binding proteins, and breast cancer risk, in northern and southern Sweden. *Cancer Causes Control*, **13**, 307-16.
- Kaklamani, V.G., Linos, A., Kaklamani, E., Markaki, I., Koumantaki, Y. & Mantzoros, C.S. (1999). Dietary fat and carbohydrates are independently associated with circulating insulin-like growth factor 1 and insulin-like

- growth factor-binding protein 3 concentrations in healthy adults. *J Clin Oncol*, **17**, 3291-8.
- Kapiotis, S., Hermann, M., Held, I., Seelos, C., Ehringer, H. & Gmeiner, B.M. (1997). Genistein, the dietary-derived angiogenesis inhibitor, prevents LDL oxidation and protects endothelial cells from damage by atherogenic LDL. *Arterioscler Thromb Vasc Biol*, **17**, 2868-74.
- Karey, K.P. & Sirbasku, D.A. (1988). Differential responsiveness of human breast cancer cell lines MCF-7 and T47D to growth factors and 17 beta-estradiol. *Cancer Res*, **48**, 4083-92.
- Karr, S.C., Lampe, J.W., Hutchins, A.M. & Slavin, J.L. (1997). Urinary isoflavonoid excretion in humans is dose dependent at low to moderate levels of soy-protein consumption. *Am J Clin Nutr*, **66**, 46-51.
- Kato, S., Masuhiro, Y., Watanabe, M., Kobayashi, Y., Takeyama, K.I., Endoh, H. & Yanagisawa, J. (2000). Molecular mechanism of a cross-talk between oestrogen and growth factor signalling pathways. *Genes Cells*, **5**, 593-601.
- Katzenellenbogen, B.S., Montano, M.M., Ekena, K., Herman, M.E. & McInerney, E.M. (1997). William L. McGuire Memorial Lecture. Antiestrogens: mechanisms of action and resistance in breast cancer. *Breast Cancer Res Treat*, **44**, 23-38.
- Kawase, T., Orikasa, M., Ogata, S. & Burns, D.M. (1995). Protein tyrosine phosphorylation induced by epidermal growth factor and insulin-like growth factor-I in a rat clonal dental pulp-cell line. *Arch Oral Biol*, **40**, 921-9.
- Keinan-Boker, L., van Der Schouw, Y.T., Grobbee, D.E. & Peeters, P.H. (2004). Dietary phytoestrogens and breast cancer risk. *Am J Clin Nutr*, **79**, 282-8.
- Kelly, G.E., Joannou, G.E., Reeder, A.Y., Nelson, C. & Waring, M.A. (1995). The variable metabolic response to dietary isoflavones in humans. *Proc Soc Exp Biol Med*, **208**, 40-3.
- Kelly, J.J., Rajkovic, I.A., O'Sullivan, A.J., Sernia, C. & Ho, K.K. (1993). Effects of different oral oestrogen formulations on insulin-like growth factor-I, growth hormone and growth hormone binding protein in post-menopausal women. *Clin Endocrinol (Oxf)*, **39**, 561-7.
- Ketelslegers, J.M., Maiter, D., Maes, M., Underwood, L.E. & Thissen, J.P. (1995). Nutritional regulation of insulin-like growth factor-I. *Metabolism*, **44**, 50-7.
- Key, T., Appleby, P., Barnes, I. & Reeves, G. (2002). Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies. *J Natl Cancer Inst*, **94**, 606-16.
- Key, T.J. (1999). Serum oestradiol and breast cancer risk. *Endocr Relat Cancer*, **6**, 175-80.
- Key, T.J., Sharp, G.B., Appleby, P.N., Beral, V., Goodman, M.T., Soda, M. & Mabuchi, K. (1999). Soya foods and breast cancer risk: a prospective study in Hiroshima and Nagasaki, Japan. *Br J Cancer*, **81**, 1248-56.
- Khalil, D.A., Lucas, E.A., Juma, S., Smith, B.J., Payton, M.E. & Arjmandi, B.H. (2002). Soy protein supplementation increases serum insulin-like growth factor-I in young and old men but does not affect markers of bone metabolism. *J Nutr*, **132**, 2605-8.
- Kim, H., Peterson, T.G. & Barnes, S. (1998). Mechanisms of action of the soy isoflavone genistein: emerging role for its effects via transforming growth factor beta signaling pathways. *Am J Clin Nutr*, **68**, 1418S-1425S.

- Kim, H.J., Kwon, S.H., Kim, S.W., Park, D.J., Shin, C.S., Park, K.S., Kim, S.Y., Cho, B.Y. & Lee, H.K. (2001). Diagnostic value of serum IGF-I and IGFBP-3 in growth hormone disorders in adults. *Horm Res*, **56**, 117-23.
- King, R.A. & Bursill, D.B. (1998). Plasma and urinary kinetics of the isoflavones daidzein and genistein after a single soy meal in humans. *Am J Clin Nutr*, **67**, 867-72.
- King, R.A., Mano, M.M. & Head, R.J. (1998). Assessment of isoflavonoid concentrations in Australian bovine milk samples. *J Dairy Res*, **65**, 479-89.
- Knight, D.C., Howes, J.B. & Eden, J.A. (1999). The effect of Promensil, an isoflavone extract, on menopausal symptoms. *Climacteric*, **2**, 79-84.
- Kostelac, D., Rechkemmer, G. & Briviba, K. (2003). Phytoestrogens modulate binding response of estrogen receptors alpha and beta to the estrogen response element. *J Agric Food Chem*, **51**, 7632-5.
- Krajcik, R.A., Borofsky, N.D., Massardo, S. & Orentreich, N. (2002). Insulin-like growth factor I (IGF-I), IGF-binding proteins, and breast cancer. *Cancer Epidemiol Biomarkers Prev*, **11**, 1566-73.
- Kuiper, G.G., Carlsson, B., Grandien, K., Enmark, E., Haggblad, J., Nilsson, S. & Gustafsson, J.A. (1997). Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology*, **138**, 863-70.
- Kuiper, G.G., Enmark, E., Peltö-Huikko, M., Nilsson, S. & Gustafsson, J.A. (1996). Cloning of a novel receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci U S A*, **93**, 5925-30.
- Kuiper, G.G. & Gustafsson, J.A. (1997). The novel estrogen receptor-beta subtype: potential role in the cell- and promoter-specific actions of estrogens and anti-estrogens. *FEBS Lett*, **410**, 87-90.
- Kuiper, G.G., Lemmen, J.G., Carlsson, B., Corton, J.C., Safe, S.H., van der Saag, P.T., van der Burg, B. & Gustafsson, J.A. (1998). Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology*, **139**, 4252-63.
- Kumar, N.B., Cantor, A., Allen, K., Riccardi, D. & Cox, C.E. (2002). The specific role of isoflavones on estrogen metabolism in premenopausal women. *Cancer*, **94**, 1166-74.
- Kurzer, M.S. (2002). Hormonal effects of soy in premenopausal women and men. *J Nutr*, **132**, 570S-573S.
- Laban, C., Bustin, S.A. & Jenkins, P.J. (2003). The GH-IGF-I axis and breast cancer. *Trends Endocrinol Metab*, **14**, 28-34.
- Lahti, E.I., Knip, M. & Laatikainen, T.J. (1994). Plasma insulin-like growth factor I and its binding proteins 1 and 3 in postmenopausal patients with breast cancer receiving long term tamoxifen. *Cancer*, **74**, 618-24.
- Lamartiniere, C.A., Cotroneo, M.S., Fritz, W.A., Wang, J., Mentor-Marcel, R. & Elgavish, A. (2002a). Genistein chemoprevention: timing and mechanisms of action in murine mammary and prostate. *J Nutr*, **132**, 552S-558S.
- Lamartiniere, C.A., Moore, J., Holland, M. & Barnes, S. (1995). Neonatal genistein chemoprevents mammary cancer. *Proc Soc Exp Biol Med*, **208**, 120-3.
- Lamartiniere, C.A., Wang, J., Smith-Johnson, M. & Eltoum, I.E. (2002b). Daidzein: bioavailability, potential for reproductive toxicity, and breast cancer chemoprevention in female rats. *Toxicol Sci*, **65**, 228-38.

- Lampe, J.W., Gustafson, D.R., Hutchins, A.M., Martini, M.C., Li, S., Wahala, K., Grandits, G.A., Potter, J.D. & Slavin, J.L. (1999). Urinary isoflavonoid and lignan excretion on a Western diet: relation to soy, vegetable, and fruit intake. *Cancer Epidemiol Biomarkers Prev*, **8**, 699-707.
- Lampe, J.W., Karr, S.C., Hutchins, A.M. & Slavin, J.L. (1998). Urinary equol excretion with a soy challenge: influence of habitual diet. *Proc Soc Exp Biol Med*, **217**, 335-9.
- Lampe, J.W., Skor, H.E., Li, S., Wahala, K., Howald, W.N. & Chen, C. (2001). Wheat bran and soy protein feeding do not alter urinary excretion of the isoflavan equol in premenopausal women. *J Nutr*, **131**, 740-4.
- Larsson, S.C., Wolk, K., Brismar, K. & Wolk, A. (2005). Association of diet with serum insulin-like growth factor I in middle-aged and elderly men. *Am J Clin Nutr*, **81**, 1163-7.
- Laughlin, G.A., Barrett-Connor, E., Criqui, M.H. & Kritz-Silverstein, D. (2004). The prospective association of serum insulin-like growth factor I (IGF-I) and IGF-binding protein-1 levels with all cause and cardiovascular disease mortality in older adults: the Rancho Bernardo Study. *J Clin Endocrinol Metab*, **89**, 114-20.
- Lavigne, J.A., Wimbrow, H.H., Clevidence, B.A., Albert, P.S., Reichman, M.E., Campbell, W.S., Barrett, J.C., Hursting, S.D., Judd, J.T. & Taylor, P.R. (2004). Effects of alcohol and menstrual cycle on insulin-like growth factor-I and insulin-like growth factor binding protein-3. *Cancer Epidemiol Biomarkers Prev*, **13**, 2264-7.
- Le Bail, J.C., Varnat, F., Nicolas, J.C. & Habrioux, G. (1998). Estrogenic and antiproliferative activities on MCF-7 human breast cancer cells by flavonoids. *Cancer Lett*, **130**, 209-16.
- Lee, A.V., Jackson, J.G., Gooch, J.L., Hilsenbeck, S.G., Coronado-Heinsohn, E., Osborne, C.K. & Yee, D. (1999). Enhancement of insulin-like growth factor signaling in human breast cancer: estrogen regulation of insulin receptor substrate-1 expression in vitro and in vivo. *Mol Endocrinol*, **13**, 787-96.
- Lee, A.V., Weng, C.N., Jackson, J.G. & Yee, D. (1997a). Activation of estrogen receptor-mediated gene transcription by IGF-I in human breast cancer cells. *J Endocrinol*, **152**, 39-47.
- Lee, H.P., Day, N.E. & Shanmugaratnam, K. (1988). Trends in cancer incidence in Singapore 1968-1982. *IARC Sci.Publ.*, 1-161.
- Lee, H.P., Gourley, L., Duffy, S.W., Esteve, J., Lee, J. & Day, N.E. (1991). Dietary effects on breast-cancer risk in Singapore. *Lancet*, **337**, 1197-200.
- Lee, H.P., Gourley, L., Duffy, S.W., Esteve, J., Lee, J. & Day, N.E. (1992). Risk factors for breast cancer by age and menopausal status: a case-control study in Singapore. *Cancer Causes Control*, **3**, 313-22.
- Lee, P.D., Giudice, L.C., Conover, C.A. & Powell, D.R. (1997b). Insulin-like growth factor binding protein-1: recent findings and new directions. *Proc Soc Exp Biol Med*, **216**, 319-57.
- Li, B.D., Khosravi, M.J., Berkel, H.J., Diamandi, A., Dayton, M.A., Smith, M. & Yu, H. (2001). Free insulin-like growth factor-I and breast cancer risk. *Int J Cancer*, **91**, 736-9.
- Li, D., Yee, J.A., McGuire, M.H., Murphy, P.A. & Yan, L. (1999). Soybean isoflavones reduce experimental metastasis in mice. *J Nutr*, **129**, 1075-8.

- Li, Y. & Sarkar, F.H. (2002). Inhibition of nuclear factor kappaB activation in PC3 cells by genistein is mediated via Akt signaling pathway. *Clin Cancer Res*, **8**, 2369-77.
- Lichtenstein, A.H. (2001). Got soy? *Am J Clin Nutr*, **73**, 667-8.
- Lien, E.A., Solheim, E., Lea, O.A., Lundgren, S., Kvinnsland, S. & Ueland, P.M. (1989). Distribution of 4-hydroxy-N-desmethyltamoxifen and other tamoxifen metabolites in human biological fluids during tamoxifen treatment. *Cancer Res*, **49**, 2175-83.
- Linassier, C., Pierre, M., Le Pecq, J.B. & Pierre, J. (1990). Mechanisms of action in NIH-3T3 cells of genistein, an inhibitor of EGF receptor tyrosine kinase activity. *Biochem Pharmacol*, **39**, 187-93.
- Lissin, L.W. & Cooke, J.P. (2000). Phytoestrogens and cardiovascular health. *J Am Coll Cardiol*, **35**, 1403-10.
- Liu, D.X. & Wang, C.C. (1997). [Antigen strategy: a new way for gene therapy]. *Sheng Li Ke Xue Jin Zhan*, **28**, 214-8.
- Lonning, P.E., Hall, K., Aakvaag, A. & Lien, E.A. (1992). Influence of tamoxifen on plasma levels of insulin-like growth factor I and insulin-like growth factor binding protein I in breast cancer patients. *Cancer Res*, **52**, 4719-23.
- Lonning, P.E., Helle, S.I., Johannessen, D.C., Adlercreutz, H., Lien, E.A., Tally, M., Ekse, D., Fotsis, T., Anker, G.B. & Hall, K. (1995). Relations between sex hormones, sex hormone binding globulin, insulin-like growth factor-I and insulin-like growth factor binding protein-1 in post-menopausal breast cancer patients. *Clin Endocrinol (Oxf)*, **42**, 23-30.
- Low, Y.L., Taylor, J.I., Grace, P.B., Dowsett, M., Scollen, S., Dunning, A.M., Mulligan, A.A., Welch, A.A., Luben, R.N., Khaw, K.T., Day, N.E., Wareham, N.J. & Bingham, S.A. (2005). Phytoestrogen exposure correlation with plasma estradiol in postmenopausal women in European Prospective Investigation of Cancer and Nutrition-Norfolk may involve diet-gene interactions. *Cancer Epidemiol Biomarkers Prev*, **14**, 213-20.
- Lu, L.J. & Anderson, K.E. (1998). Sex and long-term soy diets affect the metabolism and excretion of soy isoflavones in humans. *Am J Clin Nutr*, **68**, 1500S-1504S.
- Lu, L.J., Anderson, K.E., Grady, J.J., Kohen, F. & Nagamani, M. (2000a). Decreased ovarian hormones during a soya diet: implications for breast cancer prevention. *Cancer Res*, **60**, 4112-21.
- Lu, L.J., Anderson, K.E., Grady, J.J. & Nagamani, M. (1996a). Effects of soya consumption for one month on steroid hormones in premenopausal women: implications for breast cancer risk reduction. *Cancer Epidemiol Biomarkers Prev*, **5**, 63-70.
- Lu, L.J., Anderson, K.E., Grady, J.J. & Nagamani, M. (2001). Effects of an isoflavone-free soy diet on ovarian hormones in premenopausal women. *J Clin Endocrinol Metab*, **86**, 3045-52.
- Lu, L.J., Cree, M., Josyula, S., Nagamani, M., Grady, J.J. & Anderson, K.E. (2000b). Increased urinary excretion of 2-hydroxyestrone but not 16alpha-hydroxyestrone in premenopausal women during a soya diet containing isoflavones. *Cancer Res*, **60**, 1299-305.
- Lu, L.J., Lin, S.N., Grady, J.J., Nagamani, M. & Anderson, K.E. (1996b). Altered kinetics and extent of urinary daidzein and genistein excretion in women during chronic soya exposure. *Nutr Cancer*, **26**, 289-302.

- Lund, T.D., Munson, D.J., Haldy, M.E., Setchell, K.D., Lephart, E.D. & Handa, R.J. (2004). Equol is a novel anti-androgen that inhibits prostate growth and hormone feedback. *Biol Reprod*, **70**, 1188-95.
- Lydeking-Olsen, E., Beck-Jensen, J.E., Setchell, K.D. & Holm-Jensen, T. (2004). Soymilk or progesterone for prevention of bone loss—a 2 year randomized, placebo-controlled trial. *Eur J Nutr*, **43**, 246-57.
- Ma, J., Giovannucci, E., Pollak, M., Chan, J.M., Gaziano, J.M., Willett, W. & Stampfer, M.J. (2001). Milk intake, circulating levels of insulin-like growth factor-I, and risk of colorectal cancer in men. *J Natl Cancer Inst*, **93**, 1330-6.
- Magee, P.J., Raschke, M., Steiner, C., Duffin, J.G., Pool-Zobel, B.L., Jokela, T., Wahala, K. & Rowland, I.R. (2006). Equol: a comparison of the effects of the racemic compound with that of the purified S-enantiomer on the growth, invasion, and DNA integrity of breast and prostate cells in vitro. *Nutr Cancer*, **54**, 232-42.
- Magee, P.J. & Rowland, I.R. (2004). Phyto-oestrogens, their mechanism of action: current evidence for a role in breast and prostate cancer. *Br J Nutr*, **91**, 513-31.
- Maggiolini, M., Bonfiglio, D., Marsico, S., Panno, M.L., Cenni, B., Picard, D. & Ando, S. (2001). Estrogen receptor alpha mediates the proliferative but not the cytotoxic dose-dependent effects of two major phytoestrogens on human breast cancer cells. *Mol Pharmacol*, **60**, 595-602.
- Markiewicz, L., Garey, J., Adlercreutz, H. & Gurdip, E. (1993). In vitro bioassays of non-steroidal phytoestrogens. *J Steroid Biochem Mol Biol*, **45**, 399-405.
- Martin, M.B. & Stoica, A. (2002). Insulin-like growth factor-I and estrogen interactions in breast cancer. *J Nutr*, **132**, 3799S-3801S.
- Martin, M.E., Haourigui, M., Pelissero, C., Benassayag, C. & Nunez, E.A. (1996). Interactions between phytoestrogens and human sex steroid binding protein. *Life Sci*, **58**, 429-36.
- Martini, M.C., Dancisak, B.B., Haggans, C.J., Thomas, W. & Slavin, J.L. (1999). Effects of soy intake on sex hormone metabolism in premenopausal women. *Nutr Cancer*, **34**, 133-9.
- Martino, S., Costantino, J., McNabb, M., Mershon, J., Bryant, K., Powles, T. & Secrest, R.J. (2004). The role of selective estrogen receptor modulators in the prevention of breast cancer: comparison of the clinical trials. *Oncologist*, **9**, 116-25.
- Maskarinec, G., Franke, A.A., Williams, A.E., Hebshi, S., Oshiro, C., Murphy, S. & Stanczyk, F.Z. (2004). Effects of a 2-year randomized soy intervention on sex hormone levels in premenopausal women. *Cancer Epidemiol Biomarkers Prev*, **13**, 1736-44.
- Maskarinec, G., Takata, Y. & Kaaks, R. (2005a). The relation between nutritional factors and insulin-like growth factor-I in premenopausal women of different ethnicity. *Eur J Nutr*, **44**, 105-13.
- Maskarinec, G., Takata, Y., Murphy, S.P., Franke, A.A. & Kaaks, R. (2005b). Insulin-like growth factor-1 and binding protein-3 in a 2-year soya intervention among premenopausal women. *Br J Nutr*, **94**, 362-7.
- Maskarinec, G., Williams, A.E., Inouye, J.S., Stanczyk, F.Z. & Franke, A.A. (2002). A randomized isoflavone intervention among premenopausal women. *Cancer Epidemiol Biomarkers Prev*, **11**, 195-201.
- Massa, G., Igout, A., Rombauts, L., Frankenne, F. & Vanderschueren-Lodeweyckx, M. (1993). Effect of oestrogen status on serum levels of

- growth hormone-binding protein and insulin-like growth factor-I in non-pregnant and pregnant women. *Clin Endocrinol (Oxf)*, **39**, 569-75.
- Matsukawa, Y., Marui, N., Sakai, T., Satomi, Y., Yoshida, M., Matsumoto, K., Nishino, H. & Aoike, A. (1993). Genistein arrests cell cycle progression at G2-M. *Cancer Res*, **53**, 1328-31.
- Matsumura, A., Ghosh, A., Pope, G.S. & Darbre, P.D. (2005). Comparative study of oestrogenic properties of eight phytoestrogens in MCF7 human breast cancer cells. *J Steroid Biochem Mol Biol*, **94**, 431-43.
- Matthews, J. & Gustafsson, J.A. (2003). Estrogen signaling: a subtle balance between ER alpha and ER beta. *Mol Interv*, **3**, 281-92.
- McCance, R.A., Widdowson, E.M., Paul, A.A., Southgate, D.A.T., Medical Research, C., Great Britain. Ministry of Agriculture, F. & Food. Medical Research, C. (1978). *McCance and Widdowson's The composition of foods*. Special report series ; no. 297. H.M.S.O.: London.
- McCann, S.E., Muti, P., Vito, D., Edge, S.B., Trevisan, M. & Freudenheim, J.L. (2004). Dietary lignan intakes and risk of pre- and postmenopausal breast cancer. *Int J Cancer*, **111**, 440-3.
- McCarty, M.F. (2006). Isoflavones made simple - genistein's agonist activity for the beta-type estrogen receptor mediates their health benefits. *Med Hypotheses*, **66**, 1093-114.
- McDowell, I.F., McEneny, J. & Trimble, E.R. (1995). A rapid method for measurement of the susceptibility to oxidation of low-density lipoprotein. *Ann Clin Biochem*, **32** (Pt 2), 167-74.
- McKeown, N.M., Day, N.E., Welch, A.A., Runswick, S.A., Luben, R.N., Mulligan, A.A., McTaggart, A. & Bingham, S.A. (2001). Use of biological markers to validate self-reported dietary intake in a random sample of the European Prospective Investigation into Cancer United Kingdom Norfolk cohort. *Am J Clin Nutr*, **74**, 188-96.
- McMichael-Phillips, D.F., Harding, C., Morton, M., Roberts, S.A., Howell, A., Potten, C.S. & Bundred, N.J. (1998). Effects of soy-protein supplementation on epithelial proliferation in the histologically normal human breast. *Am J Clin Nutr*, **68**, 1431S-1435S.
- McPherson, K., Steel, C.M. & Dixon, J.M. (2000). ABC of breast diseases. Breast cancer-epidemiology, risk factors, and genetics. *Bmj*, **321**, 624-8.
- Mendez, P., Azcoitia, I. & Garcia-Segura, L.M. (2003). Estrogen receptor alpha forms estrogen-dependent multimolecular complexes with insulin-like growth factor receptor and phosphatidylinositol 3-kinase in the adult rat brain. *Brain Res Mol Brain Res*, **112**, 170-6.
- Merz-Demlow, B.E., Duncan, A.M., Wangen, K.E., Xu, X., Carr, T.P., Phipps, W.R. & Kurzer, M.S. (2000). Soy isoflavones improve plasma lipids in normocholesterolemic, premenopausal women. *Am J Clin Nutr*, **71**, 1462-9.
- Messina, M., Barnes, S. & Setchell, K.D. (1997). Phyto-oestrogens and breast cancer. *Lancet*, **350**, 971-2.
- Messina, M., McCaskill-Stevens, W. & Lampe, J.W. (2006). Addressing the soy and breast cancer relationship: review, commentary, and workshop proceedings. *J Natl Cancer Inst*, **98**, 1275-84.
- Messina, M.J. (1999). Legumes and soybeans: overview of their nutritional profiles and health effects. *Am J Clin Nutr*, **70**, 439S-450S.
- Meyer, B.J., Larkin, T.A., Owen, A.J., Astheimer, L.B., Tapsell, L.C. & Howe, P.R. (2004). Limited lipid-lowering effects of regular consumption of whole soybean foods. *Ann Nutr Metab*, **48**, 67-78.

- Middleton, E., Jr., Kandaswami, C. & Theoharides, T.C. (2000). The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease, and cancer. *Pharmacol Rev*, **52**, 673-751.
- Migliaccio, A., Di Domenico, M., Castoria, G., de Falco, A., Bontempo, P., Nola, E. & Auricchio, F. (1996). Tyrosine kinase/p21ras/MAP-kinase pathway activation by estradiol-receptor complex in MCF-7 cells. *Embo J*, **15**, 1292-300.
- Miller, A.B., Berrino, F., Hill, M., Pietinen, P., Riboli, E. & Wahrendorf, J. (1994). Diet in the aetiology of cancer: a review. *Eur J Cancer*, **30A**, 207-20; discussion 220-8.
- Mitchell, J.H. & Collins, A.R. (1999). Effects of a soy milk supplement on plasma cholesterol levels and oxidative DNA damage in men--a pilot study. *Eur J Nutr*, **38**, 143-8.
- Monti, E. & Sinha, B.K. (1994). Antiproliferative effect of genistein and adriamycin against estrogen-dependent and -independent human breast carcinoma cell lines. *Anticancer Res*, **14**, 1221-6.
- Morito, K., Hirose, T., Kinjo, J., Hirakawa, T., Okawa, M., Nohara, T., Ogawa, S., Inoue, S., Muramatsu, M. & Masamune, Y. (2001). Interaction of phytoestrogens with estrogen receptors alpha and beta. *Biol Pharm Bull*, **24**, 351-6.
- Morton, M.S., Arisaka, O., Miyake, N., Morgan, L.D. & Evans, B.A. (2002). Phytoestrogen concentrations in serum from Japanese men and women over forty years of age. *J Nutr*, **132**, 3168-71.
- Morton, M.S., Chan, P.S., Cheng, C., Blacklock, N., Matos-Ferreira, A., Abranches-Monteiro, L., Correia, R., Lloyd, S. & Griffiths, K. (1997). Lignans and isoflavonoids in plasma and prostatic fluid in men: samples from Portugal, Hong Kong, and the United Kingdom. *Prostate*, **32**, 122-8.
- Morton, M.S., Wilcox, G., Wahlqvist, M.L. & Griffiths, K. (1994). Determination of lignans and isoflavonoids in human female plasma following dietary supplementation. *J Endocrinol*, **142**, 251-9.
- Mouridsen, H., Gershanovich, M., Sun, Y., Perez-Carrion, R., Boni, C., Monnier, A., Apffelstaedt, J., Smith, R., Sleenboom, H.P., Janicke, F., Pluzanska, A., Dank, M., Becquart, D., Bapsy, P.P., Salminen, E., Snyder, R., Lassus, M., Verbeek, J.A., Staffler, B., Chaudri-Ross, H.A. & Dugan, M. (2001). Superior efficacy of letrozole versus tamoxifen as first-line therapy for postmenopausal women with advanced breast cancer: results of a phase III study of the International Letrozole Breast Cancer Group. *J Clin Oncol*, **19**, 2596-606.
- Mucci, L.A., Tamimi, R., Lagiou, P., Trichopoulou, A., Benetou, V., Spanos, E. & Trichopoulos, D. (2001). Are dietary influences on the risk of prostate cancer mediated through the insulin-like growth factor system? *BJU Int*, **87**, 814-20.
- Mulligan, A.A., Welch, A.A., McTaggart, A.A., Bhaniani, A. & Bingham, S.A. (2006). Intakes and sources of soya foods and isoflavones in a UK population cohort study (EPIC-Norfolk). *Eur J Clin Nutr*.
- Murkies, A., Dalais, F.S., Briganti, E.M., Burger, H.G., Healy, D.L., Wahlqvist, M.L. & Davis, S.R. (2000). Phytoestrogens and breast cancer in postmenopausal women: a case control study. *Menopause*, **7**, 289-96.
- Murkies, A.L., Lombard, C., Strauss, B.J., Wilcox, G., Burger, H.G. & Morton, M.S. (1995). Dietary flour supplementation decreases post-menopausal hot flushes: effect of soy and wheat. *Maturitas*, **21**, 189-95.

- Murphy, P.A., Song, T., Buseman, G., Barua, K., Beecher, G.R., Trainer, D. & Holden, J. (1999). Isoflavones in retail and institutional soy foods. *J Agric Food Chem*, **47**, 2697-704.
- Muthyala, R.S., Ju, Y.H., Sheng, S., Williams, L.D., Doerge, D.R., Katzenellenbogen, B.S., Helferich, W.G. & Katzenellenbogen, J.A. (2004). Equol, a natural estrogenic metabolite from soy isoflavones: convenient preparation and resolution of R- and S-equols and their differing binding and biological activity through estrogen receptors alpha and beta. *Bioorg Med Chem*, **12**, 1559-67.
- Muti, P., Quattrin, T., Grant, B.J., Krogh, V., Micheli, A., Schunemann, H.J., Ram, M., Freudenheim, J.L., Sieri, S., Trevisan, M. & Berrino, F. (2002). Fasting glucose is a risk factor for breast cancer: a prospective study. *Cancer Epidemiol Biomarkers Prev*, **11**, 1361-8.
- Nabholtz, J.M., Buzdar, A., Pollak, M., Harwin, W., Burton, G., Mangalik, A., Steinberg, M., Webster, A. & von Euler, M. (2000). Anastrozole is superior to tamoxifen as first-line therapy for advanced breast cancer in postmenopausal women: results of a North American multicenter randomized trial. Arimidex Study Group. *J Clin Oncol*, **18**, 3758-67.
- Naciff, J.M., Jump, M.L., Torontali, S.M., Carr, G.J., Tiesman, J.P., Overmann, G.J. & Daston, G.P. (2002). Gene expression profile induced by 17alpha-ethynyl estradiol, bisphenol A, and genistein in the developing female reproductive system of the rat. *Toxicol Sci*, **68**, 184-99.
- Nagata, C., Kabuto, M., Kurisu, Y. & Shimizu, H. (1997). Decreased serum estradiol concentration associated with high dietary intake of soy products in premenopausal Japanese women. *Nutr Cancer*, **29**, 228-33.
- Nagata, C., Shimizu, H., Takami, R., Hayashi, M., Takeda, N. & Yasuda, K. (2003). Dietary soy and fats in relation to serum insulin-like growth factor-1 and insulin-like growth factor-binding protein-3 levels in premenopausal Japanese women. *Nutr Cancer*, **45**, 185-9.
- Nagata, C., Takatsuka, N., Inaba, S., Kawakami, N. & Shimizu, H. (1998). Effect of soymilk consumption on serum estrogen concentrations in premenopausal Japanese women. *J Natl Cancer Inst*, **90**, 1830-5.
- Nagel, S.C., vom Saal, F.S. & Welshons, W.V. (1998). The effective free fraction of estradiol and xenoestrogens in human serum measured by whole cell uptake assays: physiology of delivery modifies estrogenic activity. *Proc Soc Exp Biol Med*, **217**, 300-9.
- Nestel, P.J., Pomeroy, S., Kay, S., Komesaroff, P., Behrsing, J., Cameron, J.D. & West, L. (1999). Isoflavones from red clover improve systemic arterial compliance but not plasma lipids in menopausal women. *J Clin Endocrinol Metab*, **84**, 895-8.
- Newton, C.J., Buric, R., Trapp, T., Brockmeier, S., Pagotto, U. & Stalla, G.K. (1994). The unliganded estrogen receptor (ER) transduces growth factor signals. *J Steroid Biochem Mol Biol*, **48**, 481-6.
- Nicholls, J., Lasley, B.L., Nakajima, S.T., Setchell, K.D. & Schneeman, B.O. (2002). Effects of soy consumption on gonadotropin secretion and acute pituitary responses to gonadotropin-releasing hormone in women. *J Nutr*, **132**, 708-14.
- Nilsson, A., Hill, J.L. & Davids, H.L. (1967). An in vitro study of formononetin and biochanin A metabolism in rumen fluid from sheep. *Biochim Biophys Acta*, **148**, 92-8.
- Nomura, A., Henderson, B.E. & Lee, J. (1978). Breast cancer and diet among the Japanese in Hawaii. *Am J Clin Nutr*, **31**, 2020-5.

- Norat, T., Dossus, L., Rinaldi, S., Overvad, K., Gronbaek, H., Tjonneland, A., Olsen, A., Clavel-Chapelon, F., Boutron-Ruault, M.C., Boeing, H., Lahmann, P.H., Linseisen, J., Nagel, G., Trichopoulou, A., Trichopoulos, D., Kalapothaki, V., Sieri, S., Palli, D., Panico, S., Tumino, R., Sacerdote, C., Bueno-de-Mesquita, H.B., Peeters, P.H., van Gils, C.H., Agudo, A., Amiano, P., Ardanoz, E., Martinez, C., Quiros, R., Tormo, M.J., Bingham, S., Key, T.J., Allen, N.E., Ferrari, P., Slimani, N., Riboli, E. & Kaaks, R. (2007). Diet, serum insulin-like growth factor-I and IGF-binding protein-3 in European women. *Eur J Clin Nutr*, **61**, 91-8.
- O'Brian, C.A., Liskamp, R.M., Solomon, D.H. & Weinstein, I.B. (1985). Inhibition of protein kinase C by tamoxifen. *Cancer Res*, **45**, 2462-5.
- Olsson, H., Landin-Olsson, M. & Gullberg, B. (1983). Retrospective assessment of menstrual cycle length in patients with breast cancer, in patients with benign breast disease, and in women without breast disease. *J Natl Cancer Inst*, **70**, 17-20.
- ONS. (31 October 2003). National Statistics Online - Breast Cancer. <http://www.statistics.gov.uk/cc/nugget.asp?id=575>. Office for National Statistics, UK.
- Ozasa, K., Nakao, M., Watanabe, Y., Hayashi, K., Miki, T., Mikami, K., Mori, M., Sakauchi, F., Washio, M., Ito, Y., Suzuki, K., Kubo, T., Wakai, K. & Tamakoshi, A. (2005). Association of serum phytoestrogen concentration and dietary habits in a sample set of the JACC Study. *J Epidemiol*, **15 Suppl 2**, S196-202.
- Paech, K., Webb, P., Kuiper, G.G., Nilsson, S., Gustafsson, J., Kushner, P.J. & Scanlan, T.S. (1997). Differential ligand activation of estrogen receptors ERalpha and ERbeta at AP1 sites. *Science*, **277**, 1508-10.
- Pagliacci, M.C., Smacchia, M., Migliorati, G., Grignani, F., Riccardi, C. & Nicoletti, I. (1994). Growth-inhibitory effects of the natural phyto-oestrogen genistein in MCF-7 human breast cancer cells. *Eur J Cancer*, **30A**, 1675-82.
- Paik, S. (1992). Expression of IGF-I and IGF-II mRNA in breast tissue. *Breast Cancer Res Treat*, **22**, 31-8.
- Palmieri, C., Cheng, G.J., Saji, S., Zelada-Hedman, M., Warri, A., Weihua, Z., Van Noorden, S., Wahlstrom, T., Coombes, R.C., Warner, M. & Gustafsson, J.A. (2002). Estrogen receptor beta in breast cancer. *Endocr Relat Cancer*, **9**, 1-13.
- Panno, M.L., Salerno, M., Pezzi, V., Sisci, D., Maggiolini, M., Mauro, L., Morrone, E.G. & Ando, S. (1996). Effect of oestradiol and insulin on the proliferative pattern and on oestrogen and progesterone receptor contents in MCF-7 cells. *J Cancer Res Clin Oncol*, **122**, 745-9.
- Park, B.W., Kim, K.S., Heo, M.K., Ko, S.S., Hong, S.W., Yang, W.I., Kim, J.H., Kim, G.E. & Lee, K.S. (2003). Expression of estrogen receptor-beta in normal mammary and tumor tissues: is it protective in breast carcinogenesis? *Breast Cancer Res Treat*, **80**, 79-85.
- Paruthiyil, S., Parmar, H., Kerekatte, V., Cunha, G.R., Firestone, G.L. & Leitman, D.C. (2004). Estrogen receptor beta inhibits human breast cancer cell proliferation and tumor formation by causing a G2 cell cycle arrest. *Cancer Res*, **64**, 423-8.
- Peeters, P.H., Keinan-Boker, L., van der Schouw, Y.T. & Grobbee, D.E. (2003). Phytoestrogens and breast cancer risk. Review of the epidemiological evidence. *Breast Cancer Res Treat*, **77**, 171-83.

- Persky, V.W., Turyk, M.E., Wang, L., Freels, S., Chatterton, R., Jr., Barnes, S., Erdman, J., Jr., Sepkovic, D.W., Bradlow, H.L. & Potter, S. (2002). Effect of soy protein on endogenous hormones in postmenopausal women. *Am J Clin Nutr*, **75**, 145-53.
- Peterson, G. & Barnes, S. (1991). Genistein inhibition of the growth of human breast cancer cells: independence from estrogen receptors and the multi-drug resistance gene. *Biochem Biophys Res Commun*, **179**, 661-7.
- Peterson, J., Lagiou, P., Samoli, E., Lagiou, A., Katsouyanni, K., La Vecchia, C., Dwyer, J. & Trichopoulos, D. (2003). Flavonoid intake and breast cancer risk: a case-control study in Greece. *Br J Cancer*, **89**, 1255-9.
- Peterson, T.G., Coward, L., Kirk, M., Falany, C.N. & Barnes, S. (1996). The role of metabolism in mammary epithelial cell growth inhibition by the isoflavones genistein and biochanin A. *Carcinogenesis*, **17**, 1861-9.
- Petrakis, N.L., Barnes, S., King, E.B., Lowenstein, J., Wiencke, J., Lee, M.M., Milke, R., Kirk, M. & Coward, L. (1996). Stimulatory influence of soy protein isolate on breast secretion in pre- and postmenopausal women. *Cancer Epidemiol Biomarkers Prev*, **5**, 785-94.
- Peyrat, J.P., Bonnetterre, J., Hecquet, B., Vennin, P., Louchez, M.M., Fournier, C., Lefebvre, J. & Demaille, A. (1993). Plasma insulin-like growth factor-1 (IGF-1) concentrations in human breast cancer. *Eur J Cancer*, **29A**, 492-7.
- Pietinen, P., Stumpf, K., Mannisto, S., Kataja, V., Uusitupa, M. & Adlercreutz, H. (2001). Serum enterolactone and risk of breast cancer: a case-control study in eastern Finland. *Cancer Epidemiol Biomarkers Prev*, **10**, 339-44.
- Pike, M.C., Spicer, D.V., Dahmouch, L. & Press, M.F. (1993). Estrogens, progestogens, normal breast cell proliferation, and breast cancer risk. *Epidemiol Rev*, **15**, 17-35.
- Piller, R., Chang-Claude, J. & Linseisen, J. (2006). Plasma enterolactone and genistein and the risk of premenopausal breast cancer. *Eur J Cancer Prev*, **15**, 225-32.
- Pino, A.M., Valladares, L.E., Palma, M.A., Mancilla, A.M., Yanez, M. & Albala, C. (2000). Dietary isoflavones affect sex hormone-binding globulin levels in postmenopausal women. *J Clin Endocrinol Metab*, **85**, 2797-800.
- Pollak, M., Costantino, J., Polychronakos, C., Blauer, S.A., Guyda, H., Redmond, C., Fisher, B. & Margolese, R. (1990). Effect of tamoxifen on serum insulinlike growth factor I levels in stage I breast cancer patients. *J Natl Cancer Inst*, **82**, 1693-7.
- Pool-Zobel, B.L., Adlercreutz, H., Glei, M., Liegibel, U.M., Sittlington, J., Rowland, I., Wahala, K. & Rechkemmer, G. (2000). Isoflavonoids and lignans have different potentials to modulate oxidative genetic damage in human colon cells. *Carcinogenesis*, **21**, 1247-52.
- Potter, S.M. (1995). Overview of proposed mechanisms for the hypocholesterolemic effect of soy. *J Nutr*, **125**, 606S-611S.
- Power, K.A. & Thompson, L.U. (2003). Ligand-induced regulation of ERalpha and ERbeta is indicative of human breast cancer cell proliferation. *Breast Cancer Res Treat*, **81**, 209-21.
- Powles, T., Eeles, R., Ashley, S., Easton, D., Chang, J., Dowsett, M., Tidy, A., Viggers, J. & Davey, J. (1998). Interim analysis of the incidence of breast cancer in the Royal Marsden Hospital tamoxifen randomised chemoprevention trial. *Lancet*, **352**, 98-101.
- Probst-Hensch, N.M., Wang, H., Goh, V.H., Seow, A., Lee, H.P. & Yu, M.C. (2003). Determinants of circulating insulin-like growth factor I and insulin-

- like growth factor binding protein 3 concentrations in a cohort of Singapore men and women. *Cancer Epidemiol Biomarkers Prev*, **12**, 739-46.
- Pumford, S.L., Morton, M.M., Turkes, A. & Griffiths, K. (2002). Determination of the isoflavonoids genistein and daidzein in biological samples by gas chromatography-mass spectrometry. *Ann Clin Biochem*, **39**, 281-92.
- Qin, L.Q., Xu, J.Y., Wang, P.Y. & Hoshi, K. (2006). Soyfood intake in the prevention of breast cancer risk in women: a meta-analysis of observational epidemiological studies. *J Nutr Sci Vitaminol (Tokyo)*, **52**, 428-36.
- Rajaram, S., Baylink, D.J. & Mohan, S. (1997). Insulin-like growth factor-binding proteins in serum and other biological fluids: regulation and functions. *Endocr Rev*, **18**, 801-31.
- Ranke, M.B., Savage, M.O., Chatelain, P.G., Preece, M.A., Rosenfeld, R.G. & Wilton, P. (1999). Long-term treatment of growth hormone insensitivity syndrome with IGF-I. Results of the European Multicentre Study. The Working Group on Growth Hormone Insensitivity Syndromes. *Horm Res*, **51**, 128-34.
- Renahan, A.G., Harvie, M. & Howell, A. (2006). Insulin-like growth factor (IGF)-I, IGF binding protein-3, and breast cancer risk: eight years on. *Endocr Relat Cancer*, **13**, 273-8.
- Renahan, A.G., Zwahlen, M., Minder, C., O'Dwyer, S.T., Shalet, S.M. & Egger, M. (2004). Insulin-like growth factor (IGF)-I, IGF binding protein-3, and cancer risk: systematic review and meta-regression analysis. *Lancet*, **363**, 1346-53.
- Rinaldi, S., Kaaks, R., Zeleniuch-Jacquotte, A., Arslan, A.A., Shore, R.E., Koenig, K.L., Dossus, L., Riboli, E., Stattin, P., Lukanova, A. & Toniolo, P. (2005a). Insulin-like growth factor-I, IGF binding protein-3, and breast cancer in young women: a comparison of risk estimates using different peptide assays. *Cancer Epidemiol Biomarkers Prev*, **14**, 48-52.
- Rinaldi, S., Peeters, P.H., Berrino, F., Dossus, L., Biessy, C., Olsen, A., Tjonneland, A., Overvad, K., Clavel-Chapelon, F., Boutron-Ruault, M.C., Tehard, B., Nagel, G., Linseisen, J., Boeing, H., Lahmann, P.H., Trichopoulou, A., Trichopoulos, D., Koliva, M., Palli, D., Panico, S., Tumino, R., Sacerdote, C., van Gils, C.H., van Noord, P., Grobbee, D.E., Bueno-de-Mesquita, H.B., Gonzalez, C.A., Agudo, A., Chirlaque, M.D., Barricarte, A., Larranaga, N., Quiros, J.R., Bingham, S., Khaw, K.T., Key, T., Allen, N.E., Lukanova, A., Slimani, N., Saracci, R., Riboli, E. & Kaaks, R. (2006). IGF-I, IGFBP-3 and breast cancer risk in women: The European Prospective Investigation into Cancer and Nutrition (EPIC). *Endocr Relat Cancer*, **13**, 593-605.
- Rinaldi, S., Toniolo, P., Muti, P., Lundin, E., Zeleniuch-Jacquotte, A., Arslan, A., Micheli, A., Lenner, P., Dossus, L., Krogh, V., Shore, R.E., Koenig, K.L., Riboli, E., Stattin, P., Berrino, F., Hallmans, G., Lukanova, A. & Kaaks, R. (2005b). IGF-I, IGFBP-3 and breast cancer in young women: a pooled re-analysis of three prospective studies. *Eur J Cancer Prev*, **14**, 493-6.
- Rollison, D.E., Newschaffer, C.J., Tao, Y., Pollak, M. & Helzlsouer, K.J. (2006). Premenopausal levels of circulating insulin-like growth factor I and the risk of postmenopausal breast cancer. *Int J Cancer*, **118**, 1279-84.
- Rowland, I., Faughnan, M., Hoey, L., Wahala, K., Williamson, G. & Cassidy, A. (2003). Bioavailability of phyto-oestrogens. *Br J Nutr*, **89 Suppl 1**, S45-58.

- Rowland, I.R., Wiseman, H., Sanders, T.A., Adlercreutz, H. & Bowey, E.A. (2000). Interindividual variation in metabolism of soy isoflavones and lignans: influence of habitual diet on equol production by the gut microflora. *Nutr Cancer*, **36**, 27-32.
- Ruan, W., Catanese, V., Wieczorek, R., Feldman, M. & Kleinberg, D.L. (1995). Estradiol enhances the stimulatory effect of insulin-like growth factor-I (IGF-I) on mammary development and growth hormone-induced IGF-I messenger ribonucleic acid. *Endocrinology*, **136**, 1296-302.
- Ruan, W. & Kleinberg, D.L. (1999). Insulin-like growth factor I is essential for terminal end bud formation and ductal morphogenesis during mammary development. *Endocrinology*, **140**, 5075-81.
- Ruiz-Larrea, M.B., Mohan, A.R., Paganga, G., Miller, N.J., Bolwell, G.P. & Rice-Evans, C.A. (1997). Antioxidant activity of phytoestrogenic isoflavones. *Free Radic Res*, **26**, 63-70.
- Rutanen, E.M. & Pekonen, F. (1991). Assays for IGF binding proteins. *Acta Endocrinol (Copenh)*, **124 Suppl 2**, 70-3.
- Samman, S., Lyons Wall, P.M., Chan, G.S., Smith, S.J. & Petocz, P. (1999). The effect of supplementation with isoflavones on plasma lipids and oxidisability of low density lipoprotein in premenopausal women. *Atherosclerosis*, **147**, 277-83.
- Santell, R.C., Kieu, N. & Helferich, W.G. (2000). Genistein inhibits growth of estrogen-independent human breast cancer cells in culture but not in athymic mice. *J Nutr*, **130**, 1665-9.
- Sarkar, F.H. & Li, Y. (2003). Soy isoflavones and cancer prevention. *Cancer Invest*, **21**, 744-57.
- Sartippour, M.R., Rao, J.Y., Apple, S., Wu, D., Henning, S., Wang, H., Elashoff, R., Rubio, R., Heber, D. & Brooks, M.N. (2004). A pilot clinical study of short-term isoflavone supplements in breast cancer patients. *Nutr Cancer*, **49**, 59-65.
- Sasano, N., Tamahashi, N., Namiki, T. & Stemmermann, G.N. (1975). Consecutive radiography of breast slices for estimation of glandular volume and detection of small subclinical lesions. A comparison between Japan and Hawaii Japanese. *Tohoku J Exp.Med.*, **117**, 217-224.
- Sathyamoorthy, N. & Wang, T.T. (1997). Differential effects of dietary phytoestrogens daidzein and equol on human breast cancer MCF-7 cells. *Eur J Cancer*, **33**, 2384-9.
- Sathyamoorthy, N., Wang, T.T. & Phang, J.M. (1994). Stimulation of pS2 expression by diet-derived compounds. *Cancer Res*, **54**, 957-61.
- Schernhammer, E.S., Holly, J.M., Hunter, D.J., Pollak, M.N. & Hankinson, S.E. (2006). Insulin-like growth factor-I, its binding proteins (IGFBP-1 and IGFBP-3), and growth hormone and breast cancer risk in The Nurses Health Study II. *Endocr Relat Cancer*, **13**, 583-92.
- Schernhammer, E.S., Holly, J.M., Pollak, M.N. & Hankinson, S.E. (2005). Circulating levels of insulin-like growth factors, their binding proteins, and breast cancer risk. *Cancer Epidemiol Biomarkers Prev*, **14**, 699-704.
- Schmitt, E., Dekant, W. & Stopper, H. (2001). Assaying the estrogenicity of phytoestrogens in cells of different estrogen sensitive tissues. *Toxicol In Vitro*, **15**, 433-9.
- Scholar, E.M. & Toews, M.L. (1994). Inhibition of invasion of murine mammary carcinoma cells by the tyrosine kinase inhibitor genistein. *Cancer Lett*, **87**, 159-62.

- Schuerch, C., 3rd, Rosen, P.P., Hirota, T., Itabashi, M., Yamamoto, H., Kinne, D.W. & Beattie, E.J., Jr. (1982). A pathologic study of benign breast diseases in Tokyo and New York. *Cancer*, **50**, 1899-903.
- Schult, T.M., Ensrud, K.E., Blackwell, T., Ettinger, B., Wallace, R. & Tice, J.A. (2004). Effect of isoflavones on lipids and bone turnover markers in menopausal women. *Maturitas*, **48**, 209-18.
- Setchell, K.D., Brown, N.M., Desai, P.B., Zimmer-Nechimias, L., Wolfe, B., Jakate, A.S., Creutzinger, V. & Heubi, J.E. (2003). Bioavailability, disposition, and dose-response effects of soy isoflavones when consumed by healthy women at physiologically typical dietary intakes. *J Nutr*, **133**, 1027-35.
- Setchell, K.D., Brown, N.M. & Lydeking-Olsen, E. (2002). The clinical importance of the metabolite equol—a clue to the effectiveness of soy and its isoflavones. *J Nutr*, **132**, 3577-84.
- Setchell, K.D., Clerici, C., Lephart, E.D., Cole, S.J., Heenan, C., Castellani, D., Wolfe, B.E., Nechemias-Zimmer, L., Brown, N.M., Lund, T.D., Handa, R.J. & Heubi, J.E. (2005). S-equol, a potent ligand for estrogen receptor beta, is the exclusive enantiomeric form of the soy isoflavone metabolite produced by human intestinal bacterial flora. *Am J Clin Nutr*, **81**, 1072-9.
- Shao, Z.M., Alpaugh, M.L., Fontana, J.A. & Barsky, S.H. (1998a). Genistein inhibits proliferation similarly in estrogen receptor-positive and negative human breast carcinoma cell lines characterized by P21WAF1/CIP1 induction, G2/M arrest, and apoptosis. *J Cell Biochem*, **69**, 44-54.
- Shao, Z.M., Wu, J., Shen, Z.Z. & Barsky, S.H. (1998b). Genistein exerts multiple suppressive effects on human breast carcinoma cells. *Cancer Res*, **58**, 4851-7.
- Shen, F., Xue, X. & Weber, G. (1999). Tamoxifen and genistein synergistically down-regulate signal transduction and proliferation in estrogen receptor-negative human breast carcinoma MDA-MB-435 cells. *Anticancer Res*, **19**, 1657-62.
- Shi, R., Yu, H., McLarty, J. & Glass, J. (2004). IGF-I and breast cancer: a meta-analysis. *Int J Cancer*, **111**, 418-23.
- Shimizu, H., Ross, R.K., Bernstein, L., Yatani, R., Henderson, B.E. & Mack, T.M. (1991). Cancers of the prostate and breast among Japanese and white immigrants in Los Angeles County. *Br J Cancer*, **63**, 963-6.
- Shoff, S.M., Newcomb, P.A., Mares-Perlman, J.A., Klein, B.E., Haffner, S.M., Storer, B.E. & Klein, R. (1998). Usual consumption of plant foods containing phytoestrogens and sex hormone levels in postmenopausal women in Wisconsin. *Nutr Cancer*, **30**, 207-12.
- Shu, X.O., Jin, F., Dai, Q., Wen, W., Potter, J.D., Kushi, L.H., Ruan, Z., Gao, Y.T. & Zheng, W. (2001). Soyfood intake during adolescence and subsequent risk of breast cancer among Chinese women. *Cancer Epidemiol Biomarkers Prev*, **10**, 483-8.
- Sierens, J., Hartley, J.A., Campbell, M.J., Leathem, A.J. & Woodside, J.V. (2001). Effect of phytoestrogen and antioxidant supplementation on oxidative DNA damage assessed using the comet assay. *Mutat Res*, **485**, 169-76.
- Sierens, J., Hartley, J.A., Campbell, M.J., Leathem, A.J. & Woodside, J.V. (2002). In vitro isoflavone supplementation reduces hydrogen peroxide-induced DNA damage in sperm. *Teratog Carcinog Mutagen*, **22**, 227-34.

- Simons, L.A., von Konigsmark, M., Simons, J. & Celermajer, D.S. (2000). Phytoestrogens do not influence lipoprotein levels or endothelial function in healthy, postmenopausal women. *Am J Cardiol*, **85**, 1297-301.
- Smith, W.J., Underwood, L.E. & Clemmons, D.R. (1995). Effects of caloric or protein restriction on insulin-like growth factor-I (IGF-I) and IGF-binding proteins in children and adults. *J Clin Endocrinol Metab*, **80**, 443-9.
- So, F.V., Guthrie, N., Chambers, A.F. & Carroll, K.K. (1997). Inhibition of proliferation of estrogen receptor-positive MCF-7 human breast cancer cells by flavonoids in the presence and absence of excess estrogen. *Cancer Lett*, **112**, 127-33.
- Society, A.C. (2006). Cancer Facts and Figures 2006 pp. 35. American Cancer Society: Atlanta.
- Somekawa, Y., Chiguchi, M., Ishibashi, T. & Aso, T. (2001). Soy intake related to menopausal symptoms, serum lipids, and bone mineral density in postmenopausal Japanese women. *Obstet Gynecol*, **97**, 109-15.
- Spanswick, V., Hartley, J., Ward, T. & Hartley, J. (1999). Measurement of drug-induced DNA interstrand crosslinking using the single cell gel electrophoresis (comet) assay. In *Methods in molecular medicine: cytotoxic drug resistance mechanisms*, In: Brown R, B.U., editors. (ed) pp. 143-54. Humana Press: USA7.
- Speirs, V. & Kerin, M.J. (2000). Prognostic significance of oestrogen receptor beta in breast cancer. *Br J Surg*, **87**, 405-9.
- Spentzos, D., Mantzoros, C., Regan, M.M., Morrissey, M.E., Duggan, S., Flickner-Garvey, S., McCormick, H., DeWolf, W., Balk, S. & Bubley, G.J. (2003). Minimal effect of a low-fat/high soy diet for asymptomatic, hormonally naive prostate cancer patients. *Clin Cancer Res*, **9**, 3282-7.
- Stewart, A.J., Johnson, M.D., May, F.E. & Westley, B.R. (1990). Role of insulin-like growth factors and the type I insulin-like growth factor receptor in the estrogen-stimulated proliferation of human breast cancer cells. *J Biol Chem*, **265**, 21172-8.
- Strom, A., Hartman, J., Foster, J.S., Kietz, S., Wimalasena, J. & Gustafsson, J.A. (2004). Estrogen receptor beta inhibits 17beta-estradiol-stimulated proliferation of the breast cancer cell line T47D. *Proc Natl Acad Sci U S A*, **101**, 1566-71.
- Su, S.J., Yeh, T.M., Chuang, W.J., Ho, C.L., Chang, K.L., Cheng, H.L., Liu, H.S., Hsu, P.Y. & Chow, N.H. (2005). The novel targets for anti-angiogenesis of genistein on human cancer cells. *Biochem Pharmacol*, **69**, 307-18.
- Sugumar, A., Liu, Y.C., Xia, Q., Koh, Y.S. & Matsuo, K. (2004). Insulin-like growth factor (IGF)-I and IGF-binding protein 3 and the risk of premenopausal breast cancer: a meta-analysis of literature. *Int J Cancer*, **111**, 293-7.
- Surmacz, E. & Bartucci, M. (2004). Role of estrogen receptor alpha in modulating IGF-I receptor signaling and function in breast cancer. *J Exp Clin Cancer Res*, **23**, 385-94.
- Swan, G. (2004). Findings from the latest National Diet and Nutrition Survey. *Proc Nutr Soc*, **63**, 505-12.
- Takatsuka, N., Nagata, C., Kurisu, Y., Inaba, S., Kawakami, N. & Shimizu, H. (2000). Hypocholesterolemic effect of soymilk supplementation with usual diet in premenopausal normolipidemic Japanese women. *Prev Med*, **31**, 308-14.

- Tannenbaum, G.S., Gurd, W., Lapointe, M. & Pollak, M. (1992). Tamoxifen attenuates pulsatile growth hormone secretion: mediation in part by somatostatin. *Endocrinology*, **130**, 3395-401.
- Tanos, V., Brzezinski, A., Drize, O., Strauss, N. & Peretz, T. (2002). Synergistic inhibitory effects of genistein and tamoxifen on human dysplastic and malignant epithelial breast cells in vitro. *Eur J Obstet Gynecol Reprod Biol*, **102**, 188-94.
- Taylor, C.M., Blanchard, B. & Zava, D.T. (1984). Estrogen receptor-mediated and cytotoxic effects of the antiestrogens tamoxifen and 4-hydroxytamoxifen. *Cancer Res*, **44**, 1409-14.
- Taylor, E.F., Burley, V.J., Greenwood, D.C. & Cade, J.E. (2007). Meat consumption and risk of breast cancer in the UK Women's Cohort Study. *Br J Cancer*, **96**, 1139-46.
- Tew, B.Y., Xu, X., Wang, H.J., Murphy, P.A. & Hendrich, S. (1996). A diet high in wheat fiber decreases the bioavailability of soybean isoflavones in a single meal fed to women. *J Nutr*, **126**, 871-7.
- Thanos, J., Cotterchio, M., Boucher, B.A., Kreiger, N. & Thompson, L.U. (2006). Adolescent dietary phytoestrogen intake and breast cancer risk (Canada). *Cancer Causes Control*, **17**, 1253-61.
- Thierry van Dessel, H.J., Chandrasekhar, Y., Yap, O.W., Lee, P.D., Hintz, R.L., Faessen, G.H., Braat, D.D., Fauser, B.C. & Giudice, L.C. (1996). Serum and follicular fluid levels of insulin-like growth factor I (IGF-I), IGF-II, and IGF-binding protein-1 and -3 during the normal menstrual cycle. *J Clin Endocrinol Metab*, **81**, 1224-31.
- This, P., De La Rochefordiere, A., Clough, K., Fourquet, A. & Magdelenat, H. (2001). Phytoestrogens after breast cancer. *Endocr Relat Cancer*, **8**, 129-34.
- Thissen, J.P., Ketelslegers, J.M. & Underwood, L.E. (1994). Nutritional regulation of the insulin-like growth factors. *Endocr Rev*, **15**, 80-101.
- Thomas, H.V., Davey, G.K. & Key, T.J. (1999). Oestradiol and sex hormone-binding globulin in premenopausal and post-menopausal meat-eaters, vegetarians and vegans. *Br J Cancer*, **80**, 1470-5.
- Thomas, H.V., Reeves, G.K. & Key, T.J. (1997). Endogenous estrogen and postmenopausal breast cancer: a quantitative review. *Cancer Causes Control*, **8**, 922-8.
- Thomas, T., Trend, B., Butterfield, J.R., Janne, O.A. & Kiang, D.T. (1989). Regulation of ornithine decarboxylase gene expression in MCF-7 breast cancer cells by antiestrogens. *Cancer Res*, **49**, 5852-7.
- Thompson, L.P., K. Saarinen, N. Chen, J. and Ward, W. (2005). The Combination of Soy and Flaxseed or Their Phytoestrogens Can Better Reduce the Growth of Breast Tumors than Soy or Genistein Alone While Causing Little Effects on Bone Health in Ovariectomized Nude Mice. (abstract). In *6th International symposium on the role of soy in preventing and treating chronic disease* pp. 5: Chicago, IL.
- Thomsen, A.R., Mortensen, A., Breinholt, V.M., Lindecrona, R.H., Penalvo, J.L. & Sorensen, I.K. (2005). Influence of Prevastein, an isoflavone-rich soy product, on mammary gland development and tumorigenesis in Tg.NK (MMTV/c-neu) mice. *Nutr Cancer*, **52**, 176-88.
- Thorburn, J. & Thorburn, A. (1994). The tyrosine kinase inhibitor, genistein, prevents alpha-adrenergic-induced cardiac muscle cell hypertrophy by inhibiting activation of the Ras-MAP kinase signaling pathway. *Biochem Biophys Res Commun*, **202**, 1586-91.

- Thurnham, D.I., Smith, E. & Flora, P.S. (1988). Concurrent liquid-chromatographic assay of retinol, alpha-tocopherol, beta-carotene, alpha-carotene, lycopene, and beta-cryptoxanthin in plasma, with tocopherol acetate as internal standard. *Clin Chem*, **34**, 377-81.
- Tikkanen, M.J., Wahala, K., Ojala, S., Vihma, V. & Adlercreutz, H. (1998). Effect of soybean phytoestrogen intake on low density lipoprotein oxidation resistance. *Proc Natl Acad Sci U S A*, **95**, 3106-10.
- Toms, J.R. (2004). CancerStats Monograph 2004. Cancer Research UK 2004.: London.
- Tong, D., Schuster, E., Seifert, M., Czerwenka, K., Leodolte, S. & Zeillinger, R. (2002). Expression of estrogen receptor beta isoforms in human breast cancer tissues and cell lines. *Breast Cancer Res Treat*, **71**, 249-55.
- Toniolo, P., Bruning, P.F., Akhmedkhanov, A., Bonfrer, J.M., Koenig, K.L., Lukanova, A., Shore, R.E. & Zeleniuch-Jacquotte, A. (2000). Serum insulin-like growth factor-I and breast cancer. *Int J Cancer*, **88**, 828-32.
- Touillaud, M.S., Thiebaut, A.C., Fournier, A., Niravong, M., Boutron-Ruault, M.C. & Clavel-Chapelon, F. (2007). Dietary lignan intake and postmenopausal breast cancer risk by estrogen and progesterone receptor status. *J Natl Cancer Inst*, **99**, 475-86.
- Treloar, A.E., Boynton, R.E., Behn, B.G. & Brown, B.W. (1967). Variation of the human menstrual cycle through reproductive life. *Int J Fertil*, **12**, 77-126.
- Trock, B.J., Hilakivi-Clarke, L. & Clarke, R. (2006). Meta-analysis of soy intake and breast cancer risk. *J Natl Cancer Inst*, **98**, 459-71.
- Ueda, M., Niho, N., Imai, T., Shibutani, M., Mitsumori, K., Matsui, T. & Hirose, M. (2003). Lack of significant effects of genistein on the progression of 7,12-dimethylbenz(a)anthracene-induced mammary tumors in ovariectomized Sprague-Dawley rats. *Nutr Cancer*, **47**, 141-7.
- Ueno, T.U., S. (2001). Identification of the specific intestinal bacteria capable of metabolising soy isoflavone to equol. *Ann. Nutr. Metab.*, **45**, 114(abs.).
- van Gils, C.H., Peeters, P.H., Bueno-de-Mesquita, H.B., Boshuizen, H.C., Lahmann, P.H., Clavel-Chapelon, F., Thiebaut, A., Kesse, E., Sieri, S., Palli, D., Tumino, R., Panico, S., Vineis, P., Gonzalez, C.A., Ardanaz, E., Sanchez, M.J., Amiano, P., Navarro, C., Quiros, J.R., Key, T.J., Allen, N., Khaw, K.T., Bingham, S.A., Psaltopoulou, T., Koliva, M., Trichopoulou, A., Nagel, G., Linseisen, J., Boeing, H., Berglund, G., Wirfalt, E., Hallmans, G., Lenner, P., Overvad, K., Tjonneland, A., Olsen, A., Lund, E., Engeset, D., Alsaker, E., Norat, T., Kaaks, R., Slimani, N. & Riboli, E. (2005). Consumption of vegetables and fruits and risk of breast cancer. *Jama*, **293**, 183-93.
- Van Wyk, J.J., Svoboda, M.E. & Underwood, L.E. (1980). Evidence from radioligand assays that somatomedin-C and insulin-like growth factor-I are similar to each other and different from other somatomedins. *J Clin Endocrinol Metab*, **50**, 206-8.
- Vedrine, N., Mathey, J., Morand, C., Brandolini, M., Davicco, M.J., Guy, L., Remesy, C., Coxam, V. & Manach, C. (2006). One-month exposure to soy isoflavones did not induce the ability to produce equol in postmenopausal women. *Eur J Clin Nutr*, **60**, 1039-45.
- Verheus, M., van Gils, C.H., Keinan-Boker, L., Grace, P.B., Bingham, S.A. & Peeters, P.H. (2007). Plasma phytoestrogens and subsequent breast cancer risk. *J Clin Oncol*, **25**, 648-55.

- Verkasalo, P.K., Appleby, P.N., Davey, G.K. & Key, T.J. (2001). Soy milk intake and plasma sex hormones: a cross-sectional study in pre- and postmenopausal women (EPIC-Oxford). *Nutr Cancer*, **40**, 79-86.
- Veronesi, U., Maisonneuve, P., Rotmensz, N., Costa, A., Sacchini, V., Travaglini, R., D'Aiuto, G., Lovison, F., Gucciardo, G., Muraca, M.G., Pizzichetta, M.A., Conforti, S., Decensi, A., Robertson, C. & Boyle, P. (2003). Italian randomized trial among women with hysterectomy: tamoxifen and hormone-dependent breast cancer in high-risk women. *J Natl Cancer Inst*, **95**, 160-5.
- Vladusic, E.A., Hornby, A.E., Guerra-Vladusic, F.K., Lakins, J. & Lupu, R. (2000). Expression and regulation of estrogen receptor beta in human breast tumors and cell lines. *Oncol Rep*, **7**, 157-67.
- Vogel, V.G., Costantino, J.P., Wickerham, D.L., Cronin, W.M., Cecchini, R.S., Atkins, J.N., Bevers, T.B., Fehrenbacher, L., Pajon, E.R., Jr., Wade, J.L., 3rd, Robidoux, A., Margolese, R.G., James, J., Lippman, S.M., Runowicz, C.D., Ganz, P.A., Reis, S.E., McCaskill-Stevens, W., Ford, L.G., Jordan, V.C. & Wolmark, N. (2006). Effects of tamoxifen vs raloxifene on the risk of developing invasive breast cancer and other disease outcomes: the NSABP Study of Tamoxifen and Raloxifene (STAR) P-2 trial. *Jama*, **295**, 2727-41.
- Voskuil, D.W., Bosma, A., Vrieling, A., Rookus, M.A. & van 't Veer, L.J. (2004). Insulin-like growth factor (IGF)-system mRNA quantities in normal and tumor breast tissue of women with sporadic and familial breast cancer risk. *Breast Cancer Res Treat*, **84**, 225-33.
- Voskuil, D.W., Vrieling, A., van't Veer, L.J., Kampman, E. & Rookus, M.A. (2005). The insulin-like growth factor system in cancer prevention: potential of dietary intervention strategies. *Cancer Epidemiol Biomarkers Prev*, **14**, 195-203.
- Vrieling, A., Rookus, M.A., Kampman, E., Bonfrer, J.M., Korse, C.M., van Doorn, J., Lampe, J.W., Cats, A., Witteman, B.J., van Leeuwen, F.E., van't Veer, L.J. & Voskuil, D.W. (2007). Isolated isoflavones do not affect the circulating insulin-like growth factor system in men at increased colorectal cancer risk. *J Nutr*, **137**, 379-83.
- Vrieling, A., Voskuil, D.W., Bueno de Mesquita, H.B., Kaaks, R., van Noord, P.A., Keinan-Boker, L., van Gils, C.H. & Peeters, P.H. (2004). Dietary determinants of circulating insulin-like growth factor (IGF)-I and IGF binding proteins 1, -2 and -3 in women in the Netherlands. *Cancer Causes Control*, **15**, 787-96.
- Vuilleumier, J. & Keck, E. (1989). Fluorometric assay of vitamin C in biological materials using a centrifugal analyser with fluorescence attachment. *J Micronutr Anal*, **5**, 25-34.
- Wang, C. & Kurzer, M.S. (1997). Phytoestrogen concentration determines effects on DNA synthesis in human breast cancer cells. *Nutr Cancer*, **28**, 236-47.
- Wang, C. & Kurzer, M.S. (1998). Effects of phytoestrogens on DNA synthesis in MCF-7 cells in the presence of estradiol or growth factors. *Nutr Cancer*, **31**, 90-100.
- Wang, H.S., Lee, J.D. & Soong, Y.K. (1995). Serum levels of insulin-like growth factor I and insulin-like growth factor-binding protein-1 and -3 in women with regular menstrual cycles. *Fertil Steril*, **63**, 1204-9.

- Wang, T.T., Sathyamoorthy, N. & Phang, J.M. (1996). Molecular effects of genistein on estrogen receptor mediated pathways. *Carcinogenesis*, **17**, 271-5.
- Wangen, K.E., Duncan, A.M., Merz-Demlow, B.E., Xu, X., Marcus, R., Phipps, W.R. & Kurzer, M.S. (2000). Effects of soy isoflavones on markers of bone turnover in premenopausal and postmenopausal women. *J Clin Endocrinol Metab*, **85**, 3043-8.
- Wei, H., Cai, Q., Tian, L. & Lebwohl, M. (1998). Tamoxifen reduces endogenous and UV light-induced oxidative damage to DNA, lipid and protein in vitro and in vivo. *Carcinogenesis*, **19**, 1013-8.
- Wei, H., Wei, L., Frenkel, K., Bowen, R. & Barnes, S. (1993). Inhibition of tumor promoter-induced hydrogen peroxide formation in vitro and in vivo by genistein. *Nutr Cancer*, **20**, 1-12.
- Westley, B.R. & May, F.E. (1994). Role of insulin-like growth factors in steroid modulated proliferation. *J Steroid Biochem Mol Biol*, **51**, 1-9.
- Winston, R., Kao, P.C. & Kiang, D.T. (1994). Regulation of insulin-like growth factors by antiestrogen. *Breast Cancer Res Treat*, **31**, 107-15.
- Wiseman, H., Casey, K., Bowey, E.A., Duffy, R., Davies, M., Rowland, I.R., Lloyd, A.S., Murray, A., Thompson, R. & Clarke, D.B. (2004). Influence of 10 wk of soy consumption on plasma concentrations and excretion of isoflavonoids and on gut microflora metabolism in healthy adults. *Am J Clin Nutr*, **80**, 692-9.
- Wiseman, H., O'Reilly, J.D., Adlercreutz, H., Mallet, A.I., Bowey, E.A., Rowland, I.R. & Sanders, T.A. (2000). Isoflavone phytoestrogens consumed in soy decrease F(2)-isoprostane concentrations and increase resistance of low-density lipoprotein to oxidation in humans. *Am J Clin Nutr*, **72**, 395-400.
- Witte, J.S., Ursin, G., Siemiatycki, J., Thompson, W.D., Paganini-Hill, A. & Haile, R.W. (1997). Diet and premenopausal bilateral breast cancer: a case-control study. *Breast Cancer Res Treat*, **42**, 243-51.
- Wolff, S. (1994). Ferrous ion oxidation in presence of ferric ion indicator xylenol orange for measurement of hydroperoxides. *Methods Enzymol*, **233**, 182-189.
- Woodside, J.V. & Campbell, M.J. (2001). Isoflavones and breast cancer. *J Br Menopause Soc*, **7**, S17-S21.
- Woodside, J.V., Campbell, M.J., Denholm, E.E., Newton, L., Honour, J.W., Morton, M.S., Young, I.S. & Leathem, A.J. (2006). Short-term phytoestrogen supplementation alters insulin-like growth factor profile but not lipid or antioxidant status. *J Nutr Biochem*, **17**, 211-5.
- Wosikowski, K., Kung, W., Hasmann, M., Loser, R. & Eppenberger, U. (1993). Inhibition of growth-factor-activated proliferation by anti-estrogens and effects on early gene expression of MCF-7 cells. *Int J Cancer*, **53**, 290-7.
- Wu, A.H., Stanczyk, F.Z., Hendrich, S., Murphy, P.A., Zhang, C., Wan, P. & Pike, M.C. (2000). Effects of soy foods on ovarian function in premenopausal women. *Br J Cancer*, **82**, 1879-86.
- Wu, A.H., Wan, P., Hankin, J., Tseng, C.C., Yu, M.C. & Pike, M.C. (2002). Adolescent and adult soy intake and risk of breast cancer in Asian-Americans. *Carcinogenesis*, **23**, 1491-6.
- Wu, A.H., Ziegler, R.G., Horn-Ross, P.L., Nomura, A.M., West, D.W., Kolonel, L.N., Rosenthal, J.F., Hoover, R.N. & Pike, M.C. (1996). Tofu and risk of breast cancer in Asian-Americans. *Cancer Epidemiol Biomarkers Prev*, **5**, 901-6.

- Wu, A.H., Ziegler, R.G., Nomura, A.M. & et, a. (1998). Soy intake and risk of breast cancer in Asians and Asian Americans. *Am.J Clin.Nutr.*, **68**, 1437S-1443S.
- Wu, J., Oka, J., Higuchi, M., Tabata, I., Toda, T., Fujioka, M., Fuku, N., Teramoto, T., Okuhira, T., Ueno, T., Uchiyama, S., Urata, K., Yamada, K. & Ishimi, Y. (2006). Cooperative effects of isoflavones and exercise on bone and lipid metabolism in postmenopausal Japanese women: a randomized placebo-controlled trial. *Metabolism*, **55**, 423-33.
- Wu, Y., Cui, K., Miyoshi, K., Hennighausen, L., Green, J.E., Setser, J., LeRoith, D. & Yakar, S. (2003). Reduced circulating insulin-like growth factor I levels delay the onset of chemically and genetically induced mammary tumors. *Cancer Res*, **63**, 4384-8.
- Xu, X., Wang, H.J., Murphy, P.A., Cook, L. & Hendrich, S. (1994). Daidzein is a more bioavailable soymilk isoflavone than is genistein in adult women. *J Nutr*, **124**, 825-32.
- Yakar, S., Pennisi, P., Kim, C.H., Zhao, H., Toyoshima, Y., Gavrilova, O. & LeRoith, D. (2005). Studies involving the GH-IGF axis: Lessons from IGF-I and IGF-I receptor gene targeting mouse models. *J Endocrinol Invest*, **28**, 19-22.
- Yamamoto, S., Sobue, T., Kobayashi, M., Sasaki, S. & Tsugane, S. (2003). Soy, isoflavones, and breast cancer risk in Japan. *J Natl Cancer Inst*, **95**, 906-13.
- Yan, L., Li, D. & Yee, J.A. (2002). Dietary supplementation with isolated soy protein reduces metastasis of mammary carcinoma cells in mice. *Clin Exp Metastasis*, **19**, 535-40.
- Young, I.S. & Trimble, E.R. (1991). Measurement of malondialdehyde in plasma by high performance liquid chromatography with fluorimetric detection. *Ann Clin Biochem*, **28 (Pt 5)**, 504-8.
- Yu, H., Jin, F., Shu, X.O., Li, B.D., Dai, Q., Cheng, J.R., Berkel, H.J. & Zheng, W. (2002). Insulin-like growth factors and breast cancer risk in Chinese women. *Cancer Epidemiol Biomarkers Prev*, **11**, 705-12.
- Yuan, J.M., Wang, Q.S., Ross, R.K., Henderson, B.E. & Yu, M.C. (1995). Diet and breast cancer in Shanghai and Tianjin, China. *Br J Cancer*, **71**, 1353-8.
- Zava, D.T. & Duwe, G. (1997). Estrogenic and antiproliferative properties of genistein and other flavonoids in human breast cancer cells in vitro. *Nutr Cancer*, **27**, 31-40.
- Zhang, Y., Song, T.T., Cunnick, J.E., Murphy, P.A. & Hendrich, S. (1999). Daidzein and genistein glucuronides in vitro are weakly estrogenic and activate human natural killer cells at nutritionally relevant concentrations. *J Nutr*, **129**, 399-405.
- Zheng, W., Dai, Q., Custer, L.J., Shu, X.O., Wen, W.Q., Jin, F. & Franke, A.A. (1999). Urinary excretion of isoflavonoids and the risk of breast cancer. *Cancer Epidemiol Biomarkers Prev*, **8**, 35-40.
- Zhou, J.R., Gugger, E.T., Tanaka, T., Guo, Y., Blackburn, G.L. & Clinton, S.K. (1999). Soybean phytochemicals inhibit the growth of transplantable human prostate carcinoma and tumor angiogenesis in mice. *J Nutr*, **129**, 1628-35.
- Zhou, J.R., Yu, L., Mai, Z. & Blackburn, G.L. (2004). Combined inhibition of estrogen-dependent human breast carcinoma by soy and tea bioactive components in mice. *Int J Cancer*, **108**, 8-14.

- Ziegler, R.G., Hoover, R.N., Pike, M.C., Hildesheim, A., Nomura, A.M., West, D.W., Wu-Williams, A.H., Kolonel, L.N., Horn-Ross, P.L., Rosenthal, J.F. & Hyer, M.B. (1993). Migration patterns and breast cancer risk in Asian-American women. *J Natl Cancer Inst*, **85**, 1819-27.
- Zubik, L. & Meydani, M. (2003). Bioavailability of soybean isoflavones from aglycone and glucoside forms in American women. *Am J Clin Nutr*, **77**, 1459-65.